

LIMIT COMPUTABILITY AND CONSTRUCTIVE MEASURE

DENIS R. HIRSCHFELDT AND SEBASTIAAN A. TERWIJN

ABSTRACT. In this paper we study constructive measure and dimension in the class Δ_2^0 of limit computable sets. We prove that the lower cone of any Turing-incomplete set in Δ_2^0 has Δ_2^0 -dimension 0, and in contrast, that although the upper cone of a noncomputable set in Δ_2^0 always has Δ_2^0 -measure 0, upper cones in Δ_2^0 have nonzero Δ_2^0 -dimension. In particular the Δ_2^0 -dimension of the Turing degree of \emptyset' (the Halting Problem) is 1. Finally, it is proved that the low sets do not have Δ_2^0 -measure 0, which means that the low sets do not form a small subset of Δ_2^0 . This result has consequences for the existence of bi-immune sets.

1. INTRODUCTION

In his study of randomness [27], Schnorr introduced the notion of a Schnorr null set as a more constructive version of Martin-Löf's [21] notion of null set. We briefly review the relevant definitions. For motivation and discussion of these notions we refer the reader to Schnorr's book [27], the monograph by Li and Vitányi [17], and the recent surveys [5, 8, 31].

For $\sigma \in 2^{<\omega}$ and $X \in 2^\omega$, we write $\sigma \sqsubset X$ to mean that σ is an initial segment of X . A set $\mathcal{A} \subseteq 2^\omega$ is a Σ_1^0 -class if there is a c.e. set $A \subseteq 2^{<\omega}$ such that $\mathcal{A} = \bigcup_{\sigma \in A} [\sigma]$, where $[\sigma] = \{X \in 2^\omega : \sigma \sqsubset X\}$. Whenever we mention a Σ_1^0 -class \mathcal{A} , we assume we have fixed such a set of generators A , and identify \mathcal{A} with A . Note that we can assume that A is *prefix-free*, that is, if $\sigma \in A$ and $\sigma \prec \tau$ then $\tau \notin A$.

Let μ be the usual Lebesgue measure on 2^ω . A set $\mathcal{A} \subseteq 2^\omega$ is called *Martin-Löf null* (or Σ_1^0 -null) if there is a uniformly c.e. sequence $\{\mathcal{U}_i\}_{i \in \omega}$ of Σ_1^0 -classes (called a *test*) such that $\mu(\mathcal{U}_i) \leq 2^{-i}$ and $\mathcal{A} \subseteq \bigcap_i \mathcal{U}_i$. The set \mathcal{A} is *Schnorr null* if in addition the measures $\mu(\mathcal{U}_i)$ are uniformly computable reals. A test with this extra property is called a *total test* or a *Schnorr test*. Equivalently, \mathcal{A} is Schnorr null if there is a test $\{\mathcal{U}_i\}_{i \in \omega}$ such that $\mu(\mathcal{U}_i) = 2^{-i}$ and $\mathcal{A} \subseteq \bigcap_i \mathcal{U}_i$.

The corresponding randomness notions are defined by saying that $A \in 2^\omega$ is Σ_1^0 -random (or 1-random or *Martin-Löf random*) if $\{A\}$ is not Σ_1^0 -null, and A is *Schnorr random* if $\{A\}$ is not Schnorr null.

A different treatment of measure is the one of Ville [32] using martingales. A *martingale* is a function $d : 2^{<\omega} \rightarrow \mathbb{Q}^+$ that satisfies for every $\sigma \in 2^{<\omega}$ the averaging condition $2d(\sigma) = d(\sigma 0) + d(\sigma 1)$, and d is a *supermartingale* if merely $2d(\sigma) \geq d(\sigma 0) + d(\sigma 1)$. A (super)martingale d *succeeds on* a set A if $\limsup_{n \rightarrow \infty} d(A \upharpoonright n) = \infty$. We say that d *succeeds on*, or *covers*, a class $\mathcal{A} \subseteq 2^\omega$ if d succeeds on every $A \in \mathcal{A}$. The *success set* $S[d]$ of d is the class of all sets on which d succeeds. Ville

The first author was supported by National Science Foundation Grants DMS-02-00465 and DMS-05-00590. The second author was supported by the Austrian Research Fund FWF under grants P17503-N12 and P18713-N12.

proved that the class of null sets of the form $S[d]$, with d of arbitrary complexity, coincides with the class of classical (Lebesgue) null sets.

Schnorr gave characterizations of the above notions of effectively null set in terms of martingales. In particular, he introduced null sets of the form

$$S_h[d] = \left\{ X : \limsup_{n \rightarrow \infty} \frac{d(X \upharpoonright n)}{h(n)} = \infty \right\},$$

where d is a martingale and h is a nondecreasing unbounded function (called an *order*), and proved the following theorem.

Theorem 1.1 (Schnorr [27], Sätze 9.4, 9.5). *A set $\mathcal{A} \subseteq 2^\omega$ is Schnorr null if and only if there are a computable martingale d and a computable order h such that $\mathcal{A} \subseteq S_h[d]$.*

Schnorr also addressed null sets of *exponential* order, that is, of the form $S_h[d]$ with $h(n) = 2^{\varepsilon n}$ and $\varepsilon \in (0, 1]$. Although he did not make an explicit reference to Hausdorff dimension, it turns out that the theory of Hausdorff dimension can be cast precisely in terms of such null sets of exponential order, so that Schnorr's notion of effective measure in a natural way leads us into the theory of dimension.

Lutz [18] used effective martingales to develop his theory of resource bounded measure. He defined $A \in 2^\omega$ to be *computably random* if there is no computable martingale d such that $A \in S[d]$.¹ This framework for studying measure and randomness at the level of complexity classes can be used to constructivize Hausdorff dimension along the same lines:

Definition 1.2. For a complexity class \mathcal{C} , a set $\mathcal{A} \subseteq 2^\omega$ has \mathcal{C} -dimension α if

$$\alpha = \inf \left\{ s \in \mathbb{Q} : \exists d \in \mathcal{C} \left(d \text{ is a supermartingale and } \mathcal{A} \subseteq S_{2^{(1-s)n}}[d] \right) \right\}.$$

Lutz [19, 20] used a variant of martingales called *gales* in his presentation. In this paper we stick to martingales and the null sets of the form $S_h[d]$ used by Schnorr in our treatment of Hausdorff dimension. That this makes no difference was pointed out by several authors, including those of [2, 3, 31]. If \mathcal{C} consists of all functions, then the notion of \mathcal{C} -dimension is equivalent to classical Hausdorff dimension. We say that a function $2^{<\omega} \rightarrow \mathbb{Q}^+$ is in Σ_1^0 if it is approximable from below by a nondecreasing computable function. If $\mathcal{C} = \Sigma_1^0$ then the notion of \mathcal{C} -dimension is equivalent to Lutz's definition [20] of constructive Σ_1^0 -dimension.

In this paper we are interested in the quantitative structure of Δ_2^0 . The appropriate measures to use in this context are those for which Δ_2^0 itself does not have measure 0, but for which every element of Δ_2^0 does have measure 0. Since there are Σ_1^0 -random sets in Δ_2^0 , Martin-Löf's Σ_1^0 -measure is too weak for our purposes. For Σ_2^0 -measure, obtained by relativizing Σ_1^0 -measure to the halting set \emptyset' , the class Δ_2^0 has measure 0, so this measure is too strong. However, relativizing the notions of Schnorr null and computably null to \emptyset' gives measures that meet our requirements:

Definition 1.3. A set $\mathcal{A} \subseteq 2^\omega$ has Δ_2^0 -measure 0 (or is Δ_2^0 -null) if there is a \emptyset' -computable martingale that succeeds on \mathcal{A} .

A set $\mathcal{A} \subseteq 2^\omega$ has *Schnorr Δ_2^0 -measure 0* (or is *Schnorr Δ_2^0 -null*) if there is a \emptyset' -computable Schnorr test that covers \mathcal{A} .

¹Schnorr also considered this definition in relativized form [27, p. 55].

A first study of the quantitative structure of Δ_2^0 using these measures was made in Terwijn [29, 30].

Relativizing computable randomness yields Δ_2^0 -*randomness*, while relativizing Schnorr randomness yields *Schnorr Δ_2^0 -randomness*. The relations between the various notions are as follows:

$$\begin{array}{ccc}
\Delta_2^0\text{-random} & & \\
\Downarrow & & \\
\text{Schnorr } \Delta_2^0\text{-random} & \implies & \Delta_2^0\text{-dimension 1} \\
\Downarrow & & \Downarrow \\
\Sigma_1^0\text{-random} & \implies & \Sigma_1^0\text{-dimension 1} \\
\Downarrow & & \\
\text{computably random} & & \Downarrow \\
\Downarrow & & \\
\text{Schnorr random} & \implies & \text{computable dimension 1}
\end{array}$$

No other implications hold than the ones indicated. That there are Schnorr random sets that are not computably random was proved by Wang [33]. (See Nies, Stephan, and Terwijn [22] for more information on the separation between the various randomness notions.) The strictness of the other implications in the first column follows from elementary observations and results in Schnorr [27], and is discussed in [8, 30]. That there are no more implications between the first and the second column follows from the next proposition. The strictness of the two implications in the second column follows by similar means.²

Proposition 1.4. *There are sets A such that A is not Schnorr random and A has Δ_2^0 -dimension 1.*

Proof. Let R be Δ_2^0 -random, and let $D = \{2^x : x \in \omega\}$ be an exponentially sparse computable domain. Then $A = R \cup D$ is not Schnorr random, since no Schnorr random set contains an infinite computable subset, but no Δ_2^0 -martingale can succeed on A exponentially fast. \square

Clearly, the “ Δ_2^0 -dimension 1” in Proposition 1.4 can be improved to “ Δ_n^0 -dimension 1” by the same proof, if one is considering higher orders of randomness.

The rest of this paper is organized as follows. Ambos-Spies, Merkle, Reimann, and Stephan [2] investigated resource bounded dimension in the exponential time class E. Among other things, they proved that under polynomial time many-one reducibility the complete degree in E has dimension 1, and that the set of possible dimensions of p-m-degrees in E is dense in $[0, 1]$. In Section 2 we show that under Turing reducibility in Δ_2^0 the complete degree has Δ_2^0 -dimension 1, and all other degrees have Δ_2^0 -dimension 0. In Section 3 we present a proof that the low sets do not have Δ_2^0 -measure 0 by showing that for every \emptyset' -computable martingale there is a low set that is not covered by it. This means that the low sets do not form a small subset of Δ_2^0 .

Our notation generally follows Odifreddi [23, 24] and Soare [28]. We write $\leq_{\text{T}} A$ for the lower cone $\{B : B \leq_{\text{T}} A\}$ and $A \leq^{\text{T}}$ for the upper cone $\{B : A \leq_{\text{T}} B\}$.

²It is easy to see (cf. [20]) that the class of computable sets has Σ_1^0 -dimension 0, but is not computably null, so in particular this class has computable dimension 1. Also, Lutz [20] has shown that there are sets in Δ_2^0 of any given rational Σ_1^0 -dimension, but it is obvious that every set in Δ_2^0 has Δ_2^0 -dimension 0.

2. Δ_2^0 -DIMENSION

The next theorem is a strengthening of Theorem 5.5 in [30], which states that the lower cone of every $A <_{\mathbb{T}} \emptyset'$ has Schnorr Δ_2^0 -measure 0. We will make use of the following definition and lemma.

Definition 2.1. For functions f and g and rational $q \in (0, 1]$, we say that f is q -dominated by g if

$$(1) \quad \liminf_{n \rightarrow \infty} \frac{|\{i \leq n : g(i) \geq f(i)\}|}{n} \geq q.$$

Lemma 2.2. For every $q \in (0, 1]$ there is a function $f \leq_{\mathbb{T}} \emptyset'$ such that f is not q -dominated by any function $g <_{\mathbb{T}} \emptyset'$.

Proof. Let $h \leq_{\mathbb{T}} \emptyset'$ be a function not dominated by any function $g <_{\mathbb{T}} \emptyset'$, for example, $h(x) = \mu s(\emptyset'_s \upharpoonright x = \emptyset' \upharpoonright x)$ (the smallest s such that all the $y \in \emptyset'$ smaller than x are enumerated into \emptyset' within s steps). Without loss of generality, $q = \frac{1}{c}$ for some $c \in \omega$. Define $f(x) = h(\lfloor \log_c x \rfloor)$ (where \log_c is the logarithm with base c). If g satisfies (1) then for almost every k there is a natural number $x \in [c^k, c^{k+1})$ such that $g(x) \geq f(x)$. But then the function \hat{g} defined by $\hat{g}(k) = \max\{g(x) : x \in [c^k, c^{k+1})\}$ dominates h , a contradiction. \square

Theorem 2.3. Let $A \in \Delta_2^0$ be any Turing-incomplete set. Then the Δ_2^0 -dimension of the lower cone $\leq^{\mathbb{T}} A$ is 0.

Proof. Let $q \in (0, 1]$ be rational and suppose that $A <_{\mathbb{T}} \emptyset'$. We define uniformly in \emptyset' for every $e \in \omega$ a martingale d_e such that

$$R_e : \quad \Phi_e^A \text{ total and } 0, 1\text{-valued} \implies \Phi_e^A \in S_{2^{(1-q)n}}[d_e].$$

By the usual sum trick this suffices to prove the theorem: The sum $d(\sigma) = \sum_{e \in \omega} 2^{-e} d_e(\sigma)$ is again a \emptyset' -computable martingale, and if $X \in S_{2^{(1-q)n}}[d_e]$ then for all $q' > q$ we have $X \in S_{2^{(1-q')n}}[d]$, which shows that $\{B : B \leq_{\mathbb{T}} A\}$ has Δ_2^0 -dimension $\leq q$. Since $q > 0$ was arbitrary the theorem follows.

By Lemma 2.2, let $f \leq_{\mathbb{T}} \emptyset'$ be a function that is not q -dominated by any function $g <_{\mathbb{T}} \emptyset'$.

We now define d_e in stages s . At stage s we define d_e on all strings $\sigma \in 2^{<\omega}$ of length s . The value $d_e(\sigma)$ will depend only on $|\sigma|$. (Such martingales were called ‘oblivious’ in Ambos-Spies, Mayordomo, Wang, and Zheng [1].)

Stage $s = 0$. Define $d_e(\lambda) = 1$, where λ is the empty string.

Stage $s + 1$. Given $d_e(\sigma)$ with $|\sigma| = s$, use the oracle \emptyset' to search for a string $\tau \sqsubset A$ with $|\tau| \leq f(s)$ such that $\Phi_{e,|\tau|}^\tau(s) \downarrow$. If such τ does not exist, or if $\Phi_{e,|\tau|}^\tau(s) \downarrow \notin \{0, 1\}$, do not make a bet; that is, let $d_e(\sigma i) = d_e(\sigma)$ for $i \in \{0, 1\}$. If τ exists and $\Phi_{e,|\tau|}^\tau(s) \downarrow = i \in \{0, 1\}$, define $d_e(\sigma i) = 2d_e(\sigma)$; that is, bet all our capital on $\Phi_e^A(|\sigma|) = i$. This concludes the definition of d_e .

It is clear that d_e is defined on all strings for every e , uniformly in \emptyset' . We check that R_e is satisfied. Suppose that Φ_e^A is total and computes a set. Then the function

$$g_e(n) = \mu t \left((\exists \tau \sqsubset A) [|\tau| = t \wedge \Phi_{e,t}^\tau(n) \downarrow] \right)$$

is A -computable. By the choice of f , there are infinitely many N such that for more than $(1 - q)N$ many $n < N$ we have $f(n) \geq g_e(n)$. For these n , in the definition of

d_e the string τ is found and a bet is placed successfully. (Note that we never make a wrong bet.) Hence $\Phi_e^A \in S_{2^{(1-q)^n}}[d_e]$. \square

Theorem 2.4. *For every noncomputable $A \in \Delta_2^0$, the upper cone A^{\leq_T} has Schnorr Δ_2^0 -measure 0.*

Proof. This theorem is an effectivization of the well-known result of de Leeuw, Moore, Shannon and Shapiro [16] and Sacks [26] that the upper cone of a noncomputable set has Lebesgue measure 0.³ Lutz and Terwijn [29] showed that there is a \emptyset' -computable martingale that succeeds on A^{\leq_T} when $A \in \Delta_2^0$ is noncomputable. We give here a direct proof using total \emptyset' -computable tests, which gives the stronger result of the theorem. Fix a noncomputable $A \in \Delta_2^0$, and define for every i and n the open sets

$$\mathcal{E}_{i,n} = \{B : A \upharpoonright n = \Phi_i^B \upharpoonright n\}.$$

For every i we have $\{B : A = \Phi_i^B\} = \bigcap_n \mathcal{E}_{i,n}$, so $\mu(\bigcap_n \mathcal{E}_{i,n}) = 0$. Furthermore, the $\mathcal{E}_{i,n}$ are uniformly \emptyset' -computable because A is \emptyset' -computable, and the $\mu(\mathcal{E}_{i,n})$ are uniformly \emptyset' -computable reals. So if we let $f(k)$ be the least n such that $\mu(\mathcal{E}_{i,n}) \leq 2^{-k}$ and define $\mathcal{F}_{i,k} = \mathcal{E}_{i,f(k)}$, then $\mathcal{F}_{i,0}, \mathcal{F}_{i,1}, \dots$ is a total \emptyset' -computable test, and we still have $\{B : A = \Phi_i^B\} = \bigcap_k \mathcal{F}_{i,k}$.

Because the tests $\bigcap_k \mathcal{F}_{i,k}$ are \emptyset' -uniform in i , it follows from an easily proved effective union lemma that $\bigcup_i \bigcap_k \mathcal{F}_{i,k} = A^{\leq_T}$ is also of Schnorr measure 0 relative to \emptyset' . \square

The next theorem shows that for every $A \in \Delta_2^0$ the Δ_2^0 -dimension of the upper cone of A is maximal. In particular, although the Schnorr Δ_2^0 -measure of $\text{deg}_T(\emptyset')$ is 0, there is no Δ_2^0 -martingale that succeeds on this Turing degree exponentially fast.

Theorem 2.5. *The Δ_2^0 -dimension of $\text{deg}_T(\emptyset')$ is 1.*

*Proof.*⁴ Given a martingale $d \in \Delta_2^0$ and a rational $\varepsilon > 0$, we build a $B \equiv_T \emptyset'$ such that $B \notin S_{2^{\varepsilon n}}[d]$. The idea is simple: We code \emptyset' on an exponentially sparse computable domain D , and define B by \emptyset' -effectively diagonalizing against d outside D and taking the coded version of \emptyset' on D . Since D is exponentially sparse, d cannot succeed fast on B , and we have $B \leq_T \emptyset' \leq_T B \oplus D \leq_T B$ since D is computable. Note that this idea works for every computably sparse domain D , so that in fact $\text{deg}_T(\emptyset')$ is not included in any null set of the form $S_h[d]$ for a Δ_2^0 -martingale d and a computable order h . Theorem 2.4 shows that the same is not true for all \emptyset' -computable orders h . For the theorem as stated it suffices to take $D = \{2^m - 1 : m \in \omega\}$ and define

$$B(x) = \begin{cases} \emptyset'(n) & \text{if } x = 2^m - 1 \\ 0 & \text{if } x \notin D \text{ and } d((B \upharpoonright x)0) < d((B \upharpoonright x)1) \\ 1 & \text{otherwise.} \end{cases}$$

³For another approach to effectivizing this result, see Hirschfeldt, Nies, and Stephan [10].

⁴This proof is a few years old. Coding techniques similar to the one used in it have meanwhile been used in the context of Hausdorff dimension independently by several authors, cf. e.g. Reimann [25].

Then $d(B \upharpoonright n - 1) \leq d(B \upharpoonright n)$ except possibly when $n = 2^m - 1$ for some n , so

$$\limsup_{n \rightarrow \infty} \frac{d(B \upharpoonright n)}{2^{\varepsilon n}} \leq \limsup_{m \rightarrow \infty} \frac{d(B \upharpoonright 2^m)}{2^{\varepsilon 2^m}} \leq \limsup_{m \rightarrow \infty} \frac{2^m}{2^{\varepsilon 2^m}} < 1,$$

and hence $B \notin S_{2^{\varepsilon n}}[d]$. \square

It follows from Theorems 2.3 and 2.5 that the only possibilities for the Δ_2^0 -dimension of a Turing degree are 0 or 1:

Corollary 2.6. *For $A \in \Delta_2^0$, the Δ_2^0 -dimension of $\deg_T(A)$ is 1 if A is Turing complete, and 0 otherwise.*

3. THE MEASURE OF THE LOW SETS

It is known that the class of sets that are bounded by a 1-generic set has Σ_1^0 -measure 0 (by effectivizing Theorem 4.2 in Kurtz [14], cf. [30], or by Demuth and Kučera [4]). In particular the subclass of the low sets consisting of the Δ_2^0 1-generic sets has Σ_1^0 -measure 0. In this section we prove that the low sets do not form a small subset of Δ_2^0 , that is, that they do not have Δ_2^0 -measure 0. It is easily verified that the computable sets have Δ_2^0 -measure 0, and that most sets in Δ_2^0 are bi-immune for the computable sets.⁵ Although the low sets do not have Δ_2^0 -measure 0, Downey, Hirschfeldt, Lempp, and Solomon [6] were able to construct a Δ_2^0 set A that is bi-immune for the low sets (i.e. there is no infinite low subset of either A or its complement). The set they constructed in fact truth-table reduces to \emptyset' . It is not difficult to see that the sets that tt-reduce to \emptyset' have Schnorr Δ_2^0 -measure 0, i.e. there is a total \emptyset' -computable test covering them. So in this sense the set constructed in [6] does not exhibit the typical behavior of a Δ_2^0 -set. Theorem 3.1 shows that indeed it is not the case that almost every set in Δ_2^0 is bi-immune for the low sets.

Theorem 3.1. *The low sets do not have Δ_2^0 -measure 0.*

Proof. Let M be a universal Σ_1^0 -martingale and let N be an arbitrary Δ_2^0 -martingale. We will exhibit a low set B on which N does not succeed. Define a new martingale L by

$$\begin{aligned} L(\emptyset) &= \frac{1}{2}(M(\emptyset) + N(\emptyset)), \\ L(\sigma) &= \frac{1}{2}(M(\sigma) + N(\sigma(0)\sigma(2) \dots \sigma(2n))) \text{ if } |\sigma| = 2n + 1 \text{ or } 2n + 2. \end{aligned}$$

So L is essentially a sum of the behaviour of M and the behaviour of N restricted to the even bits. We leave it to the reader to check that L is indeed a martingale.

Now let $A \in \Delta_2^0$ be such that L does not succeed on A . (Such an A exists because Δ_2^0 does not have Δ_2^0 -measure 0.) Then M is bounded on A , and hence A is Σ_1^0 -random. Also, N is bounded on the set B defined by $B(n) = A(2n)$ for every n . We claim that B is low, being half of a Σ_1^0 -random set below \emptyset' . Thus we have exhibited a low set on which the arbitrary Δ_2^0 -martingale N does not succeed.

To prove the claim that B is low, suppose that C is the odd part of A , i.e. the unique set with $A = B \oplus C$. Since A is Σ_1^0 -random, by a result of van Lambalgen [15] the set C is Σ_1^0 -random relative to B . Nies and Stephan (see [7, Theorem 3.4])

⁵Indeed, this is even true for Σ_1^0 -measure: The class of sets that are not bi-immune for the computable sets has Σ_1^0 -measure 0.

showed that this implies that B is low. For completeness, we include a proof of this fact.

Let $\text{use}(\Phi_e^B(e))$ be the partial function that for every e measures the number of computation steps of $\Phi_e^B(e)$, if this is defined. Since C is a Δ_2^0 set, it has a computable approximation C_s such that $\lim_s C_s(n) = C(n)$ for every n . Let the convergence modulus of this approximation be the function $m(n) = \mu t (\forall s \geq t) [C_s(n) = C_t(n)]$. Now if there were infinitely many e such that $\text{use}(\Phi_e^B(e)) \geq m(e)$ then B could compute infinitely many points of C , contradicting the fact that C is Σ_1^0 -random relative to B . Hence, for almost all e , whenever $\Phi_e^B(e)$ is defined we have that $\text{use}(\Phi_e^B(e)) < m(e)$. Since $m \leq_T \emptyset'$ we obtain that $B' \leq_T \emptyset'$. \square

Corollary 3.2. *The Δ_2^0 -Hausdorff dimension of the low sets is 1.*

ACKNOWLEDGMENTS

The second author thanks Bjørn Kjos-Hanssen for several discussions [12], and in particular for pointing out a mistake in [30].⁶ The present paper has greatly benefitted from the comments and suggestions of an anonymous referee, who in particular shortened and improved the proof of Theorem 3.1.

REFERENCES

- [1] K. Ambos-Spies, E. Mayordomo, Y. Wang, and X. Zheng, *Resource-bounded balanced genericity, stochasticity and weak randomness*, Proc. 13th Symp. on Theoret. Aspects of Comp. Sci., Lect. Notes in Comp. Sci. 1046 (1996) 63–74, Springer-Verlag.
- [2] K. Ambos-Spies, W. Merkle, J. Reimann, and F. Stephan, *Hausdorff dimension in exponential time*, Computational Complexity 2001, 210–217, IEEE Computer Society, 2001.
- [3] C. S. Calude, L. Staiger, and S. A. Terwijn, *On partial randomness*, Ann. Pure Appl. Logic 138 (2006) 20–30.
- [4] O. Demuth and A. Kučera, *Remarks on 1-genericity, semigenericity and related concepts*, Comment. Math. Univ. Carolinae, 28 (1987), 85–94.
- [5] R. G. Downey, *Some computability-theoretic aspects of reals and randomness*, in *The Notre Dame Lectures* (P. Cholak, ed.), to appear in the Lect. Notes Log. 18, Assoc. for Symbol. Logic, Urbana, IL, 2005, 97–148.
- [6] R. G. Downey, D. R. Hirschfeldt, S. Lempp, and R. Solomon, *A Δ_2^0 set with no infinite low subset in either it or its complement*, Journal of Symbolic Logic 66 (2001) 1371–1381.
- [7] R. G. Downey, D. R. Hirschfeldt, J. S. Miller, and A. Nies, *Relativizing Chaitin's halting probability*, to appear in the Journal of Mathematical Logic.
- [8] R. G. Downey, D. R. Hirschfeldt, A. Nies, and S. A. Terwijn, *Calibrating randomness*, to appear in the Bulletin of Symbolic Logic.
- [9] K. Falconer, *Fractal Geometry, Mathematical Foundations & Applications*, Wiley & Sons, 1992.
- [10] D. R. Hirschfeldt, A. Nies, and F. Stephan, *Using random sets as oracles*, to appear.
- [11] J. M. Hitchcock, J. H. Lutz and S. A. Terwijn, *The arithmetical complexity of dimension and randomness*, ACM Trans. on Comput. Logic, in press.
- [12] B. Kjos-Hanssen, personal communication, July 2003.
- [13] A. Kučera, *An alternative, priority-free solution to Post's Problem*, in: Lecture Notes in Computer Science 233, 493–500, Springer, 1986.

⁶In Theorem 4.1 in [30] it was proven that the lower cone of every incomplete c.e. set has Δ_2^0 -measure 0. Also, the upper cone of every noncomputable set in Δ_2^0 has Δ_2^0 -measure 0 (Theorem 2.3). Now in Corollary 4.2 it was falsely claimed that from the proofs of these theorems it follows that almost every set in Δ_2^0 is incomparable with every noncomputable and incomplete c.e. set. Kjos-Hanssen pointed out that this contradicts Kučera's result that every 1-random set in Δ_2^0 bounds a noncomputable c.e. set, cf. [13].

- [14] S. A. Kurtz, *Randomness and genericity in the degrees of unsolvability*, PhD Thesis, University of Illinois, 1981.
- [15] M. van Lambalgen, *Random sequences*, PhD Thesis, University of Amsterdam, 1987.
- [16] K. de Leeuw, E. F. Moore, C. F. Shannon, and N. Shapiro, *Computability by probabilistic machines*, in *Automata Studies*, Annals of Mathematics Studies 34, Princeton University Press, 1956, 183–212.
- [17] M. Li and P. Vitányi, *An Introduction to Kolmogorov Complexity and its Applications*, 2nd edition, Springer-Verlag, 1997.
- [18] J. H. Lutz, *Category and measure in complexity classes*, SIAM J. Comput. 19 (1990) 1100–1131.
- [19] J. H. Lutz, *Dimension in complexity classes*, SIAM J. Comput. 32 (2003), 1236–1259 (extended abstract in *15th Annual IEEE Conference on Computational Complexity (Florence, 2000)* (F. Titsworth, ed.), IEEE Computer Soc., Los Alamitos, CA, 2000, 158–169).
- [20] J. H. Lutz, *The dimensions of individual strings and sequences*, Inform. and Comput. 187 (2003), 49–79 (preliminary version: *Gales and the constructive dimension of individual sequences*, in *Proc. 27th International Colloquium on Automata, Languages, and Programming* (U. Montanari, J. D. P. Rolim, E. Welzl, eds.), Springer-Verlag, 2000, 902–913).
- [21] P. Martin-Löf, *The definition of random sequences*, Information and Control 9 (1966) 602–619.
- [22] A. Nies, F. Stephan, and S. A. Terwijn, *Randomness, relativization and Turing Degrees*, Journal of Symbolic Logic 70(2) (2005) 515–535.
- [23] P. G. Odifreddi, *Classical Recursion Theory*, Vol. 1, Studies in logic and the foundations of mathematics Vol. 125, North-Holland, 1989.
- [24] P. G. Odifreddi, *Classical Recursion Theory*, Vol. 2, Studies in logic and the foundations of mathematics Vol. 143, North-Holland, 1999.
- [25] J. Reimann, *Computability and fractal dimension*, PhD Thesis, University of Heidelberg, 2004.
- [26] G. E. Sacks, *Degrees of Unsolvability*, Annals of Mathematics Studies 55, Princeton University Press, 1963.
- [27] C.-P. Schnorr, *Zufälligkeit und Wahrscheinlichkeit*, Lect. Notes in Math. 218, Springer-Verlag, 1971.
- [28] R. Soare, *Recursively Enumerable Sets and Degrees*, Springer, 1987.
- [29] S. A. Terwijn, *Computability and Measure*, PhD Thesis, University of Amsterdam/ILLC, 1998.
- [30] S. A. Terwijn, *On the quantitative structure of Δ_2^0* , in: U. Berger, H. Osswald, and P. Schuster (eds.), *Reuniting the Antipodes — Constructive and Nonstandard Views of the Continuum*, Synthèse Library, Kluwer, 2000, 271–284.
- [31] S. A. Terwijn, *Complexity and Randomness*, Rendiconti del Seminario Matematico di Torino 62 (2004) 1–38 (preliminary version: Research Report CDMTCS-212, University of Auckland, March 2003).
- [32] J. Ville, *Étude Critique de la Notion de Collectif*, Gauthier-Villars, 1939.
- [33] Y. Wang, *A separation of two randomness concepts*, Information Processing Letters 69(3) (1999) 115–118.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, IL 60637, USA
E-mail address: drh@math.uchicago.edu

INSTITUTE OF DISCRETE MATHEMATICS AND GEOMETRY, TECHNICAL UNIVERSITY OF VIENNA,
 WIEDNER HAUPTSTRASSE 8-10/E104, 1040 VIENNA, AUSTRIA
E-mail address: terwijn@logic.at