

ESSENTIAL DIMENSION VIA PRISMATIC COHOMOLOGY

BENSON FARB, MARK KISIN AND JESSE WOLFSON

ABSTRACT. For X a smooth, proper complex variety such that $h_X^{0,i} \neq 0$, we show that for almost all primes p , the restriction of the mod p cohomology $H^i(X, \mathbb{F}_p)$ to any Zariski open is nonzero. The proof uses the prismatic cohomology of Bhatt-Scholze.

We use this result to obtain lower bounds on the p -essential dimension of covers of complex varieties. For example, we prove the p -incompressibility of the mod p homology cover of an abelian variety, confirming a conjecture of Brosnan for sufficiently large p . By combining these techniques with the theory of toroidal compactifications of Shimura varieties, we show that for any Hermitian symmetric domain X , there exist p -congruence covers that are p -incompressible.

CONTENTS

1. Introduction	1
2. Essential dimension and characteristic classes	5
2.1. Review of mod p cohomology	5
2.2. Restriction of mod p étale cohomology	8
2.3. Characteristic classes	12
3. Torus embeddings and Shimura varieties	14
3.1. Cohomology of generalized Heisenberg groups	14
3.2. Toric varieties	18
3.3. Shimura Varieties	23
References	29

1. INTRODUCTION

Let $f : Y \rightarrow X$ be a finite map of complex algebraic varieties. The *essential dimension* $\text{ed}(Y/X)$ of f is the smallest integer e such that, over some dense open of X , the map f arises as the pullback of a map of varieties of dimension e . The motivation for studying this invariant goes back to classical questions about reducing the number of parameters in a solution to a general n^{th} degree polynomial. It first appeared in work of Kronecker [Kr1861] and Klein [Kl1884] on the quintic, and was formally defined by Tschebotarow [Tsc34], and in a modern context by Buhler-Reichstein [BR97], [BR99].

An idea which goes back to Arnol'd [Arn70a], [Arn70b], is to use characteristic classes to obtain a lower bound for essential dimension: if f has covering group G

The authors are partially supported by NSF grants DMS-1811772 (BF), DMS-1902158 (MK) and DMS-1811846 and DMS-1944862 (JW).

then the cohomology of G sometimes contributes to the cohomology of X - such classes on X are called *characteristic*. If they contribute to the cohomology of X in some degree i , and X is affine, then f cannot arise as a pullback of a G -cover of dimension $< i$. The main difficulty with this approach is that to get a lower bound for $\text{ed}(Y/X)$ one needs to know that (some of) the classes coming from G continue to be nontrivial on any Zariski open in X . Indeed in *loc. cit.* Arnol'd was able to address the question only without restricting to open subsets. For the universal S_n -cover, this problem was solved by Buhler-Reichstein [BR99]. Following a suggestion of Serre, they showed the Stiefel-Whitney classes which arise in that situation are nonzero at the generic point.

In this paper we introduce a new method, which solves this restriction problem in many cases. In particular, it allows us to give lower bounds on $\text{ed}(Y/X)$ in many cases when X is *proper*. Previously, lower bounds on the essential dimension of coverings of proper varieties were known only in very special cases [CT02, FKW19, FS21]. In fact our results apply to the so called *p-essential dimension* $\text{ed}(Y/X; p)$ (p a prime), where one is allowed to pull back the covering not just to Zariski opens, but to auxiliary coverings of prime to p degree [RY00]. A first example of our results is the following:

Theorem 1. *Let X be a smooth proper complex variety, and $Y \rightarrow X$ its mod p homology cover. Suppose that X has good unramified reduction at p , and let b_1 denote the first Betti number of X . Then for $p > \max\{\frac{1}{2}b_1, 3\}$,*

$$\text{ed}(Y/X; p) \geq \min\{\dim X, \frac{1}{2}b_1\}.$$

Recall that the mod p homology cover is defined as the covering corresponding to the maximal elementary abelian p -group quotient of the fundamental group of X . We refer the reader to 2.2.12 for the definition of unramified good reduction. This is a condition that holds for almost all p . If X is defined over a number field F , the condition means that there is an unramified prime of F over p at which X has good reduction. The condition holds for *all* primes if X is a generic member of a sufficiently good moduli space, see 2.2.14 below.

There are clearly a plethora of varieties to which Theorem 1 applies. In many cases it implies that, for sufficiently large p , the mod p homology cover $Y \rightarrow X$ is *p-incompressible*, i.e. that $\text{ed}(Y/X; p) = \dim X$. This includes the following examples:

- X an abelian variety.
- $X = C_1 \times \cdots \times C_r$ for curves C_i with $\text{genus}(C_i) \geq 1$.
- Locally symmetric varieties associated to certain cocompact (noncongruence!) lattices in $\text{SU}(n, 1)$. See [BW00, p167, Cor. 5.9] for the construction of these examples.

When X is an abelian variety, this confirms (almost all of) a conjecture of Brosnan [FS21, Conj. 6.1], which was previously known only for either a very general abelian variety, or in dimension at most 3 for a positive density set of primes [FS21]. The result for a product of curves was known only for a generic product of elliptic curves, by a result of Gabber [CT02]. If X is a generic abelian variety, or a product of generic curves, then X has unramified good reduction at all p , and this condition imposes no extra restriction in Theorem 1. We leave it to the reader to find further examples.

Our results are not actually limited to elementary abelian p -covers. One can apply them to G -covers for any finite group G , but the conditions which have to hold are rather more restrictive:

Theorem 2. *Let X be a smooth, proper complex variety, G a finite group, and $Y \rightarrow X$ a G -cover. Suppose that X has unramified good reduction at p , and let $i < p - 2$. If $H^0(X, \Omega_X^i) \neq 0$ and the map $H^i(G, \mathbb{F}_p) \rightarrow H^i(X, \mathbb{F}_p)$ is surjective then*

$$\mathrm{ed}(Y/X; p) \geq i.$$

We also work with open varieties, and not just proper ones, in which case the formulation of the above results involves logarithmic differentials, see 2.3.3, 2.3.8 below. Note that the condition on surjectivity in Theorem 2 is quite restrictive. If X is an étale $K(\pi, 1)$, then there always exists a G , and a G -cover satisfying the condition, but this is not true in general.

In §3.2 we explain how to apply our results to torus torsors over abelian schemes. Here the fundamental group is a *generalized Heisenberg group* - a central extension of finitely-generated free abelian groups - and there is a natural notion of “reduction mod p ” for such a group. We show that the corresponding covers are p -incompressible for large enough p , see 3.2.7.

We then use Theorem 2 to deduce the p -incompressibility of certain covers of locally symmetric varieties. These have the form $\Gamma_1 \backslash X \rightarrow \Gamma \backslash X$, where X is a Hermitian symmetric domain and Γ is an arithmetic lattice in the corresponding real semisimple Lie group. Thus, Γ is a congruence subgroup of $G(\mathbb{Q})$ for a semisimple \mathbb{Q} -group G . We consider *principal p -congruence covers* which means, roughly speaking, that there are no congruences mod p involved in the definition of Γ , and Γ_1 is the subgroup of elements which are trivial mod p . A sample of our results is the following.

Theorem 3. *Suppose X is an irreducible Hermitian domain and that the \mathbb{Q} -rank of G (the rank of its maximal \mathbb{Q} -split torus) is equal to its \mathbb{R} -rank. For any principal p -congruence cover $\Gamma_1 \backslash X \rightarrow \Gamma \backslash X$, we have*

$$\mathrm{ed}(\Gamma_1 \backslash X \rightarrow \Gamma \backslash X; p) = \dim X,$$

provided p satisfies the following conditions if X is not a tube domain:

- *If X is of classical type, then $p > \frac{3}{2} \dim X$.*
- *If X is of type E_6 , then p is sufficiently large.*

Note that for *any* irreducible Hermitian domain there are many examples where Theorem 3 applies. Our results actually apply to many examples of quotients of *reducible* Hermitian domains by irreducible lattices, for example the case of Hilbert modular varieties. See Lemma 3.3.8 and the discussion in 3.3.9.

Theorem 3 is *not* proved directly using our results on characteristic classes. Rather, we relate principal p -congruence covers to the covers of torus torsors over abelian varieties, with covering group a generalized mod p Heisenberg group mentioned above, and we then apply our results about these latter covers. The connection between these two kinds of covers makes use of the theory of toroidal compactifications of Shimura varieties.

We remark that when X is a tube domain, the corresponding torus torsor is just a torus; the abelian variety is 0-dimensional. In this case, one does not need our results on characteristic classes (this is why there is no restriction on p in this case).

The theorem can be deduced from a result of Burda [Bur12] on coverings of tori. The result when X is a tube domain has also been proven by Brosnan-Fakhruddin [BF20], who used the fixed-point method instead of Burda's results.

The conditions on G and p in Theorem 3 are completely different from those considered in [FKW19, FS21]. For example, in many cases when the results of *loc. cit.* apply, they give p -incompressibility not for almost all primes, but only for an explicit set of primes of positive density. They are also restricted to groups of classical type. On the other hand, proper varieties pose no special problem for these results, whereas they cannot be handled by Theorem 3. This also shows, that one should not expect the bounds on p in Theorem 3 to be sharp, as there are also many cases which are covered by *both* Theorem 3 and [FKW19, FS21], but where the latter results impose no similar lower bound on p .

As mentioned above, our lower bounds on essential dimension rely on results that assert that cohomology classes do not vanish on restriction to any Zariski open. An example of such a statement is the following:

Theorem 4. *Let X be a smooth, proper, complex variety, with unramified good reduction at p . Let $i < p - 2$, and $W \subset X$ any Zariski open. Then the image of the restriction map*

$$H^i(X, \mathbb{F}_p) \rightarrow H^i(W, \mathbb{F}_p)$$

has dimension at least $h_X^{0,i} = \dim H^0(X, \Omega_X^i)$.

To get a feel for what such a statement entails, suppose that the integral cohomology $H^*(X, \mathbb{Z})$ is torsion free. Then Theorem 4 asserts that certain classes in the image of $H^i(X, \mathbb{Z}) \rightarrow H^i(W, \mathbb{Z})$ are not divisible by p . Although this is a statement about the topology of complex algebraic varieties, it appears to be out of reach of classical methods (the analogue with \mathbb{Q} -coefficients can be proved using Hodge theory, but this does not suffice for applications to essential dimension).

Our proof of Theorem 4 makes use of *prismatic cohomology*, recently introduced by Bhatt-Scholze [BS19]. This is a cohomology theory that takes as an input a smooth formal scheme \mathcal{X} over a p -adic base and, in some sense, interpolates between the mod p (or more generally p -adic) étale cohomology of its generic fiber, and the de Rham cohomology of its special fiber \mathcal{X}_k . Using it, we translate the statement of Theorem 4 into the analogous statement for de Rham cohomology of \mathcal{X}_k , and then into a statement about differentials using the Cartier isomorphism.

To deduce Theorem 2 from Theorem 4 one considers the composite

$$H^i(G, \mathbb{F}_p) \rightarrow H^i(X, \mathbb{F}_p) \rightarrow H^i(W, \mathbb{F}_p).$$

Theorem 4 and the assumptions of Theorem 2 guarantee this map is nonzero. However if $Y|_W \rightarrow W$ arises from a covering of varieties of dimension $< i$, then we may assume that these varieties are affine, and it follows that the above map must vanish since the cohomological dimension of affine varieties is at most their dimension.

We also prove a variant of Theorem 4 where we consider the image of the map $\wedge^i H^1(X, \mathbb{F}_p) \rightarrow H^i(W, \mathbb{F}_p)$ given by the cup product and restriction. This leads to Theorem 1. Using the trace map, it is easy to deduce from Theorem 4 a version where one allows restriction to a prime-to- p covering of W , so we obtain lower bounds on p -essential dimension.

The paper is organized as follows. In §2.1 we review the results we need from prismatic and crystalline cohomology. In §2.2 we prove Theorem 4 and its variants, and in §2.3 we apply this to characteristic classes to deduce lower bounds on p -essential dimension. Throughout §2 we work with schemes or formal schemes equipped with a normal crossings divisor, and we consider coverings of the complement. This greater level of generality is needed for the applications in §3. In §3.1, we carry out an analysis of the mod p cohomology of (generalized) finite Heisenberg groups. This is used in §3.2 to apply our results on characteristic classes to torus embeddings over abelian varieties, by verifying the surjectivity assumption in Theorem 2 (or more precisely its logarithmic analogue) in this case. Most of §3.2 is actually concerned with a variant of Theorem 4 for torus embeddings over an abelian scheme, where we restrict not just to Zariski opens, but to analytic neighborhoods of the boundary. This is then applied in §3.3 to obtain our results on the p -essential dimension of congruence covers: the analytic neighborhoods map to our locally symmetric varieties and we can restrict the congruence covers to them.

Acknowledgments. We thank Dave Benson, H el ene Esnault, Keerthi Madapusi Pera, Alexander Petrov, Gopal Prasad and Chris Rogers for useful discussions and suggestions. We thank Patrick Brosnan, Najmuddin Fakhruddin and Peter Scholze for helpful comments on a draft.

2. ESSENTIAL DIMENSION AND CHARACTERISTIC CLASSES

2.1. Review of mod p cohomology.

2.1.1. In this subsection we review the results we will need on de Rham and prismatic cohomology. We begin with the former, see Deligne-Illusie [DI, 4.2.3].

Let k be a perfect field of characteristic p , and let X be a smooth k -scheme. Suppose that X is equipped with a normal crossings divisor $D \subset X$. Let $\Omega_{X/k}^\bullet(\log D)$ denote the logarithmic de Rham complex with poles in D .

Let $X^1 = X \times_{\mathrm{Spec} k, F_k} \mathrm{Spec} k$ and let $D^1 = D \times_{\mathrm{Spec} k, F_k} \mathrm{Spec} k$, where F_k denotes the Frobenius on k . Let $F_{X/k} : X \rightarrow X^1$ denote the relative Frobenius; it is a finite flat map of k -schemes taking D to D^1 .

Lemma 2.1.2. *Suppose that (X, D) admits a lift ¹ to a smooth formal scheme \tilde{X} over $W(k)$, equipped with a normal crossing divisor \tilde{D} relative to $W(k)$. Then for $j < p$, $H^0(X^1, \Omega_{X^1/k}^j(\log D^1))$ is canonically a direct summand of the de Rham cohomology $H^j(X, \Omega_{X/k}^\bullet(\log D))$.*

Proof. For $n \geq 0$, recall the truncation $\tau_{<n} F_{X/k*} \Omega_{X/k}^\bullet(\log D)$, whose cohomology agrees with that of $F_{X/k*} \Omega_{X/k}^\bullet(\log D)$ in degrees $< n$ and vanishes otherwise. By [DI87, 4.2.3], the complex of \mathcal{O}_{X^1} -modules $\tau_{<p} F_{X/k*} \Omega_{X/k}^\bullet(\log D)$ is canonically (in a way which depends on the chosen lifting of (X, D)) quasi-isomorphic to the direct sum of its cohomology sheaves, so that the Cartier isomorphism induces an isomorphism in the derived category of \mathcal{O}_{X^1} -modules

$$\bigoplus_{i < p} \Omega_{X^1/k}^i(\log D^1)[-i] \simeq \tau_{<p} F_{X/k*} \Omega_{X/k}^\bullet(\log D).$$

¹In fact for what follows only a lifting to the Witt vectors of length 2, W_2 is required, but we will not need this greater generality.

Hence $H^0(X^1, \Omega_{X^1/k}^j(\log D^1))$ is a direct summand in

$$H^j(X^1, \tau_{<p} F_{X/k*} \Omega_{X/k}^\bullet(\log D)) \simeq H^j(X, \tau_{<p} \Omega_{X/k}^\bullet(\log D))$$

which agrees with $H^j(X, \Omega_{X/k}^\bullet(\log D))$, as $j < p$ and the cohomology of the cone of $\tau_{<p} \Omega_{X/k}^\bullet(\log D) \rightarrow \Omega_{X/k}^\bullet(\log D)$ vanishes in degree $< p$. \square

2.1.3. Let K be a field of characteristic 0. By a *p-adic valuation* on K , we mean a rank one valuation v on K , with $v(p) > 0$. We suppose that K is complete with respect to such a valuation, with ring of integers \mathcal{O}_K and perfect residue field k . We now recall the facts we will need about the prismatic cohomology of smooth formal schemes over \mathcal{O}_K .

We do not recall the general formalism of prisms, as developed by Bhatt-Scholze [BS19], and Koshikawa [Kos20] in the logarithmic case, but consider only the two examples we will need. First suppose that the valuation on K is discrete. Set $A = W(k)[[u]]$, equipped with a Frobenius φ which extends the Frobenius on W by sending u to u^p . We equip A with the map $A \rightarrow \mathcal{O}_K$ sending u to some chosen uniformizer π . Its kernel is generated by an Eisenstein polynomial $E(u) \in W(k)[u]$ for π . In fact, in applications we will assume $\mathcal{O}_K = W(k)$, and $\pi = p$. Then $(A, E(u)A)$ is the so called *Breuil-Kisin* prism.

Next suppose that K is algebraically closed. Let $R = \varprojlim \mathcal{O}_K/p$ with the maps in the inverse limit being given by the absolute Frobenius. We take $A = A_{\text{inf}} = W(R)$, with its canonical Frobenius. Any element $x = (x_0, x_1, \dots) \in R$ lifts uniquely to a sequence $(\hat{x}_0, \hat{x}_1, \dots)$ in \mathcal{O}_K with $\hat{x}_i^p = \hat{x}_{i-1}$. There is a natural surjective map of rings $\theta : A \rightarrow \mathcal{O}_K$, which sends a Teichmüller element x as above to \hat{x}_0 . The kernel of θ is principal, generated by $\xi = p - \underline{p}$, where $\underline{p} = (p, p^{1/p}, \dots)$. Then $(A, \xi A)$ is an example of a perfect prism.

2.1.4. In the rest of this section we will make use of logarithmic formal schemes over \mathcal{O}_K , logarithmic schemes over K , and logarithmic adic spaces over K . In particular, when K is algebraically closed we will consider the étale cohomology of logarithmic schemes and adic spaces over K with coefficients in \mathbb{F}_p . We refer the reader to [Kat89] for a general introduction to logarithmic geometry, and to [Ill02] and [DLLZ19] for étale cohomology in this context.

2.1.5. Now assume that K is either discretely valued or algebraically closed. Let X be a formally smooth \mathcal{O}_K -scheme, equipped with a relative normal crossings divisor D . We write X_D for the formal scheme X equipped with the logarithmic structure given by D . We will denote by $X_{D,K}$ the associated logarithmic adic space. When the divisor D is empty we drop it from the notation.

By [BS19] and [Kos20], the *prismatic cohomology* of X_D is a complex of A -modules $R\Gamma_{\Delta}(X_D/A)$, equipped with a φ -semi-linear map $\varphi : R\Gamma_{\Delta}(X_D/A) \rightarrow R\Gamma_{\Delta}(X_D/A)$. As we will only be interested in mod p cohomology we set

$$\overline{R\Gamma_{\Delta}(X_D/A)} = R\Gamma_{\Delta}(X_D/A) \otimes_A^L A/pA,$$

and we will denote by $\overline{H_{\Delta}^i(X_D/A)}$ the cohomology of $\overline{R\Gamma_{\Delta}(X_D/A)}$. Then we have the following properties, where we work in the derived category $D(A)$ of the category of A -modules and again only consider mod p cohomology.

- (1) There is a canonical isomorphism of commutative algebras in $D(A)$

$$R\Gamma(\Omega_{X \otimes k/k}^\bullet(\log D \otimes k)) \simeq \overline{R\Gamma_{\Delta}(X_D/A)} \otimes_{A/pA, \varphi}^L k.$$

- (2) If K is algebraically closed then there is an isomorphism of commutative algebras in $D(A)$

$$R\Gamma_{\text{ét}}(X_{D,K}, \mathbb{F}_p) \simeq \overline{R\Gamma_{\Delta}(X_D/A)}[1/\xi]^{\varphi=1}$$

where the right hand side means the fiber of $\varphi-1$ acting on $\overline{R\Gamma_{\Delta}(X_D/A)}[1/\xi]$, $X_{D,K}$ denotes the adic space X_K associated to X , equipped with the log structure given by D_K .

- (3) The linearization

$$\varphi^*(\overline{R\Gamma_{\Delta}(X_D/A)}) \rightarrow \overline{R\Gamma_{\Delta}(X_D/A)}$$

becomes an isomorphism in $D(A)$ after inverting u (resp. ξ) if K is discrete (resp. algebraically closed). For each $i \geq 0$, there is a canonical map

$$V_i : \overline{H_{\Delta}^i(X_D/A)} \rightarrow H^i(\varphi^* \overline{R\Gamma_{\Delta}(X_D/A)})$$

with $V_i \circ \varphi = \varphi \circ V_i$ equal to u^{ie} (resp. ξ^i) if K is discrete (resp. algebraically closed), where e denotes the ramification degree of \mathcal{O}_K over $W(k)$.

- (4) Let K' be a field complete with respect to a p -adic valuation, and which is either discrete or algebraically closed. Let $A' \rightarrow \mathcal{O}_{K'}$ be the corresponding prism, as defined above. Suppose we are given map of valued fields $K \rightarrow K'$, and a map $A \rightarrow A'$ compatible with the projections to $\mathcal{O}_K \rightarrow \mathcal{O}_{K'}$, and with Frobenius maps. Then there is a canonical isomorphism in $D(A)$

$$\overline{R\Gamma_{\Delta}(X_D/A)} \otimes_A A' \simeq \overline{R\Gamma_{\Delta}(X_{D,\mathcal{O}_{K'}}/A')}$$

We shall apply this when K is discrete and K' is algebraically closed.

- (5) When X is proper over \mathcal{O}_K , then $\overline{R\Gamma_{\Delta}(X_D/A)}$ is a perfect complex of A/p -modules.

In the non-logarithmic case, when D is empty this is [BS19, Thm. 1.8]. In the logarithmic case (1), (4) and (5) are contained in [Kos20, §5]. The other properties should be in forthcoming work of Koshikawa-Yao [KY].

Using (5), when X is proper, we have the following more explicit form of (2) above.

Lemma 2.1.6. *Suppose that K is algebraically closed, and that X is proper over \mathcal{O}_K . Then for each $i \geq 0$ there is a natural isomorphism*

$$H_{\text{ét}}^i(X_{D,K}, \mathbb{F}_p) \otimes_{\mathbb{F}_p} A/pA[1/\xi] \simeq \overline{H_{\Delta}^i(X_D/A)}[1/\xi].$$

Proof. By (2) above there is an exact sequence in cohomology

$$\cdots \rightarrow H_{\text{ét}}^i(X_{D,K}, \mathbb{F}_p) \rightarrow \overline{H_{\Delta}^i(X_D/A)}[1/\xi] \xrightarrow{1-\varphi} \overline{H_{\Delta}^i(X_D/A)}[1/\xi] \rightarrow H_{\text{ét}}^{i+1}(X_{D,K}, \mathbb{F}_p)$$

Now $A/pA[1/\xi] = \text{Frac}R$ is an algebraically closed field, and $\overline{H_{\Delta}^i(X_D/A)}[1/\xi]$ is a finite dimensional $A/pA[1/\xi]$ vector space by (5) above. Hence $1-\varphi$ is surjective on $\overline{H_{\Delta}^i(X_D/A)}[1/\xi]$, and this space is spanned by its Frobenius invariants [Kat73, Prop. 4.1.1]. \square

2.2. Restriction of mod p étale cohomology. The goal of this subsection is to prove a result on restriction of étale cohomology classes to open neighborhoods. We begin with a technical lemma. As above, k denotes a perfect field of characteristic p .

Lemma 2.2.1. *Let M be a finitely generated $k[[u]]$ -module, equipped with a Frobenius semi-linear map $\varphi : M \rightarrow M$, and a linear map $V : M \rightarrow \varphi^*M$ such that $\varphi \circ V = V \circ \varphi = u^d$ for some integer d . If $d + 1 < p$, then M is torsion free.*

Proof. Suppose that M contains nontrivial torsion. Since M is finitely generated over $k[[u]]$, there is a minimal integer $m \geq 1$ such that $u^m \cdot x = 0$ for any torsion element $x \in M$. Choose a torsion element x so that $u^{m-1} \cdot x \neq 0$. Then in φ^*M we have

$$u^{d+m}(1 \otimes x) = V \circ (1 \otimes \varphi)(u^m \otimes x) = V(u^m \varphi(x)) = 0,$$

since $u^{pm} \varphi(x) = 0$, so that $\varphi(x)$ is torsion in M . On the other hand, as φ is finite flat over $k[[u]]$, the smallest power of u which kills $1 \otimes x \in \varphi^*M$ is pm . Thus $d + m \geq pm$, which implies $p \leq 1 + d/m \leq 1 + d$. \square

2.2.2. Now let K be as in 2.1.3 and suppose that K is discrete, and that $\mathcal{O}_K = W(k)$. Let C be an algebraically closed field equipped with a complete p -adic valuation, and $K \subset C$ an inclusion of valued fields. As before, for a formal scheme or formal log. scheme X over \mathcal{O}_K , we denote by X_K and X_C the associated adic spaces over K and C respectively. We denote by $X_{\mathcal{O}_C} = X \hat{\otimes}_{\mathcal{O}_K} \mathcal{O}_C$ the base change of X to \mathcal{O}_C , as a formal scheme, and by X_k the special fiber (i.e the reduced subscheme) of X .

For the next three results we let X be a proper, smooth formal scheme over \mathcal{O}_K , equipped with a relative normal crossings divisor $D \subset X$. Let

$$h_{(X,D)}^{0,i} := \dim_K H^0(X_K, \Omega_{X_K/K}^i(\log D)).$$

Proposition 2.2.3. *Let $W \subset X - D$ be a dense open formal subscheme. Then for $0 \leq i < p - 2$,*

$$\dim_{\mathbb{F}_p} \text{Im} (H_{\text{ét}}^i(X_{D,C}, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(W_C, \mathbb{F}_p)) \geq h_{(X,D)}^{0,i}.$$

Proof. As $W \subset X - D$, we will omit the divisor D from the notation when writing the cohomology of W . We take the prism A in 2.1.3 to be $W(k)[[u]]$, with $E(u) = u - p$. We can also apply the construction of 2.1.3 to C , to obtain a prism $A_C \rightarrow \mathcal{O}_C$. There is a Frobenius compatible map $A \rightarrow A_C$ sending u to $[p]$.

Set

$$M_{\Delta} = \text{Im} (\overline{H_{\Delta}^i(X_D/A)} \rightarrow \overline{H_{\Delta}^i(W/A)}),$$

which is a finitely generated $A/pA = k[[u]]$ -module. By 2.1.5(4), there is an isomorphism

$$\overline{H_{\Delta}^i(X_D/A)} \otimes_A A_C \xrightarrow{\sim} \overline{H_{\Delta}^i(X_{D,\mathcal{O}_C}/A_C)},$$

and similarly for W . Thus using Lemma 2.1.6 we have maps

$$(2.2.4) \quad \begin{aligned} \overline{H_{\Delta}^i(X_D/A)} \otimes_A A_C[1/\xi] &\simeq H_{\text{ét}}^i(X_{D,C}, \mathbb{F}_p) \otimes_{\mathbb{F}_p} A_C/pA_C[1/\xi] \\ &\rightarrow H_{\text{ét}}^i(W_C, \mathbb{F}_p) \otimes_{\mathbb{F}_p} A_C/pA_C[1/\xi] \rightarrow \overline{H_{\Delta}^i(W/A)} \otimes_A A_C[1/\xi], \end{aligned}$$

the composite being the natural map. Hence

$$\dim_{\mathbb{F}_p} \text{Im} (H_{\text{ét}}^i(X_{D,C}, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(W_C, \mathbb{F}_p)) \geq \dim_{k((u))} M_{\Delta}[1/u].$$

By Lemma 2.2.1, 2.1.5(3) and our assumptions on i imply that M_Δ is a finitely generated, free $k[[u]]$ -module. Hence it suffices to show $\dim_k M_\Delta / uM_\Delta \geq h_{(X,D)}^{0,i}$.

Using Lemma 2.2.1 again, we see that $\overline{H_\Delta^j(X_D/A)}$ is u -torsion free for $0 \leq j \leq i+1$. Hence 2.1.5(1) implies that we have maps

$$(2.2.5) \quad \begin{aligned} H^i(X_k, \Omega_{X_k/k}^\bullet(\log D)) &\simeq \overline{H_\Delta^i(X_D/A)} \otimes_{A,\varphi} k \rightarrow M_\Delta \otimes_{A,\varphi} k \\ &\rightarrow \overline{H_\Delta^i(W/A)} \otimes_{A,\varphi} k \rightarrow H^i(W_k, \Omega_{W_k/K}^\bullet(\log D)), \end{aligned}$$

where the composite map is the natural map. This shows the image of (2.2.5) has dimension $\leq \dim_k M_\Delta / uM_\Delta$, and it suffices to show that this dimension is $\geq h_{(X,D)}^{0,i}$. Since $W \subset X$ is dense, the map

$$H^0(X_k, \Omega_{X_k/k}^i(\log D)) \rightarrow H^0(W_k, \Omega_{X_k/k}^i),$$

is injective. Hence by Lemma 2.1.2, the image of (2.2.5) has dimension at least $\dim_k H^0(X_k, \Omega_{X_k/k}^i) \geq h_{(X,D)}^{0,i}$, where the last inequality follows from the upper semi-continuity of h^0 . \square

Corollary 2.2.6. *With the above assumptions, suppose also that $\overline{H_\Delta^{i+1}(X_D/A)}$ is u -torsion free. Then the conclusion of Proposition 2.2.3 holds for $i \leq p-2$. In particular, the conclusion holds for $i \leq p-2$ if X is the formal completion of an abelian scheme over \mathcal{O}_K , and D is empty.*

Proof. The stronger assumption $i+1 < p-1$ was used in the proof of Proposition 2.2.3 only to know that $\overline{H_\Delta^{i+1}(X_D/A)}$ is u -torsion free, so the first claim follows. For the second claim, we remark that Anschütz-Le Bras [ALB19, Prop. 4.5.2] have shown that when X is the formal completion of an abelian scheme, the ring $\overline{H_\Delta^*(X/A)}$ is the exterior algebra on the $k[[u]]$ -module $\overline{H_\Delta^1(X/A)}$ which is free of rank $2g = 2\dim X_K$ over $k[[u]]$. \square

2.2.7. The following variant of Proposition 2.2.3 will be useful in obtaining lower bounds on the essential dimension of covers whose groups are elementary p -groups.

Proposition 2.2.8. *Let $W \subset X - D$ be a dense open formal subscheme, and $i \leq \dim X_K$. Suppose $p > \max\{i+1, 3\}$. Then*

$$\dim_{\mathbb{F}_p} \text{Im} (\wedge^i H_{\text{ét}}^1(X_{D,C}, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(W_C, \mathbb{F}_p)) \geq \binom{h_{(X,D)}^{0,1}}{i},$$

where the map is given by the cup product followed by restriction of classes to W_C .

Proof. As the proof is analogous to that of Proposition 2.2.3, we only sketch it. Let

$$M_\Delta = \text{Im} (\wedge^i \overline{H_\Delta^1(X_D/A)} \rightarrow \overline{H_\Delta^i(W/A)}).$$

Arguing as in Proposition 2.2.3, since $i < p-1$, it suffices to show $\dim_k M_\Delta / uM_\Delta \geq \binom{h_{(X,D)}^{0,1}}{i}$. Since $p > 3$, $\overline{H_\Delta^2(X_D/A)}$ is u -torsion free by Lemma 2.2.1, and we reduce to showing that the image of

$$\wedge^i H^1(X_k, \Omega_{X_k/k}^\bullet(\log D)) \rightarrow H^i(W_k, \Omega_{X_k/K}^\bullet(\log D)),$$

has dimension $\geq \binom{h_{(X,D)}^{0,1}}{i}$. This follows from the fact that the composite map

$$\wedge^i H^0(X_k, \Omega_{X_k/k}^1(\log D)) \rightarrow H^0(X_k, \Omega_{X_k/k}^i(\log D)) \rightarrow H^0(W_k, \Omega_{X_k/k}^i)$$

is injective if $i \leq \dim X_K$. \square

2.2.9. We now want to deduce analogues of the above results for schemes. Let Y be a smooth scheme over C , equipped with a normal crossings divisor $D \subset Y$. As above, we write $h_{(Y,D)}^{0,i} = \dim_C H^0(Y, \Omega_{Y/C}^i(\log D))$.

We will denote by Y^{ad} and D^{ad} the adic spaces attached to Y , and D respectively. Then $(Y^{\text{ad}}, D^{\text{ad}})$ is a logarithmic adic space. We begin by recording a lemma comparing the étale cohomology of logarithmic adic spaces and logarithmic schemes.

Lemma 2.2.10. *Let Y be a proper scheme over C , equipped with a normal crossings divisor $D \subset Y$, and set $U = Y - D$. Then for $i \geq 0$ there are natural isomorphisms*

$$H_{\text{ét}}^i(Y_{D^{\text{ad}}}, \mathbb{F}_p) \simeq H_{\text{ét}}^i(Y_D, \mathbb{F}_p) \simeq H_{\text{ét}}^i(U, \mathbb{F}_p)$$

Proof. For second isomorphism see [Ill02, Thm. 7.4]. The first isomorphism follows from [Ill02, Thm 7.2], [DLLZ19, Lem. 4.6.2] and [Hub96, Prop. 2.1.4, Thm 3.8.1]. More precisely, the first two references show that there is a bounded complex of \mathbb{F}_p -vector spaces M on the étale site $Y_{\text{ét}}$ of Y , such that

- $H_{\text{ét}}^i(Y, M)$ is naturally isomorphic to $H_{\text{ét}}^i(Y_D, \mathbb{F}_p)$.
- If M^{ad} denotes the pullback of M to $Y_{\text{ét}}^{\text{ad}}$, then $H_{\text{ét}}^i(Y^{\text{ad}}, M^{\text{ad}})$ is naturally isomorphic to $H_{\text{ét}}^i(Y_{D^{\text{ad}}}, \mathbb{F}_p)$.

The two results in Huber's book then provide a natural isomorphism $H_{\text{ét}}^i(Y, M) \simeq H_{\text{ét}}^i(Y^{\text{ad}}, M^{\text{ad}})$. \square

Proposition 2.2.11. *Let X be a proper smooth scheme over \mathcal{O}_K , equipped with a relative normal crossings divisor $D \subset X$. Set $U = X \setminus D$, and let $W \subset U_C$ be a dense open subscheme. If $0 \leq i < p - 2$, then*

$$\dim_{\mathbb{F}_p} \text{Im} (H_{\text{ét}}^i(U_C, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(W, \mathbb{F}_p)) \geq h_{(X_C, D_C)}^{0,i}.$$

If X is an abelian scheme, then the same statement holds for $i \leq p - 2$.

If $p > \max\{i + 1, 3\}$ and $i \leq \dim X_K$, then

$$\dim_{\mathbb{F}_p} \text{Im} (\wedge^i H_{\text{ét}}^1(U_C, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(W, \mathbb{F}_p)) \geq \binom{h_{(X_C, D_C)}^{0,1}}{i}.$$

Proof. Let k_C be the residue field of C . We may replace X by its base change to $W(k_C)$, and assume that C and K have the same residue field. Denote by \widehat{X} and \widehat{D} the formal completions of X and D . Let Z be the closure of $X_C - W$ in $X_{\mathcal{O}_C}$, and let $\widehat{W} \subset \widehat{X}$ be the formal open subscheme, which is the complement of $Z \otimes k$ in \widehat{X} . Note that we have $\widehat{W}_C \subset W^{\text{ad}}$. Thus we have a commutative diagram of natural maps

$$\begin{array}{ccc} H_{\text{ét}}^i(X_{D,C}, \mathbb{F}_p) & \longrightarrow & H_{\text{ét}}^i(W, \mathbb{F}_p) \\ \downarrow \sim & & \downarrow \\ & & H_{\text{ét}}^i(W^{\text{ad}}, \mathbb{F}_p) \\ \downarrow & & \downarrow \\ H_{\text{ét}}^i(\widehat{X}_{D,C}, \mathbb{F}_p) & \longrightarrow & H_{\text{ét}}^i(\widehat{W}_C, \mathbb{F}_p) \end{array}$$

Where the isomorphism on the left is given by Lemma 2.2.10. By Propositions 2.2.3, 2.2.8 and Corollary 2.2.6, we have the inequalities claimed in the proposition, but for the dimension of the images of $H_{\text{ét}}^i(\widehat{X}_{D,C}, \mathbb{F}_p)$ and $\wedge^i H_{\text{ét}}^1(\widehat{X}_{D,C}, \mathbb{F}_p)$ in $H_{\text{ét}}^i(\widehat{W}_C, \mathbb{F}_p)$. The proposition now follows from the commutative diagram above, as well as Lemma 2.2.10. \square

2.2.12. The previous results apply for schemes that are smooth over \mathcal{O}_K . We now want to reformulate these results to show that they hold for any algebraically closed field, for almost all p . Thus we now assume that C is algebraically closed, but we no longer assume it is equipped with a complete p -adic valuation.

Let Y be a proper, smooth scheme over C , and $D \subset Y$ a normal crossings divisor. We say that (Y, D) has *good reduction at p* if there exists a p -adic valuation on C (which we do not assume complete), with ring of integers \mathcal{O}_C , and Y extends to a smooth proper \mathcal{O}_C -scheme Y° with a relative normal crossings divisor $D^\circ \subset Y^\circ$ over \mathcal{O}_C , extending D . We say that (Y, D) has *unramified good reduction at p* if in addition (Y°, D°) can be chosen so that it descends to an absolutely unramified discrete valuation ring (with respect to the given valuation) $\mathcal{O} \subset \mathcal{O}_C$.

Corollary 2.2.13. *Let Y be a proper, smooth, connected scheme over C , $D \subset Y$ a normal crossings divisor, and $W \subset U = Y - D$ a dense open subscheme. Suppose that (Y, D) has unramified good reduction at p . If $0 \leq i < p - 2$, then*

$$\dim_{\mathbb{F}_p} \text{Im} (H_{\text{ét}}^i(U, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(W, \mathbb{F}_p)) \geq h_{(Y,D)}^{0,i}.$$

If Y is an abelian scheme, then the same statement holds for $i \leq p - 2$.

If $p > \max\{i + 1, 3\}$ and $i \leq \dim Y$, then

$$\dim_{\mathbb{F}_p} \text{Im} (\wedge^i H_{\text{ét}}^1(U, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(W, \mathbb{F}_p)) \geq \binom{h_{(Y,D)}^{0,1}}{i}.$$

Proof. If $C \subset C'$ is any algebraically closed field, then the étale cohomology groups in the corollary do not change if we replace (Y, D) and W by their base change to C' . Thus we may assume that C is complete. We may then also assume that $\mathcal{O} \subset \mathcal{O}_C$ in 2.2.12 is p -adically complete, and unramified. The result now follows from Proposition 2.2.11. \square

2.2.14. We remark that when (Y, D) is a generic member of a good moduli space, then (Y, D) has unramified good reduction at *all* primes, and this condition in Corollary 2.2.13 can then be suppressed.

More precisely, suppose $\mathcal{Y} \rightarrow \mathcal{M}$ is proper smooth, with \mathcal{M} a smooth, faithfully flat, connected, separated Deligne-Mumford stack over \mathbb{Z} , as in [LMB00, Definition 4.1, 4.14]. Let $\mathcal{D} \subset \mathcal{Y}$ be a relative normal crossing divisor over \mathcal{M} . If $\text{Spec } C \rightarrow \mathcal{M}$ is a map whose image is the generic point $\eta \in \mathcal{M}$, then $(Y, D) = (\mathcal{Y} \times_{\mathcal{M}} \text{Spec } C, \mathcal{D} \times_{\mathcal{M}} \text{Spec } C)$ has unramified good reduction at all primes.

To see this, note that our assumptions on \mathcal{M} imply that there exists an étale presentation $\tilde{\mathcal{M}} \rightarrow \mathcal{M}$, with $\tilde{\mathcal{M}}$ a scheme which is smooth and faithfully flat over \mathbb{Z} . Hence η lifts to a generic point $\tilde{\eta} \in \tilde{\mathcal{M}}(C)$, which admits a unique specialization to a characteristic p point $\bar{\eta} \in \mathcal{M}$. As $\tilde{\mathcal{M}}$ is smooth over \mathbb{Z} , the local ring $\mathcal{O}_{\tilde{\mathcal{M}}, \bar{\eta}}$ is a discrete valuation ring with uniformizer p , and this induces the required unramified p -adic valuation on C . This discussion applies, for example, to the universal family of principally polarized abelian varieties or curves.

Corollary 2.2.15. *Let Y be a proper, connected scheme over C , $D \subset Y$ a normal crossings divisor, $U = Y - D$, and $\eta \in Y$ the generic point. Then for $i \geq 0$, and almost all primes p ,*

$$\dim_{\mathbb{F}_p} \text{Im} (H_{\text{ét}}^i(U, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(\eta, \mathbb{F}_p)) \geq h_{(Y,D)}^{0,i}.$$

If $i \leq \dim Y$, then for almost all primes p

$$\dim_{\mathbb{F}_p} \text{Im} (\wedge^i H_{\text{ét}}^1(U, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(\eta, \mathbb{F}_p)) \geq \binom{h_{(Y,D)}^{0,1}}{i}.$$

Proof. By Corollary 2.2.13, it suffices to show that (Y, D) has unramified good reduction at p for almost all p . Since Y is of finite type, there exists finite type \mathbb{Z} -algebra $\mathcal{A} \subset C$ such that (Y, D) descends to a proper smooth \mathcal{A} -scheme $Y_{\mathcal{A}}$ equipped with a relative normal crossings divisor $D_{\mathcal{A}} \subset Y_{\mathcal{A}}$.

We may replace \mathcal{A} by its normalization, and assume it is normal. Since $\mathcal{A} \otimes \mathbb{Q}$ is reduced, it is geometrically reduced [Gro61]. Hence for almost all primes p , the \mathbb{F}_p -algebra $\mathcal{A}/p\mathcal{A}$ is reduced. Since \mathcal{A} is of finite type over \mathbb{Z} , for almost all primes p , p is the image of a prime $\mathfrak{p} \in \text{Spec } \mathcal{A}$ of height 1. Fix p such that $\mathcal{A}/p\mathcal{A}$ is reduced, and \mathfrak{p} exists. Since $\mathcal{A}/p\mathcal{A}$ is reduced, $\mathcal{A}_{\mathfrak{p}}$ is a discrete valuation ring, with uniformizer p . Extend the corresponding valuation $v_{\mathfrak{p}}$ to C . Then (Y, D) descends to $(Y_{\mathcal{A}}, D_{\mathcal{A}})$, so (Y, D) has unramified good reduction at p . \square

2.3. Characteristic classes. We continue to denote by C an algebraically closed field of characteristic 0.

2.3.1. Let X be a proper, connected, smooth C -scheme, equipped with a normal crossings divisor D , and let $U = X - D$. We fix a geometric point $\bar{\eta}$ mapping to the generic point $\eta \in X$. Let G be a finite quotient of $\pi_{1,\text{ét}}(U, \bar{\eta})$. For any i there are canonical maps

$$(2.3.2) \quad H^i(G, \mathbb{F}_p) \rightarrow H^i(\pi_{1,\text{ét}}(U, \bar{\eta}), \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(U, \mathbb{F}_p),$$

where the first map is inflation of classes from G to $\pi_{1,\text{ét}}(U, \bar{\eta})$, and the second map is induced by considering $\pi_{1,\text{ét}}(U, \bar{\eta})$ -representations as étale sheaves on U . In the proposition below we will consider the condition that the composite of the two maps above is surjective. Note that we can always choose G such that the *first map* is surjective. In particular, if U is an étale $K(\pi, 1)$ we can choose G so that the composite is surjective, but this is not the case in general.

Finally for a G -cover $Y \rightarrow U$, let $\text{ed}(Y/U; p)$ denote the p -essential dimension of Y over U [RY00].

Proposition 2.3.3. *Suppose that $i < p - 2$ and that (X, D) has unramified good reduction at p . Let G be a finite group and let $Y \rightarrow U$ be a G -cover. Suppose that $h_{(X,D)}^{0,i} \neq 0$ and that the (restriction of the) classifying map*

$$(2.3.4) \quad H^i(G, \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(U, \mathbb{F}_p)$$

is surjective. Then $\text{ed}(Y/U; p) \geq i$. If X is an abelian variety and $D = \emptyset$, the above holds for $i \leq p - 2$.

Proof. Let $U' \rightarrow U$ be a finite, connected covering which has prime to p degree over U , and let $\eta' \in U'$ be the generic point. We have to show that $\text{ed}(Y'/U') \geq i$, where $Y' = Y \times_U U'$. Consider the composite map

$$H^i(G, \mathbb{F}_p) \rightarrow H^i(\pi_{1,\text{ét}}(U, \bar{\eta}), \mathbb{F}_p) \rightarrow H_{\text{ét}}^i(U, \mathbb{F}_p) \rightarrow H^i(\eta, \mathbb{F}_p) \rightarrow H^i(\eta', \mathbb{F}_p).$$

Our assumptions imply that the composite of the first two maps is surjective. By Corollary 2.2.13, the third map is nonzero, as $h_{(X,D)}^{0,i} \neq 0$. Thus the composite of the first three maps is nonzero. The composite of the fourth map and the trace map $H^i(\eta', \mathbb{F}_p) \rightarrow H^i(\eta, \mathbb{F}_p)$ is multiplication by $\deg(\eta'/\eta)$, hence injective, as η'/η has degree prime to p . Thus the final map is an injection and the composite of all four maps is nonzero.

Suppose $\text{ed}(Y'/U') < i$. Then for some dense open $W \subset U'$, there is a map of C -schemes $W \rightarrow Z$, with $\dim Z < i$, and a G -cover $Y'_Z \rightarrow Z$, such that $Y'|_W \simeq Y'_Z \times_Z W$ as W -schemes with G -action [FKW19, 2.2.2]. Shrinking Z and W if necessary, we may assume that Z is affine. The above constructions, then give us a commutative diagram

$$\begin{array}{ccccc} H^i(G, \mathbb{F}_p) & \longrightarrow & H_{\text{ét}}^i(Z, \mathbb{F}_p) & & \\ \parallel & & \downarrow & & \\ H^i(G, \mathbb{F}_p) & \longrightarrow & H_{\text{ét}}^i(W, \mathbb{F}_p) & \longrightarrow & H_{\text{ét}}^i(\eta', \mathbb{F}_p) \end{array}$$

Since Z is affine of dimension $< i$ it follows that $H_{\text{ét}}^i(Z, \mathbb{F}_p) = 0$. This implies that the composite of the maps in the bottom row is 0. This contradicts what we saw above. \square

Corollary 2.3.5. *Let X/C be an abelian variety of dimension g . Let $p \geq g + 2$, and suppose that X has unramified good reduction at p . Let $X' = X$, viewed as a $(\mathbb{Z}/p\mathbb{Z})^{2g}$ -cover of X via the multiplication by p map $X' \rightarrow X$.*

Then $\text{ed}(X'/X; p) = g$. In particular, this equality holds for almost all p .

Remark 2.3.6. Corollary 2.3.5 resolves almost all of a conjecture of Brosnan [FS21, Conj. 6.1]. Note that [FS21, Cor. 6.7] establishes Brosnan's conjecture for $\dim X \leq 3$ and a positive density set of primes (depending on X), and one can also deduce the conjecture for a sufficiently generic abelian variety from [FS21, Lemma 6.2]. Prior work of Gabber [CT02] established the result for a very general product of elliptic curves.

Proof of Corollary 2.3.5. By definition, $g = \dim X \geq \text{ed}(X'/X; p)$, so it suffices to prove that $\text{ed}(X'/X; p) \geq g$. Let $G = (\mathbb{Z}/p\mathbb{Z})^{2g}$ be the quotient of $\pi_{1,\text{ét}}(X, \bar{\eta})$ corresponding to $X' \rightarrow X$. Note that, in our present situation, the map (2.3.4) is surjective because $H_{\text{ét}}^*(X, \mathbb{F}_p)$ is the exterior algebra on $H_{\text{ét}}^1(X, \mathbb{F}_p)$. Since $h_X^{0,g} = 1$, the inequality $\text{ed}(X'/X; p) \geq g$ follows from Proposition 2.3.3.

The final claim follows as in the proof of Corollary 2.2.15, as X has unramified good reduction at almost all primes p . \square

2.3.7. We now explain a generalization of Corollary 2.3.5, for the mod p homology cover $Y \rightarrow U$. Recall that this is the cover corresponding to the maximal quotient of $\pi_{1,\text{ét}}(U, \bar{\eta})$ which is an elementary abelian p -group. When U is projective this is just the pullback to U of the cover described in Corollary 2.3.5 applied to the Albanese variety of U .

Theorem 2.3.8. *Suppose (X, D) has unramified good reduction at p . Suppose that $i \leq \min\{h_{(X,D)}^{0,1}, \dim X\}$ and that $p > \max\{i+1, 3\}$. Then the mod p homology cover $Y \rightarrow U$ of U , satisfies $\text{ed}(Y/U; p) \geq i$.*

In particular, if $p > \max\{h_{(X,D)}^{0,1} + 1, 3\}$ then

$$\text{ed}(Y/U; p) \geq \min\{h_{(X,D)}^{0,1}, \dim X\}$$

Proof. As in the proof of Proposition 2.3.3, let $U' \rightarrow U$ be a finite, connected covering which has prime to p degree over η , and let $\eta' \in U'$ be the generic point. Let $G = \text{Gal}(Y/U)$, and consider the composite map

$$\wedge^i H^1(G, \mathbb{F}_p) \simeq \wedge^i H_{\text{ét}}^1(U, \mathbb{F}_p) \rightarrow H^i(\eta, \mathbb{F}_p) \rightarrow H^i(\eta', \mathbb{F}_p).$$

By Corollary 2.2.13, the second map is nonzero, and the last map is injective as U' has degree prime to p over η . As the composite map factors through $H^i(G, \mathbb{F}_p)$, it follows that

$$H^i(G, \mathbb{F}_p) \rightarrow H^i(\eta', \mathbb{F}_p)$$

is nonzero, which implies that $\text{ed}(Y/U; p) \geq i$, as in the proof of Proposition 2.3.3. \square

Corollary 2.3.9. *Suppose X is a projective C -scheme which has unramified good reduction at p . Let $b_1 = \dim_{\mathbb{Q}} H^1(X, \mathbb{Q})$ be the first Betti number, and suppose $p > \max\{\frac{1}{2}b_1 + 1, 3\}$. Then the mod p homology cover $Y \rightarrow X$ satisfies*

$$\text{ed}(Y/X; p) \geq \min\{\frac{1}{2}b_1, \dim X\}.$$

Proof. As X is projective, we have $h_X^{0,1} = h_X^{1,0} = \frac{1}{2}b_1$. The corollary now follows from Theorem 2.3.8. \square

3. TORUS EMBEDDINGS AND SHIMURA VARIETIES

In this section, we use the results above to obtain lower bounds on certain coverings that arise naturally in the context of torus embeddings and Shimura varieties. We assume that $p > 2$ throughout this section. In §3.1, we compute the cohomology of certain generalized finite Heisenberg groups. In §3.2 we consider certain torus bundles over abelian varieties. These have covers whose groups are the finite Heisenberg groups, and we use the results of §3.1 and Proposition 2.3.3 to show that these covers are p -incompressible. We then show a variant of this p -incompressibility result, where we restrict covers to analytic (not just Zariski) neighborhoods of the boundary in these torus bundles. Finally, in §3.3, we apply this last result to show p -incompressibility of certain congruence covers.

3.1. Cohomology of generalized Heisenberg groups.

3.1.1. In this subsection, we will be concerned with central extensions of finitely generated abelian groups, which are either free abelian, or elementary abelian p -groups. We begin by considering a central extension of elementary abelian p -groups

$$0 \rightarrow N \rightarrow E \rightarrow H \rightarrow 0.$$

Such extensions are classified by the cohomology group [BC92, §10]

$$H^2(H, N) \simeq H^2(H, \mathbb{F}_p) \otimes N \simeq \wedge^2 H^* \otimes N \oplus H^* \otimes N.$$

Here the term $\wedge^2 H^*$ is the image of $H^1(H, \mathbb{F}_p)^{\otimes 2} = H^{*\otimes 2}$ in $H^2(H, \mathbb{F}_p)$ under the cup product, while the H^* in the second term is the image of the Bockstein map $H^1(H, \mathbb{F}_p) \rightarrow H^2(H, \mathbb{F}_p)$.

Recall that if $c \in Z^2(H, N)$ is a 2-cocycle, the corresponding central extension E is defined by taking the underlying set of E to be $N \times H$ with the group law given by

$$(n_1, h_1) \cdot (n_2, h_2) = (n_1 + n_2 + c(h_1, h_2), h_1 + h_2).$$

We call the extension E a *finite Heisenberg group*² (or just *Heisenberg group* if the context is clear) if its class in $H^2(H, N)$ is represented by an alternating bilinear form $H \otimes H \rightarrow N$.

For an \mathbb{F}_p -vector space V with an action of \mathbb{F}_p^\times , we say that V has weight $n \in \mathbb{Z}/(p-1)\mathbb{Z}$ if $\alpha \in \mathbb{F}_p^\times$ acts by α^{-n} . In general, we denote by $V_n \subset V$ the direct summand of weight n . The following lemma gives a number of characterizations of Heisenberg groups, as well as showing that this property depends only on E as a group, and not as an extension.

Lemma 3.1.2. *The following conditions on the extension E are equivalent:*

- (i) E is a finite Heisenberg group.
- (ii) The class $c \in H^2(H, N)$ defining E is contained in $\wedge^2 H^* \otimes N$.
- (iii) E has exponent p .
- (iv) The extension E admits an action of \mathbb{F}_p^\times , such that H has weight -1 and N has weight -2 .

Proof. (ii) \Rightarrow (i): Suppose c satisfies (ii). Write $c = \sum_{i=1}^k (\alpha_i \cup \beta_i) \otimes n_i \in Z^2(H, N)$, where $\alpha_i, \beta_i \in Z^1(H, \mathbb{F}_p)$ and $n_i \in N$. Then c is also represented by

$$\frac{1}{2} \sum_{i=1}^k (\alpha_i \cup \beta_i - \beta_i \cup \alpha_i) \otimes n_i,$$

which is an N -valued alternating form.

(i) \Rightarrow (iv): Recall that, as a set, E is identified with $N \times H$. Define the \mathbb{F}_p^\times action on E by $\alpha \cdot (n, h) = (\alpha^2 n, \alpha h)$. This evidently induces an action of \mathbb{F}_p^\times on E as a set, and from the explicit description of the group law on E one sees that this action respects the group structure (cf. [Pey08, Prop. 6]).

(iv) \Rightarrow (iii): Consider the multiplication by p map on E . This gives a map $H \rightarrow N$ which commutes with any automorphism of E . In particular it commutes with the \mathbb{F}_p^\times -action, but H and N have distinct weights, so this map is 0.

(iii) \Rightarrow (ii): First note that (i) \Rightarrow (iii), as, if c is alternating, then $c(nh, h) = 0$ for any positive integer n . Now suppose E has exponent p , and let $c \in H^2(H, N)$ be its class. Write $c = a + b$ in (ii) with $a \in \wedge^2 H^* \otimes N$, and $b \in H^* \otimes N$. By what we just saw, the extension E_{-a} corresponding to $-a$ has exponent p . Replacing E by the Baer sum of E_{-a} and E , we may assume that $c \in H^* \otimes N$. If $c \neq 0$, there exist linear forms $s : H^* \rightarrow \mathbb{F}_p$ and $t : N \rightarrow \mathbb{F}_p$ such that $s \otimes t(c) \in \mathbb{F}_p \simeq H^2(\mathbb{F}_p, \mathbb{F}_p)$ is nonzero. If E has exponent p then so does the extension $E_{s,t}$ obtained by pulling back and pushing out E by s and t respectively. However $E_{s,t}$ corresponds to $s \otimes t(c)$ which is in the image of the Bockstein map, and hence $E_{s,t} \simeq \mathbb{Z}/p^2\mathbb{Z}$, which is a contradiction. Hence $c = 0$, which implies (ii). \square

3.1.3. We now change our setup, and consider a central extension

$$(3.1.4) \quad 0 \rightarrow N \rightarrow E \rightarrow H \rightarrow 0$$

²We remark that this is a small abuse of terminology, as this name is often reserved for the case when $\dim_{\mathbb{F}_p} N = 1$.

where N and H are finitely-generated, free abelian pro- p groups. That is $N \simeq \mathbb{Z}_p^r$ and $H \simeq \mathbb{Z}_p^s$ for some r, s .

For a finitely-generated free \mathbb{Z}_p -module V , equipped with an action of \mathbb{Z}_p^\times , we say that V has weight $n \in \mathbb{Z}$ if $\alpha \in \mathbb{Z}_p^\times$ acts by α^{-n} on V . Of course this notion makes sense for any character of \mathbb{Z}_p^\times , but we will not need it in this generality.

Lemma 3.1.5. *For E as in (3.1.4) the following hold:*

- (i) *The class of E in $H^2(H, N)$ is represented by an alternating bilinear form.*
- (ii) *There exists a \mathbb{Z}_p^\times action on E such that H has weight -1 and N has weight -2 .*

Proof. The proof of (i) is analogous to the proof of (ii) \Rightarrow (i) in Lemma 3.1.2, and the proof of (iii) is then analogous to (i) \Rightarrow (iv) in that lemma. \square

Lemma 3.1.6. *The Hochschild-Serre spectral sequence*

$$E_2^{i,j} = H^i(H, H^j(N, \mathbb{Z}_p)) \Rightarrow H^{i+j}(E, \mathbb{Z}_p)$$

degenerates at the E_3 -page. The same holds for the spectral sequence

$$E_2^{i,j} = H^i(H, H^j(N, \mathbb{F}_p)) \Rightarrow H^{i+j}(E, \mathbb{F}_p)$$

if $p \geq \min\{\text{rk}_{\mathbb{Z}} N + 1, \text{rk}_{\mathbb{Z}} H\}$.

Proof. The \mathbb{Z}_p^\times -action on E constructed in Lemma 3.1.5 acts compatibly on the spectral sequences. The cohomology ring $H^*(H, \mathbb{Z}_p)$ (resp. $H^*(H, \mathbb{F}_p)$) is the exterior algebra on $H^1(H, \mathbb{Z}_p) = H^*$, (resp. $H^1(H, \mathbb{F}_p)$) the \mathbb{Z}_p -linear dual of H (cf. [Bro94, Thm. 6.4]). Hence

$$H^i(H, H^j(N, \mathbb{Z}_p)) \simeq H^i(H, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} H^j(N, \mathbb{Z}_p) \simeq \wedge^i H^* \otimes \wedge^j N^*.$$

(An analogous expression holds for the E_2 term of the spectral sequence with \mathbb{F}_p coefficients.) It follows that, in the first spectral sequence, $E_2^{i,j}$ has weight $i + 2j$. For $r \geq 3$, the source and target of the differential $d_r : E_2^{i,j} \rightarrow E_2^{i+r, j-r+1}$ have different weights, hence this differential vanishes. This proves the first claim.

For the proof of the second claim, note that the weights of the terms in the second spectral sequence are elements of \mathbb{F}_p^\times . The source and target of d_r have weights $i + 2j$ and $i + 2j + (2 - r)$ respectively. Thus, if $r = 3, \dots, p$ the same argument as above shows that $d_r = 0$. If $r \geq p + 1$, then our assumption on p forces d_r to be 0, as $E_r^{i,j} = 0$ unless $0 \leq i \leq \text{rk}_{\mathbb{Z}} H, 0 \leq j \leq \text{rk}_{\mathbb{Z}} H$. \square

3.1.7. A reduction mod p of the extension E of (3.1.4) is a map of central extensions

$$\begin{array}{ccccccc} 0 & \longrightarrow & N & \longrightarrow & E & \longrightarrow & H \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \bar{N} & \longrightarrow & \bar{E} & \longrightarrow & \bar{H} \longrightarrow 0 \end{array}$$

which identifies \bar{N} and \bar{H} with N/pN and H/pH respectively, and such that \bar{E} is a finite Heisenberg group.

Below we will repeatedly use that for a finitely-generated, free abelian pro- p group F , and a finitely-generated elementary abelian group M ,

$$H^*(F, M) = (\wedge^* H^1(F, \mathbb{F}_p)) \otimes M.$$

Lemma 3.1.8. *Any central extension E , as in (3.1.4) admits a mod p reduction \bar{E} . The extension class of \bar{E} is uniquely determined by that of E .*

Proof. Let $\bar{N} = N/pN$, $\bar{H} = H/pH$. The natural map $H^1(\bar{H}, \bar{N}) \rightarrow H^1(H, \bar{N})$ is a bijection. Hence the composite

$$\wedge^2 H^1(\bar{H}, \mathbb{F}_p) \otimes \bar{N} \rightarrow H^2(\bar{H}, \bar{N}) \rightarrow H^2(H, \bar{N}) \simeq \wedge^2 H^1(H, \mathbb{F}_p) \otimes \bar{N}.$$

is again a bijection. This implies that the pushout of E by $N \rightarrow \bar{N}$ arises from an extension \bar{E} , whose class lies in the image of the first map above, and this class is uniquely determined by the class of E . \square

Proposition 3.1.9. *Let E be as in (3.1.4), and let \bar{E} be a mod p reduction of E . If $p > \frac{1}{2}(\text{rk}_{\mathbb{Z}} H + 3 \text{rk}_{\mathbb{Z}} N + 1)$, then the map of cohomology rings $H^*(\bar{E}, \mathbb{F}_p) \rightarrow H^*(E, \mathbb{F}_p)$ is surjective.*

Proof. Recall the Hochschild-Serre spectral sequence, which converges to the mod p cohomology of E :

$$E_2^{i,j} = \wedge^i \bar{H}^* \otimes \wedge^j \bar{N}^* = H^i(H, H^j(N, \mathbb{F}_p)) \Rightarrow H^{i+j}(E, \mathbb{F}_p).$$

By Lemma 3.1.6 and our assumptions on p , this spectral E_r degenerates on the E_3 page. There is also the analogous spectral sequence \bar{E}_r converging to $H^{i+j}(\bar{E}, \mathbb{F}_p)$:

$$\bar{E}_2^{i,j} = H^i(\bar{H}, \mathbb{F}_p) \otimes_{\mathbb{F}_p} H^j(\bar{N}, \mathbb{F}_p) = H^i(\bar{H}, H^j(\bar{N}, \mathbb{F}_p)) \Rightarrow H^{i+j}(\bar{E}, \mathbb{F}_p).$$

The cohomology ring $H^*(\bar{H}, \mathbb{F}_p)$ is a tensor product of an exterior and a symmetric algebra:

$$H^*(\bar{H}, \mathbb{F}_p) = \wedge^\bullet \bar{H}^*(1) \otimes \text{Sym}^\bullet \bar{H}^*(2),$$

where $\bar{H}^*(1), \bar{H}^*(2)$, denote the \mathbb{F}_p -dual of \bar{H} considered in degree 1 and 2 respectively, [AM04, II, Cor. 4.2]. One also has the analogous description of $H^*(\bar{N}, \mathbb{F}_p)$. Let $\tilde{E}_2^{i,j} \subset \bar{E}_2^{i,j}$ be the subgroup generated by the image of

$$\wedge^i \bar{E}_2^{1,0} \otimes \wedge^j \bar{E}_2^{0,1} \simeq \wedge^i H^1(\bar{H}, \mathbb{F}_p) \otimes \wedge^j H^1(\bar{N}, \mathbb{F}_p) \simeq \wedge^i H^1(H, \mathbb{F}_p) \otimes \wedge^j H^1(N, \mathbb{F}_p)$$

under the cup product. The inclusion $\tilde{E}_2^{i,j} \subset \bar{E}_2^{i,j}$ induces a splitting of the projection $\bar{E}_2^{\bullet,\bullet} \rightarrow E_2^{\bullet,\bullet}$ as graded groups.

Now consider the differential $d_2 : \bar{E}_2^{0,1} \rightarrow \bar{E}_2^{2,0}$. This is a map $N^* \rightarrow H^2(H, \mathbb{F}_p)$ which we consider as an element $d_2 \in H^2(H, N)$. By [HS53, Thm. 4], $-d_2$ is the class of the extension \bar{E} . Since \bar{E} is a Heisenberg group, Lemma 3.1.2 implies that $d_2 \in N \otimes \wedge^2 H^* \subset H^2(H, N)$, so $d_2(\bar{E}^{0,1}) \subset \bar{E}_2^{2,0}$. Since $\bar{E}_2^{\bullet,\bullet}$ is a differential graded algebra, this implies that $\tilde{E}_2^{\bullet,\bullet}$ is stable under d_2 . Denote the cohomology of $(\tilde{E}_2^{\bullet,\bullet}, d_2)$ by $\tilde{E}_3^{\bullet,\bullet}$. Then $\tilde{E}_3^{\bullet,\bullet} \subset \bar{E}_3^{\bullet,\bullet}$ induces a splitting of the projection $\bar{E}_3^{\bullet,\bullet} \rightarrow E_3^{\bullet,\bullet}$, as graded groups, so again the latter map is surjective.

To prove the proposition it suffices to show that for $r \geq 2$, and $0 \leq i \leq \text{rk}_{\mathbb{Z}} H$, $0 \leq j \leq \text{rk}_{\mathbb{Z}} N$, the map $\bar{E}_r^{i,j} \rightarrow E_r^{i,j}$ is surjective, as the target of this map is trivial for i, j outside this range. Thus we assume from now on that i, j satisfy these inequalities.

We now make use of the \mathbb{F}_p^\times -action on \bar{E} given by Lemma 3.1.2. This induces an \mathbb{F}_p^\times -action on the spectral sequence \bar{E}_r . The description of the cohomology rings $H^*(\bar{H}, \mathbb{F}_p), H^*(\bar{N}, \mathbb{F}_p)$ implies that the weights of $\bar{E}_2^{i,j}$ are all represented by integers in the interval $[\frac{1}{2}(i+2j), i+2j]$. Our assumptions on i, j and p imply that this interval has length less than $p-1$, i.e. the set of integers in it maps injectively to $\mathbb{Z}/(p-1)\mathbb{Z}$. In particular, the weight $i+2j$ piece $(\bar{E}_2^{i,j})_{i+2j} \subset \bar{E}_2^{i,j}$ is precisely $\tilde{E}_2^{i,j}$, and so $\tilde{E}_3^{i,j} = (E_3^{i,j})_{i+2j}$.

Now for $r \geq 3$, consider the differential $d_r : \bar{E}_r^{i,j} \rightarrow \bar{E}_r^{i+r,j-r+1}$. The weights appearing in $\bar{E}_r^{i+r,j-r+1}$ are represented by integers in $[\frac{1}{2}(i+2j-r+2), i+2j-r+2]$. Our assumptions on i, j , and p imply that the interval $[\frac{1}{2}(i+2j-r+2), i+2j]$ has length less than $p-1$ for $r = 3, \dots, j+1$. This implies that $(\bar{E}_r^{i+r,j-r+1})_{i+2j} = 0$, and so $d_r|_{(\bar{E}_r^{i,j})_{i+2j}} = 0$. On the other hand for $r > j+1$ $d_r|_{\bar{E}_r^{i,j}} = 0$ as the target of this map is trivial.

We now show by induction on r that $(\bar{E}_r^{i,j})_{i+2j}$ maps isomorphically to $E_r^{i,j}$, which in particular implies that $\bar{E}_r^{i,j} \rightarrow E_r^{i,j}$ is surjective, finishing the proof of the proposition. We have already seen this for $r = 3$. Assume the result for some $r \geq 3$. As $d_r|_{(\bar{E}_r^{i,j})_{i+2j}} = 0$ it follows that

$$(\bar{E}_r^{i,j})_{i+2j} \twoheadrightarrow H(\bar{E}_r)^{i,j}_{i+2j} \simeq (\bar{E}_{r+1}^{i,j})_{i+2j} \rightarrow E_{r+1}^{i,j} \simeq E_r^{i,j},$$

where the final isomorphism follows from Lemma 3.1.6. The composite is the natural map, which we are assuming is an isomorphism. Hence all the maps above are isomorphisms, which completes the induction. \square

3.2. Toric varieties.

3.2.1. Throughout this section we work over an algebraically closed field C . Let T be a torus over C , S an abelian variety over C , and U/S a T -torsor. Fix a geometric point \bar{x} of U , and let \bar{s} be its image in S . We will apply the results of the previous section to U . For a profinite group G , denote by G^p its pro- p completion.

Lemma 3.2.2. *The C -scheme U is an étale $K(\pi, 1)$, and $\pi_1(U, \bar{x})$ is a central extension of free abelian profinite groups*

$$(3.2.3) \quad 0 \rightarrow \pi_{1,\text{ét}}(U_{\bar{s}}, \bar{x}) \rightarrow \pi_{1,\text{ét}}(U, \bar{x}) \rightarrow \pi_{1,\text{ét}}(S, \bar{s}) \rightarrow 0,$$

whose pro- p completion is an extension

$$(3.2.4) \quad 0 \rightarrow \pi_{1,\text{ét}}(U_{\bar{s}}, \bar{x})^p \rightarrow \pi_{1,\text{ét}}(U, \bar{x})^p \rightarrow \pi_{1,\text{ét}}(S, \bar{s})^p \rightarrow 0.$$

The natural map $H^*(\pi_{1,\text{ét}}(U, \bar{x})^p, \mathbb{F}_p) \rightarrow H^*(\pi_{1,\text{ét}}(U, \bar{x}), \mathbb{F}_p)$ is an isomorphism.

Proof. Note that S and $U_{\bar{s}}$ are étale $K(\pi, 1)$'s. Indeed, for each of these varieties, both their cohomology and the cohomology of the respective étale fundamental group are exterior algebras on their cohomology in degree 1. Now the fact that U is a $K(\pi, 1)$ follows by comparing the Hochschild-Serre spectral sequence for $\pi_{1,\text{ét}}(U_{\bar{s}}, \bar{x}) \subset \pi_{1,\text{ét}}(U, \bar{x})$ and the Leray spectral sequence for the map $U \rightarrow S$.

It is well known that $\pi_{1,\text{ét}}(U, \bar{x})$ is an extension as in 3.2.3. That this is a central extension follows from the fact that the T -torsor U is Zariski locally trivial (by Hilbert's Theorem 90).

That the pro- p completion of (3.2.3) is an extension as in (3.2.4) follows easily from the fact that any central extension of two finite abelian groups of coprime order is trivial. It follows that the kernel of $\pi_{1,\text{ét}}(U, \bar{x}) \rightarrow \pi_{1,\text{ét}}(U, \bar{x})^p$ is a prime to p profinite group, which implies the final claim. \square

3.2.5. Keep the assumptions above, and consider the extension (3.2.4). We have the notion of a mod p reduction of such an extension given in 3.1.3. In particular, such a mod p reduction gives rise to a surjection of $\pi_{1,\text{ét}}(U, \bar{x}) \twoheadrightarrow \bar{E}$ onto a finite Heisenberg group \bar{E} . We write $U(\bar{E}) \rightarrow U$ for the finite cover corresponding to \bar{E} .

3.2.6. We again denote by T the torus over \mathbb{Z} with character group $X^*(T)$. We say $U \rightarrow S$ has *good reduction at p* if there exists a p -adic valuation on C , with ring of integers \mathcal{O}_C , such that $U \rightarrow S$ extends to a T -torsor over an abelian scheme over \mathcal{O}_{C° , $U^\circ \rightarrow S^\circ$. We say $U \rightarrow S$ has *unramified good reduction* if $U^\circ \rightarrow S^\circ$ can be chosen so that it descends to a T -torsor over an absolutely unramified discrete valuation ring $\mathcal{O} \subset \mathcal{O}_C$.

Proposition 3.2.7. *Suppose that $p > \dim U + \frac{1}{2}(\dim U_{\bar{s}} + 1)$, and that $U \rightarrow S$ has unramified good reduction at p . Then $\text{ed}(U(\bar{E})/U; p) = \dim U$.*

Proof. Since U has unramified good reduction, there is a p -adic valuation on C , and an absolutely unramified discrete valuation ring $\mathcal{O} \subset \mathcal{O}_C$, such that U descends to a T -torsor over an abelian scheme $U^\circ \rightarrow S^\circ$ over \mathcal{O} . Fix a basis for $X^*(T)$. Then U° corresponds to a collection of line bundles $\mathcal{L}_1, \dots, \mathcal{L}_t$ over S° , where $t = \dim U_{\bar{s}}$. Let $P_i^\circ = \text{Proj}_{\mathcal{O}_{S^\circ}}(\mathcal{O}_{S^\circ} \oplus \mathcal{L}_i^\vee)$, and set $P^\circ = P_1^\circ \times_{S^\circ} P_2^\circ \times_{S^\circ} \dots \times_{S^\circ} P_t^\circ$. Then P° is a smooth projective scheme S° -scheme, and $D^\circ = P^\circ - U^\circ$ is a normal crossings divisor. We set $P = P_C^\circ$ and $D = D_C^\circ$.

Let $d = \dim U$. Fix an identification $T \simeq \mathbb{G}_m^t$, and let z_1, \dots, z_t be the standard co-ordinates on \mathbb{G}_m^t . By Hilbert's Theorem 90, Zariski locally on S , we can identify U with $S \times T \simeq S \times \mathbb{G}_m^t$. The differential $\omega_T = \frac{dz_1}{z_1} \wedge \dots \wedge \frac{dz_t}{z_t}$ is T -invariant, and hence does not depend on the identification $T \simeq U$. (It is independent of our fixed isomorphism $T \simeq \mathbb{G}_m^t$ up to a sign). It follows that ω_T gives rise to a global section of $\Gamma(P, \Omega_P^t(\log D))$. Let $r = \dim S$, and $\omega_r \in \Gamma(S, \Omega_S^r)$ a nonzero r -form. Then $\omega_r \otimes \omega_t \in \Gamma(P, \Omega_P^d(\log D))$ is nonzero, so $h_{(P,D)}^{0,d} \neq 0$.

Now consider the maps

$$H^d(\bar{E}, \mathbb{F}_p) \rightarrow H^d(\pi_1(U, \bar{x})^p, \mathbb{F}_p) \rightarrow H^d(\pi_1(U, \bar{x}), \mathbb{F}_p) \rightarrow H^d(U, \mathbb{F}_p).$$

The first map is surjective by Proposition 3.1.9, and the second and third maps are isomorphisms by Lemma 3.2.2. As $h_{(P,D)}^{0,d} \neq 0$, the Proposition follows from Proposition 2.3.3. \square

3.2.8. We want to prove a variant of Proposition 3.2.7 which will be used in the next subsection to show p -incompressibility of certain coverings of Shimura varieties. To explain it, we recall some facts about families of toric varieties [Ful93], [KKMSD73], [Pin90, §5]. Note that the family $P \rightarrow S$, which appeared in the proof of Proposition 3.2.7 is an example of such a family.

Let k be a field. A toric variety, or torus embedding X over k is a normal k -scheme X , equipped with an action of a split torus T , such that X admits a dense open T -orbit U , which is a T -torsor. These can be described in terms of *fans*, which are certain collections Σ of *convex polyhedral cones* in $X_*(T)_\mathbb{Q}$. In particular, proper toric varieties correspond to *complete fans*, which are certain decompositions of $X_*(T)_\mathbb{Q}$ into convex polyhedral cones.

Since we are assuming that T is a split torus, it extends canonically to a torus over \mathbb{Z} , which we again denote by T . Thus, a fan Σ actually defines a torus embedding $X(\Sigma)$ over \mathbb{Z} . That is, $X(\Sigma)$ is a normal scheme, equipped with an action of T , containing a dense T -torsor. The fiber of $X(\Sigma)$ over any point $\text{Spec } k \rightarrow \text{Spec } \mathbb{Z}$ is the toric variety over k corresponding to Σ .

We will need the relative version of this notion. Let S be a scheme and $X \rightarrow S$ a map of schemes, equipped with an action of T . Then X/S is called a torus embedding over S if, Zariski locally on S , $X \rightarrow S$ can be T -equivariantly identified

with $X(\Sigma) \times_{\mathbb{Z}} S$ for a fan Σ . In this case there is an open subset $U \subset X$, which is a T -torsor, and which is dense in the fiber over every point of S . The complement $D = X - U$, is called the *boundary* of X .

We remark that it may appear more natural to make this definition with the condition on $X \rightarrow S$ imposed only étale locally on S . However these two notions are the same: As $X(\Sigma)$ contains an open dense T -torsor $U(\Sigma)$, $\text{Aut}_T(X(\Sigma) \times_{\mathbb{Z}} S) = \text{Aut}_T(U(\Sigma) \times S) = T(S)$. Hence any $X \rightarrow S$ which is étale locally isomorphic to $X(\Sigma) \times S$ gives rise to an étale T -torsor. As in the proof of Proposition 3.2.7, such a torsor is Zariski locally trivial.

Lemma 3.2.9. *Suppose that S is connected, and fix T as above. Then there is an equivalence of categories between torus embeddings $X \rightarrow S$ containing a dense open T -torsor, and pairs (U, Σ) , where U is a T -torsor over S , and Σ is a fan in $X_*(T)_{\mathbb{Q}}$.*

Proof. Given $X \rightarrow S$, we can associate to it the open T -torsor $U \subset X$, and the fan Σ corresponding to the torus embedding X_{η} , where $\eta \in S$ is the generic point. Conversely, given (U, Σ) , Σ defines a toric variety X_0 over C , which is even defined over \mathbb{Z} . We take $X = (X_0 \times U)/T$. One checks easily that these two constructions are quasi-inverses. \square

3.2.10. Now suppose that S is a C -scheme, and let $X \rightarrow S$, U and D be as above. Then D has a stratification $D_0 \subset D_1 \subset D_2 \dots$, which may be described as follows: Zariski locally, $X = X_0 \times S$ as torus embeddings, where X_0 is a torus embedding over C . We set $D_j \subset D$ to be the product of S and the closure of the j -dimensional T -orbits in X_0 . Then D_0 is étale over S , and, if X is proper, it is finite étale.

Let $X^1 \rightarrow X$ be the blow up of D_0 on X . Since D_0 is fixed by T , X^1 is again a torus embedding over S , and we denote its boundary by D^1 . If X is smooth over S , with D a relative normal crossings divisor, then the same is true for X^1 and D^1 . We can continue this construction to obtain a sequence of blow ups $X \leftarrow X^1 \leftarrow X^2 \dots$, with boundary $D^i \subset X^i$. Finally, we denote by \widehat{X}_i the completion of X along D^i .

The following lemma implies that if (X, D) has unramified good reduction at p , then so does (X^1, D^1) :

Lemma 3.2.11. *Suppose that X/S is smooth, and that D is a normal crossings divisor. If the T -torsor U has unramified good reduction at p , then (X, D) has unramified good reduction at p .*

Proof. Suppose U/S descends to a T -torsor U° over a discrete valuation ring $\mathcal{O} \subset C$. By Lemma 3.2.9, as a torus embedding X is determined by (U, Σ) for some fan Σ . Then (U°, Σ) determines a torus embedding X° over \mathcal{O} . The property that X is smooth and D is a normal crossings divisor can be read off from the fan Σ [KKMSD73, II.2, Thm. 4*]. It implies that X° is smooth over \mathcal{O} , and that its boundary D° is a relative normal crossings divisor. \square

Lemma 3.2.12. *Suppose that S and X/S are smooth, with $D \subset X$ a relative normal crossings divisor. Let $Z \subset \widehat{X}$ be a formal subscheme of codimension 1, and $Z^i \subset \widehat{X}^i$ its proper transform. Then there exists a dense open subset $W \subset S$ and an integer $i \geq 1$ such that $D_0^i|_W \not\subset Z^i$.*

Proof. Let $\pi_i : \widehat{X}^i \rightarrow \widehat{X}$. Recall that the proper transform Z^i is defined to be the union of those components of $\pi_i^{-1}(Z)$ which are not contained in $\pi_i^{-1}(D_0)$. It suffices

to show the lemma with S replaced by one of its generic points, so we may assume that $S = \text{Spec } \kappa$ is a field. By induction on the number of irreducible components of Z , we may assume that Z is an irreducible Cartier divisor in X .

Let $x \in D^0$, and let z_1, \dots, z_n be a system of local co-ordinates at x , such that the ideal of D at x is given by $z_1 z_2 \dots z_n$. Then in a formal neighborhood of x , the ideal of Z is generated by a nonzero power series $f = \sum_I a_I z^I$, where $I = (i_1, \dots, i_n)$ runs over n -tuples of non-negative integers, and $a_I \in \kappa$. Choose such an n -tuple $J = (j_1, \dots, j_n)$ such that $a_J \neq 0$ and $|J| = j_1 + \dots + j_n$ is as small as possible. If $|J| = 0$, then $x \notin Z$, and the lemma holds without blowing up X . In general we proceed by induction on $|J|$.

We may assume without loss of generality that $j_1 \neq 0$. There is a point $x^1 \in D_0^1 \subset \widehat{X}^1$ such that the functions $z_1, u_2, u_3, \dots, u_n$, with $u_i = \frac{z_i}{z_1}$ are a system of local co-ordinates at x^1 , and the ideal of D^1 is generated by $z_1 u_2 \dots u_n$. The ideal of the proper transform $Z^1 \subset \widehat{X}^1$ is generated by the function $z_1^{-|J|} f$, whose expansion contains the term $a_J u_2^{j_2} \dots u_n^{j_n}$. Thus the result follows by induction on $|J|$. \square

3.2.13. We now assume that $C = \mathbb{C}$, the complex numbers, and we denote by X^{an} the complex analytic space associated to X . Recall that a *Zariski closed* subset of a complex analytic space Y is a closed subset $Z \subset Y$, which is locally defined by (local) analytic functions on Y . We call the complement of a Zariski closed analytic subset a *Zariski open* subset of Y . Note that if $Y \rightarrow W$ is an open embedding of complex analytic spaces, it is not in general the case that a Zariski open $U \subset Y$ is a Zariski open of W ; this is only the case if the closed analytic subset $Z \subset Y$ is of the form $Z = Y \cap Z'$ for some closed analytic subset $Z' \subset W$.

Proposition 3.2.14. *Let S be an abelian scheme and let X/S be a proper torus embedding with boundary $D \subset X$, and dense open T -torsor $U \subset X$. Let $d = \dim X$, and suppose that U/S has unramified good reduction at a prime $p > d + 2$.*

Let $V \subset X^{\text{an}}$ be an analytic open subset containing D^{an} , and $V' \subset V$ a non-empty Zariski open subset of V . Then the map

$$H^d(U, \mathbb{F}_p) \rightarrow H^d(U \cap V', \mathbb{F}_p)$$

is nonzero.

Proof. After replacing X by a blow up, we may assume that X/S is smooth and $D \subset X$ is a normal crossings divisor. Such blow ups are obtained by subdividing the rational cone decomposition of $X_*(T)_{\mathbb{Q}}$ defining X , see [Ful93, p 48]. Then (X, D) has unramified good reduction at p by Lemma 3.2.11.

Next the same proof as in Proposition 3.2.7 shows that $h_{(X,D)}^{0,d} \neq 0$. In fact the differential $\omega_r \otimes \omega_s$ defined there is in $H^0(X, \Omega_X^d(\log D))$ for any torus embedding (X, D) with open dense T -torsor $U \subset X$.

We may assume that $Z = V - V'$ has codimension 1 in V . Let $\widehat{Z} \subset \widehat{X}$ denote the formal completion of Z along D . We apply Lemma 3.2.12 to \widehat{Z} . Then, after replacing X by a blow up and Z by its proper transform, we may assume there is a non-empty Zariski open $W \subset S$ such that $D_0|_W \not\subset Z$. Shrinking W if necessary, since D_0 is étale over S , we may assume that there is an irreducible component D'_0 of D_0 which does not meet Z , and that $X|_W = W \times X_0$ for a toric variety X_0 over \mathbb{C} . Then D'_0 has the form $W \times x_0$ for a T -fixed point $x_0 \in X_0$. We write $U_0 \subset X_0$ for the open T -orbit.

By [Loj64, Thm. 4], there exists a compact subset $W^- \subset W$ such that the inclusion induces a homotopy equivalence. Let $Y_0 \subset X_0$ be a ball around x_0 which is small enough that $Y_0 \setminus D_0 \rightarrow U_0$ is a homotopy equivalence, $W^- \times Y_0 \subset V$, and $W^- \times Y_0$ does not meet Z . Here we are using the compactness of W^- for the second and third properties. Let $Y^* = (W^- \times Y_0) \setminus D \simeq W^- \times (Y_0 \setminus D_0)$. Then $Y^* \rightarrow U_{W^-} \simeq W^- \times U_0$ is a homotopy equivalence, and $Y^* \subset U \cap V'$. Now consider the composite

$$H^d(U, \mathbb{F}_p) \rightarrow H^d(U|_W, \mathbb{F}_p) \rightarrow H^d(U|_{W^-}, \mathbb{F}_p) \rightarrow H^d(Y^*, \mathbb{F}_p).$$

Since $h_{(X,D)}^{0,d} \neq 0$, and (X, D) has unramified good reduction at p , we may apply Corollary 2.2.13 to deduce that the first map is nonzero. The other two maps are induced by homotopy equivalences, hence are isomorphisms. Thus the composite map is nonzero. However, as $Y^* \subset U \cap V'$, the composite map factors through $H^d(U \cap V', \mathbb{F}_p)$ which implies the Lemma. \square

3.2.15. We are nearly ready to show the variant of Proposition 3.2.7 that will be used in the next subsection. To formulate it we need a notion of essential dimension at p for complex analytic spaces. This is defined in a similar way as for algebraic varieties, but there is an important difference in that we do not insist the auxiliary coverings of order prime to p are étale. Unlike the algebraic situation, one cannot reduce ramified coverings to the unramified case, because the notion of Zariski open subsets of complex analytic spaces is not transitive.

Let $V_1 \rightarrow V$ be a finite map of (reduced) complex analytic spaces. The *essential dimension* $\text{ed}(V_1/V)$ is the smallest integer e such that for some dense Zariski open $V' \subset V$, there exists a finite map of analytic spaces $Y_1 \rightarrow Y$, and a map $V' \rightarrow Y$ such that $V_1|_{V'} \rightarrow V'$ is isomorphic to the normalization of $V' \times_Y Y_1$. The *p-essential dimension* $\text{ed}(V_1/V_2; p)$ is defined as the minimum value of $\text{ed}(\tilde{V}_1/\tilde{V})$, where $\tilde{V} \rightarrow V$ runs over finite maps of degree prime to p , and \tilde{V}_1 is the normalization of $\tilde{V} \times_V V_1$. Note that the map $\tilde{V} \rightarrow V$ is finite flat over some Zariski open; its degree is defined as the degree over any such Zariski open.

3.2.16. Keep the notation of Proposition 3.2.14, and consider a surjection onto a finite Heisenberg group $\pi_{1,\text{ét}}(U, \bar{x}) \rightarrow \bar{E}$, as in 3.2.5. We denote by $X(\bar{E}) \rightarrow X$ the normalization of $U(\bar{E}) \rightarrow U$.

Corollary 3.2.17. *With the assumptions of Proposition 3.2.14, suppose that $p > \dim U + \frac{1}{2}(\dim U_{\bar{s}} + 1)$. Write $V(\bar{E}) := X(\bar{E})^{\text{an}}|_V$. Then $\text{ed}(V(\bar{E})/V; p) = d$.*

Proof. Let $\pi : \tilde{V} \rightarrow V$ be a finite covering of degree prime to p , and $\tilde{V}' \subset \tilde{V}$ a Zariski open. We claim that the map $H^d(U, \mathbb{F}_p) \rightarrow H^d(\tilde{V}'|_U, \mathbb{F}_p)$ is nonzero.

The complement $\tilde{Z} = \tilde{V} \setminus \tilde{V}'$, is a Zariski closed subspace of everywhere positive codimension. Hence $\pi(\tilde{Z}) \subset V$ is Zariski closed with everywhere positive codimension. Let $V' = V \setminus \pi(\tilde{Z})$. We may replace \tilde{V}' by the preimage of V' , and assume that π restricts to $\tilde{V}' \rightarrow V'$. Now consider the composite

$$H^d(U, \mathbb{F}_p) \rightarrow H^d(U \cap V', \mathbb{F}_p) \rightarrow H^d(\tilde{V}'|_U, \mathbb{F}_p).$$

The first map is nonzero by Proposition 3.2.14, and the second map is injective, as in the proof of Proposition 2.3.3, hence the claim. As in the proof of Proposition 3.2.7, it follows that the composite

$$H^d(\bar{E}, \mathbb{F}_p) \rightarrow H^d(U, \mathbb{F}_p) \rightarrow H^d(\tilde{V}'|_U, \mathbb{F}_p)$$

is nonzero.

Write $\tilde{V}'(\bar{E})$ for the normalization of $V(\bar{E}) \times_V \tilde{V}'$. If $\tilde{V}'(\bar{E}) \rightarrow \tilde{V}'$ is the normalized pullback of an \bar{E} -covering $Y_1 \rightarrow Y$ of dimension $< d$, then as in the proof of Proposition 2.3.3, the map $H^d(\bar{E}, \mathbb{F}_p) \rightarrow H^d(\tilde{V}'|_U, \mathbb{F}_p)$ factors through $H^d(Y, \mathbb{F}_p)$. Shrinking Y and \tilde{V}' as necessary, we can assume that Y is Stein. By Andreotti-Frankel [AF59], $\dim Y < d$ implies that $H^d(Y, \mathbb{F}_p) = 0$. This gives a contradiction, and proves the corollary. \square

Corollary 3.2.18. *Keep the assumptions of Corollary 3.2.17, but suppose that $\dim S = 0$. Then the conclusion of 3.2.17 holds with no restriction on p .*

Proof. The restriction on p in the proof of 3.2.17 come from the application of Corollary 2.2.13 and Proposition 3.1.9, which guarantee that the maps

$$H^d(\bar{E}, \mathbb{F}_p) \rightarrow H^d(U, \mathbb{F}_p) \rightarrow H^d(U|_W, \mathbb{F}_p)$$

are injective for $W \subset S$ dense open. When $\dim S = 0$, the second map is vacuously bijective, and the first map is surjective, because in this case $\pi_{1,\text{ét}}(U, \bar{x})$ is abelian, $\bar{E} = \pi_{1,\text{ét}}(U, \bar{x})/p\pi_{1,\text{ét}}(U, \bar{x})$, and the cohomology ring $H^*(U, \mathbb{F}_p)$ is generated in degree 1. \square

3.2.19. We remark that Corollary 3.2.18 is originally due to Burda [Bur12], and our proof in this case reduces to a variant of his. The key point in the argument is the construction of the subset Y^* in the proof of Proposition 3.2.14. Burda does this by considering annuli in U with carefully chosen radii, rather than by using blow ups; see [Bur12, Thm. 18].

3.3. Shimura Varieties.

3.3.1. Let (G, X) be a Shimura datum [Del79, §1]. Recall that this consists of a reductive group G over \mathbb{Q} , together with a $G(\mathbb{R})$ -conjugacy class, X , of homomorphisms $h : \mathbb{S} = \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m \rightarrow G$ satisfying certain conditions. These imply, in particular, that X is a Hermitian domain, and that for any neat compact open $K \subset G(\mathbb{A}_f)$, the quotient

$$\text{Sh}_K(G, X) = G(\mathbb{Q}) \backslash X \times G(\mathbb{A}_f) / K$$

has the structure of a complex algebraic variety. Here \mathbb{A}_f denote the finite adeles over \mathbb{Q} .

For $h \in X$, define a G -valued cocharacter μ_h over \mathbb{C} , as follows. For a \mathbb{C} -algebra R , we have $R \otimes_{\mathbb{R}} \mathbb{C} = R \oplus c^*(R)$, where c denotes complex conjugation. The first factor gives an inclusion $R^\times \subset (R \otimes_{\mathbb{R}} \mathbb{C})^\times$, which gives a map $\mathbb{G}_m \rightarrow \mathbb{S}$ over \mathbb{C} , and μ_h is the composite of this map and h . The cocharacter μ_h is miniscule [Del79, 1.2.2], and the axioms for a Shimura variety imply that it is nontrivial.

3.3.2. Now consider the Dynkin diagram $\Delta(G)$ of G , which is equipped with an action of $\text{Gal}(\mathbb{Q}/\mathbb{Q})$. To simplify the discussion, we assume from now on that G^{ad} is \mathbb{Q} -simple, so that the $\text{Gal}(\mathbb{Q}/\mathbb{Q})$ -action on $\Delta(G)$ is *transitive*. The subgroup of $\text{Gal}(\mathbb{Q}/\mathbb{Q})$ that acts on $\Delta(G)$ trivially corresponds to a field K_Δ that is either CM or a totally real field. The action of complex conjugation $c \in \text{Gal}(K_\Delta/\mathbb{Q})$ is given by the *opposition involution* of the root system of G . Recall that this is given by $(-1) \circ w_0$, where w_0 is the longest element of the Weyl group W_G . Thus K_D is a totally real field exactly when $-1 \in W_G$.

Fix a maximal torus $T \subset G^{\text{ad}}$ and a set of positive roots $\Delta^+ \subset X^*(T)$ for G . The vertices of $\Delta(G)$ correspond to the simple roots in Δ^+ . For $\alpha \in \Delta(G)$, let $\mu_\alpha \in X_*(T)$ be the cocharacter which takes the value 1 on α and vanishes otherwise. For a subset $R \subset \Delta(G)$, define $\mu_R \in X_*(T)$ by $\mu_R := \prod_{\alpha \in R} \mu_\alpha$. If we do not specify Δ^+ , then the conjugacy class of μ_R is still well defined, and we will denote it by $[\mu_R]$.

Let $P_R \subset G$ be the parabolic associated to μ_R . Recall that P_R is characterized by the condition that its Lie algebra $\text{Lie } P_R$ is the sum of root spaces on which μ_R is non-negative [DG70, XXVI, 1.4]. As for μ_R , if we do not specify Δ^+ then the conjugacy class of P_R is still well defined, and we denote it by $[P_R]$.

Since μ_h is miniscule, it corresponds to a collection of vertices $\Sigma \subset \Delta(G)$, with each component of $\Delta(G)$ containing at most one element of Σ . Let $\tilde{\Sigma} = \Sigma \cup c(\Sigma)$.

Lemma 3.3.3. *Let $\mu_{\tilde{\Sigma}}$ be in $[\mu_{\tilde{\Sigma}}]$, $P_{\tilde{\Sigma}}$ the associated parabolic and $U_{\tilde{\Sigma}} \subset P_{\tilde{\Sigma}}$ the unipotent radical of $P_{\tilde{\Sigma}}$. Then $U_{\tilde{\Sigma}}$ is a central extension of additive groups*

$$0 \rightarrow Z(U_{\tilde{\Sigma}}) \rightarrow U_{\tilde{\Sigma}} \rightarrow U_{\tilde{\Sigma}}/Z(U_{\tilde{\Sigma}}) \rightarrow 0,$$

and $U_{\tilde{\Sigma}} = Z(U_{\tilde{\Sigma}})$ if c fixes Σ . The cocharacter $w_{\tilde{\Sigma}} = \mu_\Sigma \mu_{c(\Sigma)}$ acts with weight 2 on $Z(U_{\tilde{\Sigma}})$ and weight 1 on $U_{\tilde{\Sigma}}/Z(U_{\tilde{\Sigma}})$.

Proof. Write $G_{\mathbb{R}}^{\text{ad}} = \prod_{i=1}^r G_{\mathbb{R},i}^{\text{ad}}$, where each factor $G_{\mathbb{R},i}^{\text{ad}}$ is absolutely simple. For each i the Dynkin diagram of $G_{\mathbb{R},i}^{\text{ad}}$ corresponds to a component $\Delta(G)_i$ of $\Delta(G)$, and it suffices to prove the statement of the Lemma with $U_{\tilde{\Sigma},i} = U_{\tilde{\Sigma}} \cap G_{\mathbb{C},i}^{\text{ad}}$ in place of $U_{\tilde{\Sigma}}$. We set $\Sigma_i = \Sigma \cap \Delta(G)_i$ and $\tilde{\Sigma}_i = \tilde{\Sigma} \cap \Delta(G)_i$.

The root spaces which appear in $\text{Lie } U_{\tilde{\Sigma}}$ are exactly those which correspond to roots whose expressions as a sum of simple roots in Δ^+ contain an element of $\tilde{\Sigma}$. If Σ_i is empty, then $U_{\tilde{\Sigma},i} = 0$, and there is nothing to prove. Suppose that Σ_i is non-empty.

If c fixes Σ_i then $\tilde{\Sigma}_i = \Sigma_i$ and μ_{Σ_i} is nonzero on exactly one simple root in $\Delta(G)_i$. As μ_h is miniscule, so is μ_{Σ_i} , so the root in Σ_i can appear in the expression for a root in Δ^+ with multiplicity at most 1. This implies that $U_{\tilde{\Sigma},i}$ is abelian, and μ_{Σ_i} acts with weight 1 on $U_{\tilde{\Sigma},i}$, so $\mu_{\Sigma_i} \mu_{c(\Sigma_i)} = \mu_{\tilde{\Sigma}_i}^2$ acts with weight 2.

Now suppose that c does not fix Σ_i . Then $\tilde{\Sigma}_i$ has exactly two elements and $\mu_{\tilde{\Sigma}_i}$ is nonzero on the two corresponding simple roots, which are exchanged by c . If $\alpha \in \Sigma_i$, and e_α is the corresponding root, then e_α and $e_{c(\alpha)}$ both appear in the expression for the longest root in Δ^+ as a sum of simple roots. Using this, one sees that the root spaces appearing in $\text{Lie } Z(U_{\tilde{\Sigma},i})$ correspond to $\beta \in \Delta^+$ whose expression as a sum of simple roots contains both α and $c(\alpha)$, and that $[U_{\tilde{\Sigma}_i}, U_{\tilde{\Sigma}_i}] \subset Z(U_{\tilde{\Sigma}_i})$. Moreover the description of the root spaces appearing in the Lie algebras of $U_{\tilde{\Sigma}_i}$ and $Z(U_{\tilde{\Sigma}_i})$ implies the claim about the weights of $\mu_{\Sigma_i} \mu_{c(\Sigma_i)} = \mu_{\tilde{\Sigma}_i}$. \square

3.3.4. We will be interested in the following condition

$$(3.3.5) \quad \text{The conjugacy class } [\mu_{\tilde{\Sigma}}] \text{ contains a cocharacter defined over } \mathbb{Q}.$$

Note that this condition implies that $[\mu_{\tilde{\Sigma}}]$ is fixed by $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$. We can rephrase this condition in terms of the parabolic $P_R \subset G_{\mathbb{C}}$ associated to μ_R .

Lemma 3.3.6. *The conjugacy class $[\mu_{\tilde{\Sigma}}]$ contains a cocharacter defined over \mathbb{Q} if and only if $[P_{\tilde{\Sigma}}]$ contains a parabolic defined over \mathbb{Q} .*

Proof. If $[\mu_{\tilde{\Sigma}}]$ contains a cocharacter $\mu_{\tilde{\Sigma},\mathbb{Q}}$ defined over \mathbb{Q} , then the subspace of $\text{Lie } G$ on which $\mu_{\tilde{\Sigma},\mathbb{Q}}$ is non-negative is the Lie algebra of a parabolic in $[P_{\tilde{\Sigma}}]$, which is defined over \mathbb{Q} .

Conversely if $[P_{\tilde{\Sigma}}]$ contains a parabolic $P_{\tilde{\Sigma},\mathbb{Q}}$ defined over \mathbb{Q} , then $P_{\tilde{\Sigma},\mathbb{Q}}$ is associated (as above) to some cocharacter $\mu_{\tilde{\Sigma}}$ (not necessarily defined over \mathbb{Q}) in $[\mu_{\tilde{\Sigma}}]$. Let $U_{\tilde{\Sigma},\mathbb{Q}} \subset P_{\tilde{\Sigma},\mathbb{Q}}$ denote the unipotent radical. As $P_{\tilde{\Sigma},\mathbb{Q}}$ is its own normalizer, $\mu_{\tilde{\Sigma}}$ is determined up to conjugation by points of $P_{\tilde{\Sigma},\mathbb{Q}}$. Hence the conjugacy class of $\mu_{\tilde{\Sigma}}$ as a $P_{\tilde{\Sigma},\mathbb{Q}}/Z_G$ -valued cocharacter is defined over \mathbb{Q} , where Z_G denotes the center of G . On the other hand, the composite

$$\mathbb{G}_m \xrightarrow{\mu_{\tilde{\Sigma}}} P_{\tilde{\Sigma},\mathbb{Q}}/Z_G \rightarrow P_{\tilde{\Sigma},\mathbb{Q}}/U_{\tilde{\Sigma},\mathbb{Q}}Z_G$$

is central, hence defined over \mathbb{Q} . Now this composite can be lifted to a $P_{\tilde{\Sigma},\mathbb{Q}}/Z_G$ -valued cocharacter defined over \mathbb{Q} , and any such lift is in $[\mu_{\tilde{\Sigma}}]$, by [DG70, IX, Thm. 3.6]. \square

3.3.7. Somewhat more explicit conditions which guarantee that 3.3.5 holds are given by the following lemma.

Lemma 3.3.8. *Suppose that G^{ad} is \mathbb{Q} -simple. Then $G^{\text{ad}} = \text{Res}_{F/\mathbb{Q}}G_0$ where F is a totally real field, and G_0 is an absolutely simple group over F . If*

- (1) $\tilde{\Sigma}$ is $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ -stable, and
- (2) for some (and hence any, by (1)) embedding $F \rightarrow \mathbb{R}$, the F -rank of G_0 is equal to its \mathbb{R} -rank,

then condition 3.3.5 holds.

Proof. For the fact that G^{ad} has the form $\text{Res}_{F/\mathbb{Q}}G_0$ see [Del79, 2.3.4(a)]. Now suppose that the conditions (1) and (2) are satisfied. Fix an embedding $F \rightarrow \mathbb{R}$, so that $G_{0,\mathbb{R}}$ is a factor of $G_{\mathbb{R}}^{\text{ad}}$, and corresponds to a connected component $\Delta(G)_0$ of the Dynkin diagram of G . Let $\tilde{\Sigma}_0 = \tilde{\Sigma} \cap \Delta(G)_0$, and let $\mu_{\tilde{\Sigma}_0}$ be the corresponding $G_{\mathbb{R},0}^{\text{ad}}$ -valued cocharacter.

By [AMRT10, §3.2], the conjugacy class of the parabolic subgroup of $G_{0,\mathbb{R}}$ corresponding to $\mu_{\tilde{\Sigma}_0}$ contains a parabolic defined over \mathbb{R} , and hence the conjugacy class $[\mu_{\tilde{\Sigma}_0}]$ contains a cocharacter defined over \mathbb{R} , as in the proof of 3.3.6. By (2), after conjugation by an element of $G_0(\mathbb{R})$, this cocharacter, which we again denote by $\mu_{\tilde{\Sigma}_0}$, factors through an \mathbb{R} -split torus which is defined over F , and F -split. Thus we have a map of F -groups $\mu_{\tilde{\Sigma}_0} : \mathbb{G}_m \rightarrow G_0$.

By definition of the restriction of scalars, $\mu_{\tilde{\Sigma}_0}$ induces a map $\mu'_{\tilde{\Sigma}} : \mathbb{G}_m \rightarrow G^{\text{ad}}$ over \mathbb{Q} . To see that $\mu'_{\tilde{\Sigma}} \in [\mu_{\tilde{\Sigma}}]$, note that, by (1), $[\mu_{\tilde{\Sigma}}]$ is $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ -stable. Thus it suffices to check that the projections of $[\mu_{\tilde{\Sigma}}]$ and $[\mu'_{\tilde{\Sigma}}]$ onto G_0 are equal. But these are both equal to $[\mu_{\tilde{\Sigma}_0}]$. \square

3.3.9. Let us explain how to apply Lemma 3.3.8 in examples. First, the condition (1) implies that X has the form X_0^m for some irreducible Hermitian symmetric domain X_0 . When X_0 is not of type $D_n^{\mathbb{H}}$, then (1) is actually equivalent to this condition. In the case of type $D_n^{\mathbb{H}}$, there are two conjugacy classes of cocharacters of $G_{0,\mathbb{R}}$, which are exchanged by an outer isomorphism of G_0 , and give rise to isomorphic (via the outer isomorphism) Hermitian symmetric domains. If the outer isomorphism is induced by complex conjugation, which happens when n is odd,

then X having the form X_0^m still implies (1), but if n is even, then the condition is stronger.

The condition (2) can also be made more explicit in many cases: If G is of type B or C or D_n with n even, then $G_{\mathbb{R}}^{\text{ad}}$ is split, and (2) means that G_0 is a split group, or (in this context) that G is quasi-split. If G is of type A_n then (2) means that G_0 is the adjoint group of a unitary group over F - that is one associated to a Hermitian form on a vector space over a quadratic CM extension of F - and not just an inner form of such a group. In particular, when $n = 1$ this covers the case of Hilbert modular varieties.

The following lemma gives a way of constructing many examples when X is irreducible.

Corollary 3.3.10. *For any irreducible Hermitian domain X , and any quadratic imaginary extension L/\mathbb{Q} , there exists a Shimura datum (G, X) such that G is an absolutely simple group which splits over L , and 3.3.5 holds.*

Proof. Let $G_{\mathbb{R}}$ be the absolutely simple reductive \mathbb{R} -group associated to X , and $G_{0,\mathbb{R}}$ its quasi-split inner form. Consider the Dynkin diagram of $G_{0,\mathbb{R}}$ with its action of $\text{Gal}(\mathbb{C}/\mathbb{R}) \simeq \text{Gal}(L/\mathbb{Q})$. This corresponds to a quasi-split group G_0 over \mathbb{Q} , which splits over L . Now by [PR06, Thm. 1], there exists an inner form G of G_0 , which becomes isomorphic to $G_{\mathbb{R}}$ over \mathbb{R} (as an inner form of $G_{0,\mathbb{R}}$), and whose \mathbb{Q} -rank is equal to the \mathbb{R} -rank of $G_{0,\mathbb{R}}$. (Note that in *loc. cit.* the number of i satisfying the condition (ii) with $v = \infty$ is precisely the split rank of $G_{\mathbb{R}}$.) The corollary now follows from Lemma 3.3.8. \square

3.3.11. We will now apply the theory of toroidal compactifications of Shimura varieties to our situation. We refer the reader to the book of Ash-Mumford-Rapoport-Tai [AMRT10], and Pink's thesis [Pin90]; see also Madapusi-Pera's paper [MP19, §2] for a convenient summary. We suppose from now on that the condition 3.3.5 holds.

Let $P \in [P_{\bar{\Sigma}}]$ be defined over \mathbb{Q} , and let $U \subset P$ be its unipotent radical, and $Z \subset U$ the center of U . Abusing notation slightly, write $U(\mathbb{Z}) = U(\mathbb{Q}) \cap K$, $Z(\mathbb{Z}) = Z(\mathbb{Q}) \cap K$. Let $\xi_K = U(\mathbb{Z}) \backslash U(\mathbb{R})Z(\mathbb{C})$, and $C_K = (Z(\mathbb{C})/Z(\mathbb{Z})) \backslash \xi_K$. Then C_K is an abelian scheme and $\xi_K \rightarrow C_K$ is a torus bundle. Fix a fan Σ , in $X^*(Z(\mathbb{C})/Z(\mathbb{Z}))_{\mathbb{Q}}$ so that the resulting torus embedding is smooth with a normal crossings divisor at the boundary. As in Lemma 3.2.9, we obtain a smooth torus embedding over C_K , $\xi_K(\Sigma) \rightarrow C_K$ whose boundary $D_K(\Sigma)$ is a normal crossings divisor, see [Pin90, §3], [MP19, 2.1.7].

Note that in the constructions of *loc. cit.* one gets a smooth torus embedding over a base which is itself an abelian scheme over a Shimura variety. That is because those constructions start with the preimage under $P \rightarrow P/U$ of a factor of the reductive group P/U . However, for our specific choice of parabolic $P_{\bar{\Sigma}}$ the relevant factor of P/U is a torus, and the Shimura variety is 0-dimensional. In the notation of [Pin90, §4.7], our parabolic P is what Pink denotes by Q , and the preimage mentioned above is denoted by P_1 . These groups are denoted by P and Q_P in [MP19, §2].

3.3.12. By an *analytic neighborhood of $D_K(\Sigma)$* we mean an analytic open neighborhood $V_K \subset \xi_K(\Sigma)(\mathbb{C})$, containing $D_K(\Sigma)$. We write $V_K^* = V_K \backslash D_K(\Sigma)$.

Write $\text{Sh}_K = \text{Sh}_K(G, X)$. Fix a connected component $X^+ \subset X$, and let $\text{Sh}_K^+ \subset \text{Sh}_K$ be the corresponding connected component of Sh_K^+ . Then $\text{Sh}_K^+ = \Gamma \backslash X^+$, where

$\Gamma = K \cap G(\mathbb{Q})_+$, and $G(\mathbb{Q})_+ \subset G(\mathbb{Q})$ is the subgroup mapping to the connected component of the identity in $G^{\text{ad}}(\mathbb{R})$ [Del79, 2.1.2]. In particular, for any choice of base point, this allows us to identify the fundamental group of Sh_K^+ with Γ .

For any smooth \mathbb{C} -scheme Y , by an *ncd compactification* of Y we mean a dense embedding $Y \subset \bar{Y}$ into a proper smooth \mathbb{C} -scheme \bar{Y} , such that $\bar{Y} \setminus Y$ is a normal crossings divisor. We collect the results we need in the following proposition.

Proposition 3.3.13. *There exists an ncd compactification $\text{Sh}_K(\Sigma)^+$ of Sh_K and an analytic neighborhood V_K of $D_K(\Sigma)$ such that*

- (1) *The inclusion $V_K^* \rightarrow \xi_K(\mathbb{C})$ induces an isomorphism of topological fundamental groups.*
- (2) *There is an étale map of complex analytic spaces*

$$\pi_K : V_K \rightarrow \text{Sh}_K(\Sigma)^+$$

such that $\pi_K^{-1}(\text{Sh}_K(\Sigma)^+ \setminus \text{Sh}_K^+) = D_K(\Sigma)$, and $\pi_K : V_K^ \rightarrow \text{Sh}_K^+$ induces the natural map $U(\mathbb{Z}) \hookrightarrow \Gamma$ on fundamental groups.*

Proof. By [Pin90, 4.11], [MP19, 2.1.6] there is an open immersion of complex analytic spaces $X^+ \rightarrow U(\mathbb{R})Z(\mathbb{C})$. Now set $V_K^* = U(\mathbb{Z}) \setminus X^+$, then there is an induced open immersion $V_K^* \rightarrow \xi_K(\mathbb{C})$ (see [Pin90, 6.10], [MP19, 2.1.13]) which induces an isomorphism on fundamental groups, by construction. By [Pin90, 6.13], [MP19, 2.1.22], the latter open immersion extends to an analytic neighborhood V_K of $D_K(\Sigma)$. The existence of $\text{Sh}_K(\Sigma)^+$ and the map π_K with the properties in (2) is the main result of [Pin90, §6], see also [MP19, 2.1.26]. We remark that the ncd compactification $\text{Sh}_K(\Sigma)^+$, depends on more choices than just Σ , but as these will play no role for us, we omit them from the notation. \square

3.3.14. We continue to assume that condition 3.3.5 holds. Suppose that G admits a reductive model $G_{\mathbb{Z}_p}$ over \mathbb{Z}_p . As the scheme of parabolic subgroups is projective [DG70, XXVI, Cor. 3.5], P extends to a parabolic subgroup $P_{\mathbb{Z}_p} \subset G_{\mathbb{Z}_p}$. Denote by $U_{\mathbb{Z}_p} \subset P_{\mathbb{Z}_p}$ the unipotent radical. We will sometimes write G , P and U for $G_{\mathbb{Z}_p}$, $P_{\mathbb{Z}_p}$ and $U_{\mathbb{Z}_p}$ if this causes no confusion.

Lemma 3.3.15. *The group $U_{\mathbb{Z}_p}(\mathbb{Z}_p)$ is a central extension of finitely-generated, free abelian pro- p groups. The map $U(\mathbb{Z}_p) \rightarrow U(\mathbb{F}_p)$ is surjective, and $U(\mathbb{F}_p)$ is a reduction of $U(\mathbb{Z}_p)$ mod p .*

Proof. As in the proof of Lemma 3.3.6, one sees that $P_{\mathbb{Z}_p} \subset G_{\mathbb{Z}_p}$ corresponds to a cocharacter $\mu_{\mathbb{Z}_p}$ in $[\mu_{\bar{\Sigma}}]$, defined over \mathbb{Z}_p . As $U_{\mathbb{Z}_p}$ is unipotent, it is an iterated extension of additive groups. It then follows by Lemma 3.3.3 that $U_{\mathbb{Z}_p}$ is a central extension of additive groups, so that $U(\mathbb{Z}_p)$ is a central extension of finitely generated, free, abelian pro- p groups. As $U_{\mathbb{Z}_p}$ is smooth, $U(\mathbb{Z}_p) \rightarrow U(\mathbb{F}_p)$ is surjective.

To check that $U(\mathbb{F}_p)$ is a reduction of $U(\mathbb{Z}_p)$ mod p , we have to check that $U(\mathbb{F}_p)$ is a Heisenberg group. This condition is vacuous if U is abelian. If U is not abelian, then by Lemma 3.3.3, $\mu_{\mathbb{Z}_p}$ has weight 2 on $Z(U_{\mathbb{Z}_p})$ and weight 1 on $U_{\mathbb{Z}_p}/Z(U_{\mathbb{Z}_p})$. Hence \mathbb{F}_p^\times acting via $\mu_{\mathbb{Z}_p}$ has weight 2 on $Z(U_{\mathbb{Z}_p})(\mathbb{F}_p)$ and weight 1 on $U_{\mathbb{Z}_p}/Z(U_{\mathbb{Z}_p})(\mathbb{F}_p)$. Thus $U(\mathbb{F}_p)$ is a Heisenberg group by Lemma 3.1.2. \square

3.3.16. Now suppose that $K = K_p K^p$, with $K_p = G(\mathbb{Z}_p)$, and $K^p \subset G(\mathbb{A}_f^p)$ compact open. Let $K_p^1 = \ker(G(\mathbb{Z}_p) \rightarrow G(\mathbb{F}_p))$, and $K^1 = K_p^1 K^p$. As above, let $\Gamma = K \cap G(\mathbb{Q})_+$, and set $\Gamma_1 = K^1 \cap G(\mathbb{Q})_+$. A covering, $\Gamma_1 \setminus X^+ \rightarrow \Gamma \setminus X^+$, with Γ, Γ_1 of the above form is called a *principal p -covering*.

Theorem 3.3.17. *Assume that 3.3.5 holds, and that G extends to a reductive group scheme over \mathbb{Z}_p . If $\Sigma \neq c(\Sigma)$ then we assume that $p > \dim X + \frac{1}{2}(\dim Z + 1)$, and that p is a prime of unramified good reduction for ξ_K . Then*

$$\mathrm{ed}(\Gamma_1 \backslash X^+ \rightarrow \Gamma \backslash X^+; p) = \dim X.$$

Proof. Fix a base point $\bar{s} \in V_K$. By Proposition 3.3.13(1), we may identify $\pi_1(V_K^*, \bar{s})$ with $\pi_1(\xi_K(\mathbb{C}), \bar{s}) \simeq U(\mathbb{Z})$. By the strong approximation for unipotent groups, the pro- p completion of $U(\mathbb{Z})$ is $U(\mathbb{Z}_p)$.

The pullback of $\Gamma_1 \backslash X^+ \rightarrow \Gamma \backslash X^+$ by the map π_K in Proposition 3.3.13(2) is a covering, one of whose components $V_{K^1}^* \rightarrow V_K^*$, corresponds to the kernel of the composite

$$U(\mathbb{Z}) \rightarrow \Gamma \rightarrow K \rightarrow G(\mathbb{Z}_p) \rightarrow G(\mathbb{F}_p).$$

That is, it is the kernel of $U(\mathbb{Z}) \rightarrow U(\mathbb{F}_p)$. Using the strong approximation for unipotent groups, as above, it follows that $V_{K^1}^* \rightarrow V_K^*$ is a covering with group $U(\mathbb{F}_p)$, and is the restriction of a covering of schemes $\xi_{K^1} \rightarrow \xi_K$ with group $U(\mathbb{F}_p)$. Let $\xi_{K^1}(\Sigma)$ be the normalization of $\xi_K(\Sigma)$ in ξ_{K^1} , and set $V_{K^1} = \xi_{K^1}(\Sigma)^{\mathrm{an}}|_{V_K}$.

Next suppose $B \subset \mathrm{Sh}_K^+ = \Gamma \backslash X^+$ is a Zariski closed subset. Let \bar{B} be its Zariski closure in $\mathrm{Sh}_K^+(\Sigma)$. Then $\pi_{K^1}^{-1}(\bar{B})$ is Zariski closed in V_{K^1} . It follows that

$$\mathrm{ed}(\Gamma_1 \backslash X^+ \rightarrow \Gamma \backslash X^+; p) \geq \mathrm{ed}(V_{K^1} \rightarrow V_K; p).$$

Here we are using the notion of p -essential dimension for analytic spaces introduced in 3.2.15.

As $U(\mathbb{F}_p)$ is a mod p reduction of $U(\mathbb{Z}_p)$ by Lemma 3.3.15, the theorem follows from Corollary 3.2.17 if $c(\Sigma) \neq \Sigma$ and from Corollary 3.2.18 if $c(\Sigma) = \Sigma$. \square

Corollary 3.3.18. *With the assumptions of Theorem 3.3.17, suppose that (G, X) is of Hodge type. Then p is a prime of unramified good reduction for ξ_K , and the conclusion of the theorem holds without this assumption.*

Proof. This is a consequence of the main result of Madapusi-Pera [MP19]. \square

Corollary 3.3.19. *Let X be an irreducible symmetric domain, and let L/\mathbb{Q} be a quadratic extension. Then there exists a Shimura datum (G, X) with G an absolutely simple group which splits over L , such that for any principal p -covering $\Gamma_1 \backslash X^+ \rightarrow \Gamma \backslash X^+$, we have*

$$\mathrm{ed}(\Gamma_1 \backslash X^+ \rightarrow \Gamma \backslash X^+; p) = \dim X,$$

provided p satisfies the following conditions if X is not a tube domain:

- *If X is of classical type, then $p > \frac{3}{2}\dim X$.*
- *If X is of type E_6 , then p is sufficiently large.*

Proof. Apply Corollary 3.3.10, to obtain a Shimura datum (G, X) such that G is an absolutely simple group which splits over L , and such that 3.3.5 holds. Our definition of principal p -coverings already assumes that G admits a reductive model over \mathbb{Z}_p , so we assume this from now on.

If X is of tube type, that is $c(\Sigma) = \Sigma$ or X is of type E_6 , then the corollary follows from Theorem 3.3.17.

Suppose X is not a tube domain and is of classical type. Then (G, X) is of abelian type [Del79, 2.3.10]. Recall that this means (since G is adjoint), that there is a morphism of Shimura data $(G', X') \rightarrow (G, X)$, induced by a central isogeny $G' \rightarrow G$, with (G', X') of Hodge type. As in [Kis10, 3.4.13], we can assume that G'

again has a reductive model over \mathbb{Z}_p . Now we apply 3.3.18 to a principal p -covering coming from the group G' . We obtain the p -incompressibility of this covering for $p > \dim X + \frac{1}{2}(\dim Z + 1)$. As $\dim X = \frac{1}{2}(\dim U + \dim Z) = \frac{1}{2}\dim(U/Z) + \dim Z$, we have

$$\dim X + \frac{1}{2}(\dim Z + 1) = \dim X + \frac{1}{2}(\dim X - \frac{1}{2}(\dim U/Z) + 1) \leq \frac{3}{2}\dim X.$$

The result for the covering $\Gamma_1 \backslash X^+ \rightarrow \Gamma \backslash X^+$ now follows as in the proof of [FKW19, 4.3.12]. \square

REFERENCES

- [AM04] Alejandro Adem and R. James Milgram, *Cohomology of finite groups*, 2nd ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 309, Springer-Verlag, Berlin, 2004.
- [AF59] Aldo Andreotti and Theodore Frankel, *The Lefschetz theorem on hyperplane sections*, *Annals of Math.* **69** (1959), 713–717.
- [ALB19] Johannes Anschütz and Arthur-César Le Bras, *Prismatic Dieudonné Theory*, arXiv:1907.10525 (2019), 104 pages.
- [Arn70a] V. I. Arnol'd, *Certain topological invariants of algebraic functions*, *Trudy Moskov. Mat. Obsč.* **21** (1970), 27–46 (Russian).
- [Arn70b] ———, *Topological invariants of algebraic functions. II*, *Funkcional. Anal. i Priložen.* **4** (1970), no. 2, 1–9 (Russian).
- [AMRT10] Avner Ash, David Mumford, Michael Rapoport, and Yung-Sheng Tai, *Smooth compactifications of locally symmetric varieties*, 2nd ed., Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2010. With the collaboration of Peter Scholze.
- [BC92] D. J. Benson and Jon F. Carlson, *The cohomology of extraspecial groups*, *Bull. London Math. Soc.* **24** (1992), no. 3, 209–235.
- [BF20] Patrick Brosnan and Najmuddin Fakhruddin, *Fixed points, local monodromy, and incompressibility of congruence covers*, arXiv:2007.01752 (2020).
- [BS19] Bhargav Bhatt and Peter Scholze, *Prisms and Prismatic Cohomology*, arXiv:1905.08229 (2019), 110 pages.
- [Bro94] Kenneth S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York, 1994. Corrected reprint of the 1982 original.
- [BR97] J. Buhler and Z. Reichstein, *On the essential dimension of a finite group*, *Compositio Math.* **106** (1997), no. 2, 159–179.
- [BR99] Joe Buhler and Zinovy Reichstein, *On Tschirnhaus transformations*, *Topics in number theory* (University Park, PA, 1997), *Math. Appl.*, vol. 467, Kluwer Acad. Publ., Dordrecht, 1999, pp. 127–142.
- [Bur12] Yuri Burda, *Coverings over tori and topological approach to Klein's resolvent problem*, *Transform. Groups* **17** (2012), no. 4, 921–951.
- [BW00] A. Borel and N. Wallach, *Continuous cohomology, discrete subgroups, and representations of reductive groups*, 2nd ed., *Mathematical Surveys and Monographs*, vol. 67, American Mathematical Society, Providence, RI, 2000.
- [CT02] J.L. Colliot-Thélène, *Exposant et indice d'algèbres simples centrales non ramifiées. (With an appendix by Ofer Gabber.)*, *Enseigne. Math.* **48** (2002), no. 1–2, 127–146.
- [Del79] Pierre Deligne, *Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques*, *Automorphic forms, representations and L-functions* (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, 1979, pp. 247–289.
- [DI87] Pierre Deligne and Luc Illusie, *Relèvements modulo p^2 et décomposition du complexe de de Rham*, *Invent. Math.* **89** (1987), no. 2, 247–270.
- [DG70] Michel Demazure and Alexander Grothendieck, *Schémas en groupes*, *Lecture Notes in Mathematics*, Vol. 151, 152, 153, Springer-Verlag, Berlin-New York, 1970. Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3).
- [DLLZ19] Hansheng Diao, Kai-Wen Lan, Ruochuan Liu, and Xinwen Zhu, *Logarithmic adic spaces: Some foundational results*, arXiv:1912.09836 (2019), 81 pages.

- [Gro61] A. Grothendieck, *Éléments de géométrie algébrique*, Inst. Hautes Études Sci. Publ. Math. **4,8,11,17,20,24,32** (1961,1962,1964).
- [FS21] Najmuddin Fakhruddin and Rijui Saini, *Finite groups scheme actions and incompressibility of Galois covers: beyond the ordinary case*, arXiv:2102.05993 (2021), 22 pages.
- [FKW19] Benson Farb, Mark Kisin, and Jesse Wolfson, *The essential dimension of congruence covers*, *Compositio*, to appear (2019), 27 pages.
- [Ful93] William Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993. The William H. Roever Lectures in Geometry.
- [HS53] G. Hochschild and J.-P. Serre, *Cohomology of group extensions*, Trans. Amer. Math. Soc. **74** (1953), 110–134.
- [Hub96] Roland Huber, *Étale cohomology of rigid analytic varieties and adic spaces*, Aspects of Mathematics, E30, Friedr. Vieweg & Sohn, Braunschweig, 1996.
- [Ill02] Luc Illusie, *An overview of the work of K. Fujiwara, K. Kato, and C. Nakayama on logarithmic étale cohomology*, Astérisque **279** (2002), 271–322. Cohomologies p -adiques et applications arithmétiques, II.
- [Kat89] Kazuya Kato, *Logarithmic structures of Fontaine-Illusie*, Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988), Johns Hopkins Univ. Press, Baltimore, MD, 1989, pp. 191–224.
- [Kat73] Nicholas M. Katz, *p -adic properties of modular schemes and modular forms*, Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), Springer, Berlin, 1973, pp. 69–190. Lecture Notes in Mathematics, Vol. 350.
- [KKMSD73] G. Kempf, Finn Faye Knudsen, D. Mumford, and B. Saint-Donat, *Toroidal embeddings. I*, Lecture Notes in Mathematics, Vol. 339, Springer-Verlag, Berlin-New York, 1973.
- [Kis10] Mark Kisin, *Integral models for Shimura varieties of abelian type*, J. Amer. Math. Soc. **23** (2010), no. 4, 967–1012.
- [Kl1884] Felix Klein, *Vorlesungen über das Ikosaeder und die Auflösung der Gleichungen vom fünften Grade (Lectures on the Icosahedron and the Solution of the Equation of the Fifth Degree)*, Leipzig, Tübner, 1884.
- [Kos20] Teruhisa Koshikawa, *Logarithmic Prismatic Cohomology*, arXiv:2007.14037 (2020), 62 pages.
- [KY] Teruhisa Koshikawa and Zijian Yao, *Logarithmic Prismatic Cohomology II*, in preparation.
- [Kr1861] Leopold Kronecker, *Ueber die Gleichungen fünften Grades*, Journal für die reine und angewandte Mathematik **59** (1861), 306–310.
- [LMB00] Gérard Laumon and Laurent Moret-Bailly, *Champs algébriques*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 39, Springer-Verlag, Berlin, 2000 (French).
- [Loj64] S. Lojasiewicz, *Triangulation of semi-analytic sets*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) **18** (1964), 449–474.
- [MP19] Keerthi Madapusi Pera, *Toroidal compactifications of integral models of Shimura varieties of Hodge type*, Ann. Sci. Éc. Norm. Supér. (4) **52** (2019), no. 2, 393–514 (English, with English and French summaries).
- [Pey08] Emmanuel Peyre, *Unramified cohomology of degree 3 and Noether’s problem*, Invent. Math. **171** (2008), no. 1, 191–225.
- [Pin90] Richard Pink, *Arithmetical compactification of mixed Shimura varieties*, Bonner Mathematische Schriften [Bonn Mathematical Publications], vol. 209, Universität Bonn, Mathematisches Institut, Bonn, 1990. Dissertation, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, 1989.
- [PR06] Gopal Prasad and Andrei S. Rapinchuk, *On the existence of isotropic forms of semi-simple algebraic groups over number fields with prescribed local behavior*, Adv. Math. **207** (2006), no. 2, 646–660.

- [RY00] Zinovy Reichstein and Boris Youssin, *Essential dimensions of algebraic groups and a resolution theorem for G -varieties*, *Canad. J. Math.* **52** (2000), no. 5, 1018–1056.
With an appendix by János Kollár and Endre Szabó.
- [Tsc34] N.G. Tschebotaröw, *Die Probleme der Modernen Galoisschen Theorie*, *Comment. Math. Helv.* **6** (1934), 235–283.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO
Email address: `farb@math.uchicago.edu`

DEPARTMENT OF MATHEMATICS, HARVARD
Email address: `kisin@math.harvard.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA-IRVINE
Email address: `wolfson@uci.edu`