

ON SOME GEOMETRIC REPRESENTATIONS OF $\mathrm{GL}_n(\mathcal{O})$

URI BADER AND URI ONN

ABSTRACT. We study a family of complex representations of the group $\mathrm{GL}_n(\mathcal{O})$, where \mathcal{O} is the ring of integers of a non-archimedean local field \mathbf{F} . These representations occur in the restriction of the Grassmann representation of $\mathrm{GL}_n(\mathbf{F})$ to its maximal compact subgroup $\mathrm{GL}_n(\mathcal{O})$. We compute the Fourier decomposition of the Hecke algebra associated with the representation in terms of combinatorial data coming from lattices of submodules of finite \mathcal{O} -modules.

1. INTRODUCTION

Let \mathbf{F} be a non-archimedean local field and \mathcal{O} its ring of integers. Let \mathbb{F}_q be the residue field and \mathfrak{o} the maximal ideal of \mathcal{O} . The group $\mathrm{GL}_n(\mathbf{F})$ acts on $\mathrm{Gr}(m, n, \mathbf{F})$, the Grassmannian of m -dimensional subspaces of an n -dimensional space, and gives rise to a complex representation of $\mathrm{GL}_n(\mathbf{F})$ on $L^2(\mathrm{Gr}(m, n, \mathbf{F}))$. This work is about the restriction of this representation to the maximal compact subgroup $\mathrm{GL}_n(\mathcal{O})$. We shall refer to it as the *Grassmann representation*.

As we shall see, the analysis of this representation is deeply connected to the structure of the lattices of submodules of the finite modules $(\mathcal{O}/\mathfrak{o}^k)^n$. Invariants of representations such as multiplicities and idempotents can be expressed in terms of purely geometric or combinatorial data.

1.1. Main Results. Fix $m \leq n/2$. Let Λ_m be the set of isomorphism types of finite \mathcal{O} -modules which are quotients of \mathcal{O}^m , i.e. have rank at most m . We shall identify elements in Λ_m with partitions of at most m parts in the standard way, namely, $(\lambda_i) \leftrightarrow [\oplus \mathcal{O}/\mathfrak{o}^{\lambda_i}]$. For any $\lambda \in \Lambda_m$ let X_λ be the set of submodules of $(\mathcal{O}/\mathfrak{o}^k)^n$ of type λ (for k large enough which will not play any role). The action of $\mathrm{GL}_n(\mathcal{O})$ on each

2000 *Mathematics Subject Classification.* Primary 22D10; Secondary 20C33.

Both authors were supported by the Israel Science Foundation, ISF grant no. 100146. The second author was partially supported by NWO grant no. 613.006.573.

X_λ (via reduction modulo \wp^k) gives rise to a complex representation on the space of \mathbb{C} -valued function on X_λ which we denote by \mathcal{F}_λ .

Theorem 1. *With the above notations there exist a family $\{\mathcal{U}_\lambda\}_{\lambda \in \Lambda_m}$ of irreducible representations of $\mathrm{GL}_n(\mathcal{O})$ such that*

- (1) $L^2(\mathrm{Gr}(m, n, \mathbf{F})) = \bigoplus_{\lambda \in \Lambda_m} \mathcal{U}_\lambda$.
- (2) $\langle \mathcal{U}_\lambda, \mathcal{F}_\mu \rangle = |\{\lambda \hookrightarrow \mu\}|$. *I.e., the multiplicity of \mathcal{U}_λ in \mathcal{F}_μ is the number of nonequivalent embeddings of a module of type λ in a module of type μ .*

Let \mathcal{H}_m be the Hecke algebra associated with the representation:

$$\mathcal{H}_m = \mathcal{S}(\mathrm{Gr}(m, n, \mathbf{F}) \times \mathrm{Gr}(m, n, \mathbf{F}))^{\mathrm{GL}_n(\mathcal{O})}$$

where \mathcal{S} stands for Bruhat-Schwartz functions. One has a dense embedding $\mathcal{H}_m \hookrightarrow \mathrm{End}_{\mathrm{GL}_n(\mathcal{O})}(L^2(\mathrm{Gr}(m, n, \mathbf{F})))$ given by interpreting elements of \mathcal{H}_m as integration kernels. Thus, \mathcal{H}_m plays a central role in the analysis of the Grassmann representation. It has two natural bases indexed by elements from Λ_m : characteristic functions of classes (*geometric basis*) and idempotents (*algebraic basis*).

Theorem 2. *The idempotents in \mathcal{H}_m are computed explicitly (see §4.2) in terms of the geometric basis using combinatorial data coming from the lattice of submodules of finite quotient of \mathcal{O}^m .*

The whole study included here hinges on the profinite nature of the ring of integers \mathcal{O} , and hence of all groups, spaces and algebras defined over it:

| | |
|-----------------|--|
| Groups | $\mathrm{GL}_n(\mathcal{O}) \simeq \varprojlim \mathrm{GL}_n(\mathcal{O}/\wp^k)$ |
| Spaces | $\mathrm{Gr}(m, n, \mathbb{F}) \simeq \varprojlim \mathrm{Gr}(m, n, \mathcal{O}/\wp^k)$ |
| Representations | $L^2(\mathrm{Gr}(m, n, \mathbb{F})) \simeq \varinjlim L^2(\mathrm{Gr}(m, n, \mathcal{O}/\wp^k))$ |
| Algebras | $\mathcal{H}_m \simeq \varinjlim \mathrm{End}_{\mathrm{GL}_n(\mathcal{O})}(\mathcal{F}_{k^m})$ |

As a consequence, one can work with finite objects, and then transport the results to the limit. This also dictates the organization of this paper. Section §2 is devoted to the transition to and from the finite objects. Section §3 is a purely finite analysis and concerns the interplay between the combinatorics of the lattice of submodules of $(\mathcal{O}/\wp^k)^m$ and the representation \mathcal{F}_{k^m} . In particular, the Fourier decomposition of the finite Hecke algebra $\mathrm{End}_{\mathrm{GL}_n(\mathcal{O})}(\mathcal{F}_{k^m})$ is explicitly computed. In section §4 the decomposition is lifted to $\mathrm{GL}_n(\mathcal{O})$ and explicit formulas are given. The last section, §5, is devoted to related works. The appendix at the end of this paper is a collection of some lemmas required for the

main theorem of section §3 regarding discrete valuation rings and their modules, which we believe are known but could not find a reference.

1.2. Acknowledgements. We are most grateful to Amos Nevo for hosting and encouraging this research. We warmly thank Shai Haran for stimulating discussions, his interest and encouragement.

2. TO FINITE QUOTIENTS AND BACK

2.1. Reduction to finite quotients. Let I_k denote the kernel of the reduction of $\mathrm{GL}_n(\mathcal{O})$ modulo \wp^k , and let G_k stand for the quotient $\mathrm{GL}_n(\mathcal{O}/\wp^k)$. $\mathrm{GL}_n(\mathcal{O})$ being the inverse limit of the finite groups G_k enjoys the property that each of its continuous irreducible complex representations has a *conductor* or *level*. That is the first natural number k such that I_k acts trivially. It follows that there exist a natural filtration on the representation,

$$(0) \subset L^2(\mathrm{Gr}(m, n, \mathbf{F}))^{I_1} \subset L^2(\mathrm{Gr}(m, n, \mathbf{F}))^{I_2} \subset \cdots \subset L^2(\mathrm{Gr}(m, n, \mathbf{F}))$$

The k -th term in the filtration consists of all the irreducible components of the representation which have level at most k , and thus is in fact a representation of G_k . Denote this representation by $\bar{\rho}_{k^m}$. Since each irreducible will be captured by some $\bar{\rho}_{k^m}$, it suffices to understand the latter.

$\bar{\rho}_{k^m}$ has the following alternative description. Fix a level k , and let \mathcal{L}_n^k be the lattice of submodules of $(\mathcal{O}/\wp^k)^n$. G_k acts on \mathcal{L}_n^k with orbits:

$$X_\lambda = \{x \in \mathcal{L}_n^k \mid x \simeq \mathcal{O}/\wp^{\lambda_1} \oplus \cdots \oplus \mathcal{O}/\wp^{\lambda_n}\} \quad \lambda \in \Lambda_n^k$$

Where $\Lambda_n^k = \{(\lambda_1, \dots, \lambda_n) \mid k \geq \lambda_1 \geq \cdots \geq \lambda_n \geq 0\}$ stands for isomorphism types of elements in \mathcal{L}_n^k . For each λ define a representation of G_k arising from this action on $\mathcal{F}_\lambda = \mathcal{F}(X_\lambda)$, the vector space of \mathbb{C} -valued functions on X_λ , with inner product induced from the counting measure. Denote this representation be ρ_λ . Of particular interest is the representation which corresponds to $\lambda = k^m$ due to the following observation:

Lemma 2.1. $\rho_{k^m} \simeq \bar{\rho}_{k^m}$.

The set of types Λ_n^k is equipped with a natural partial order: $\mu \leq \nu$ whenever a module of type μ can be embedded in a module of type ν . In terms of Young diagrams it amounts to containment. Let $\tau : \mathcal{L}_n^k \rightarrow \Lambda_n^k$ be the *type* map which assigns to each module its isomorphism type. It will also be useful to use another set of coordinates for elements in Λ_n^k by transposed diagrams $\lambda' = (\lambda'_j)$, defined by $\lambda'_j = |\{i : \lambda_i \geq j\}|$.

2.2. The finite Hecke algebras. Let $\mathcal{H}_{k^m} = \text{Hom}_{G_k}(\mathcal{F}_{k^m}, \mathcal{F}_{k^m})$ be the Hecke algebra of G_k -invariant endomorphisms of \mathcal{F}_{k^m} . It is isomorphic to the convolution algebra $\mathcal{F}(X_{k^m} \times_{G_k} X_{k^m})$ by interpreting such functions as a G_k -invariant summation kernel and hence an intertwining endomorphism. We have the following bijection:

$$(\star) \quad \begin{aligned} X_{k^m} \times_{G_k} X_{k^m} &\xrightarrow{\sim} [0, k^m] = \Lambda_m^k \\ [(x, y)] &\longmapsto \tau(x \cap y) \end{aligned}$$

The fact that G_k preserves the module structure implies that this map is well defined. It is onto due to the assumption $m \leq n/2$ which give enough room to realize any type λ as intersection of two (free) modules of type k^m . It is one to one because any abstract isomorphism between $x \cap y$ and $x' \cap y'$ can be lifted to an element $g \in G_k$ such that $(x', y') = g(x, y)$. In particular $[(x, y)] = [(y, x)]$ and an application of Gelfand's trick gives:

Proposition 2.2.

- (1) $\mathcal{H}_{k^m} \simeq \mathbb{C}^{[0, k^m]}$.
- (2) \mathcal{F}_{k^m} is multiplicity free and decomposes to $|[0, k^m]| = \binom{k+m}{m}$ irreducibles.

The explicit construction of the isomorphism in part (1) of proposition 2.2 is the main theme of section §3. Before pursuing this task we describe how the finite results are lifted to the analysis of the Grassmann representation.

2.3. Lifting the finite spaces and algebras.

2.3.1. Lifting spaces. Let X_m stand for $\text{Gr}(m, n, \mathbf{F})$ and $\pi_k : X_{k^m} \rightarrow X_{(k-1)^m}$ be the natural quotient maps. As $\text{GL}_n(\mathcal{O})$ -spaces we have:

$$X_m \simeq \varprojlim X_{k^m}$$

Observe that $I_k \backslash X_m \simeq X_{k^m}$, thus supplying a proof for lemma 2.1. Using the identification (\star) above, the maps π_k descent to maps

$$\pi_k : \Lambda_m^k \rightarrow \Lambda_m^{k-1}, \quad \lambda \mapsto \lambda / \mathfrak{O}^{k-1} \lambda$$

which are easily described in transposed coordinates by

$$(\lambda'_1, \dots, \lambda'_{k-1}, \lambda'_k) \mapsto (\lambda'_1, \dots, \lambda'_{k-1})$$

Consider the set $\prod_{k \geq 0} \Lambda_m^k$, and endow it with a graph structure by connecting each $\lambda \in \Lambda_m^k$ with its image $\pi_k(\lambda) \in \Lambda_m^{k-1}$. This graph is a rooted tree, the root being the empty partition in Λ_m^0 . $\varprojlim \Lambda_m^k$ can be identified with the space of ends of this tree. The obvious sections $\Lambda_m^{k-1} \rightarrow \Lambda_m^k$ given by $(\lambda'_1, \dots, \lambda'_{k-1}) \mapsto (\lambda'_1, \dots, \lambda'_{k-1}, 0)$ give at the

limit an imbedding of Λ_m in $\varprojlim \Lambda_m^k$. Thus, Λ_m can be identified with an open and dense subset of $\varprojlim \Lambda_m^k$. Let $\overline{\mathbb{N}}$ stand for the one point compactification of \mathbb{N} . Λ_m is naturally imbedded in \mathbb{N}^m . We denote by $\overline{\Lambda}_m$ its closure in $\overline{\mathbb{N}}^m$. It is easily seen that $\overline{\Lambda}_m$ can be identified with $\varprojlim \Lambda_m^k$. We summarize this discussion by

Proposition 2.3.

$$X_m \times_{\mathrm{GL}_n(\mathcal{O})} X_m \simeq \overline{\Lambda}_m \simeq \varprojlim \Lambda_m^k$$

Proof. The only nontrivial issue left to address is the fact that the first identification is also topological. The topology on the l.h.s is the quotient topology. The quotient map from $X_m \times X_m$ to $X_{k^m} \times X_{k^m}$ is continuous and $\mathrm{GL}_n(\mathcal{O})$ -equivariant. The limit map becomes continuous and well-defined on the quotient. \square

Remark 2.4.

- (1) *As topological spaces, $\overline{\Lambda}_m \setminus \Lambda_m \simeq \overline{\Lambda}_{m-1}$, thus $\coprod_{i=0}^m \Lambda_i$ is a stratification of $\overline{\Lambda}_m$.*
- (2) *X_m carries a $\mathrm{GL}_n(\mathcal{O})$ -invariant measure. Consequently, also does $\overline{\Lambda}_m$. Apparently, Λ_m is of full measure inside $\overline{\Lambda}_m$. The computation of this measure was carried in the second author's thesis [Onn03].*

2.3.2. *Representations and algebras.* The maps π_k give rise to inclusions of the (finite dimensional) spaces $i_k : L^2(X_{(k-1)^m}) \rightarrow L^2(X_{k^m})$, where the notation $\mathcal{F}(X)$ is replaced by $L^2(X)$ to emphasize that the inner product structure is induced from the Haar measure, rather than the counting measure. The adjoint transformation, i_k^* , is the orthogonal projection on the I_{k-1} invariants. In the limit we get the space of Bruhat-Schwartz functions:

$$\mathcal{S}(X_m) \simeq \varinjlim L^2(X_m)^{I_k} \simeq \varinjlim L^2(X_{k^m})$$

$L^2(X_m)$ is the completion $\mathcal{S}(X_m)$. The inclusions i_k also give embedding of the Hecke algebras $\mathcal{H}_{k^m} = \mathrm{End}_{G_k}(L^2(X_{k^m}))$ given by

$$\mathcal{H}_{(k-1)^m} \rightarrow \mathcal{H}_{k^m}, \quad h \mapsto i_k \circ h \circ i_k^*$$

It follows (by (\star) above) that as vector spaces

$$\mathcal{H}_{k^m} \simeq \mathcal{F}(\Lambda_m^k)$$

For $\lambda \in \Lambda_m^k$, let $\delta_\lambda^k \in \mathcal{H}_{k^m}$ denote the delta function supported on λ .

Claim 2.5. *If $\lambda'_{k-1} = 0$ then $i_k \circ \delta_\lambda^{k-1} \circ i_k^* = \delta_\lambda^k$.*

Proof. The condition $\lambda'_{k-1} = 0$ implies that $\pi_k^{-1}(\lambda)$ is the singleton $\{\lambda\} \subset \Lambda_m$, and the claim follows. \square

It follows that the image of δ_λ^k inside \mathcal{H}_m stabilizes for k large enough, thus determining an element $\varinjlim \delta_\lambda^k$. We obtain that

$$\delta_\lambda = \varinjlim \delta_\lambda^k$$

where δ_λ is the delta function supported at λ , viewing λ as an element of $\bar{\Lambda}_m$ via $\Lambda_m \subset \bar{\Lambda}_m$, and using the identification

$$\mathcal{S}(\bar{\Lambda}_m) \simeq \varinjlim \mathcal{H}_k \quad \text{by proposition 2.3}$$

Here $\mathcal{S}(\bar{\Lambda}_m)$ is the space of locally constant functions on $\bar{\Lambda}_m$. The limit algebra structure obviously coincides with the operator algebra structure of $\mathcal{S}(\bar{\Lambda}_m)$, arising when viewing its elements as convolution operators on $L^2(X_m)$. Denote by $\mathcal{F}(\Lambda_m)$ the space of finitely supported functions on Λ_m . As Λ_m is discrete in $\bar{\Lambda}_m$, $\mathcal{F}(\Lambda_m)$ is imbedded in $\mathcal{S}(\bar{\Lambda}_m)$. As Λ_m is dense in $\bar{\Lambda}_m$, $\mathcal{F}(\Lambda_m)$ is dense in $\mathcal{S}(\bar{\Lambda}_m)$ too. Consequently, the algebraic structure of $\mathcal{S}(\bar{\Lambda}_m)$ is determined by $\mathcal{F}(\Lambda_m) = \text{Span}\{\delta_\lambda\}$. Analyzing the latter is the task we are undertaking here.

3. FOURIER DECOMPOSITION OF THE FINITE HECKE ALGEBRA

We are now about to compute the Fourier decomposition of the finite Hecke algebras \mathcal{H}_{k^m} . The parameter k will be fixed throughout this section. It will be convenient to denote $\phi = k^m$ and $\Phi = k^n$.

3.1. Geometric intertwiners. The following operators are fundamental in our analysis.

- **$\mathbf{T}_{\mu \succ \lambda}$ and $\mathbf{T}_{\lambda \prec \mu}$:** For each pair of types $\lambda \leq \mu$ define

$$\begin{aligned} \mathbf{T}_{\mu \succ \lambda} : \mathcal{F}_\lambda &\longrightarrow \mathcal{F}_\mu \\ h &\longmapsto \mathbf{T}_{\mu \succ \lambda} h(y) = \sum_{x \subset y} h(x) \quad y \in X_\mu \end{aligned}$$

$$\begin{aligned} \mathbf{T}_{\lambda \prec \mu} : \mathcal{F}_\mu &\longrightarrow \mathcal{F}_\lambda \\ h &\longmapsto \mathbf{T}_{\lambda \prec \mu} h(x) = \sum_{y \supset x} h(y) \quad x \in X_\lambda \end{aligned}$$

$\mathbf{T}_{\mu \succ \lambda}$ and $\mathbf{T}_{\lambda \prec \mu}$ commute with the group action and form an adjoint pair.

- **$\mathbf{T}_{\lambda \leftrightarrow \nu \leftarrow \mu}$:** For types $\lambda, \mu \leq \nu$ define the following map from \mathcal{F}_μ to \mathcal{F}_λ :

$$\mathbf{T}_{\lambda \leftrightarrow \nu \leftarrow \mu} h(x) = \sum_{\left\{ y \mid \begin{array}{l} \tau(y) = \mu \\ \tau(y+x) = \nu \end{array} \right\}} h(y) \quad x \in X_\lambda$$

- \mathbf{g}_λ : For each type $\lambda \leq \phi$, define an endomorphism \mathbf{g}_λ of \mathcal{F}_ϕ by:

$$\mathbf{g}_\lambda h(x) = \sum_{\{y | \tau(y \cap x) = \lambda\}} h(y)$$

Observe that $\mathbf{g}_\lambda = \binom{\Phi}{\phi} \delta_\lambda^k$, due to the transition from Haar measure to the counting measure. \mathbf{g}_λ is self adjoint and commutes with the group action.

In order to minimize confusion, we follow the rule that whenever an operator is labeled with a diagram, it acts from the space indexed by the *right* type of the diagram to the space indexed by the *left* type.

3.2. Bases. We shall be focused on three bases of the Hecke algebra:

$$\begin{aligned} \mathbf{Geometric:} & \quad \{\mathbf{g}_\lambda\}_{\lambda \leq \phi} \\ \mathbf{Cellular:} & \quad \{\mathbf{c}_\lambda\}_{\lambda \leq \phi} \\ \mathbf{Idempotents:} & \quad \{\mathbf{e}_\lambda\}_{\lambda \leq \phi} \end{aligned}$$

The geometric basis is just the indicating functions of the orbits $X_\phi \times_{G_k} X_\phi$ viewed as operators, and explicitly defined above. The cellular basis elements \mathbf{c}_λ are defined by $\mathbf{c}_\lambda = \mathbf{T}_{\phi \succ \lambda} \mathbf{T}_{\lambda \prec \phi}$ for $\lambda \leq \phi$. The following equality is easily verified from the definitions and proves that it is indeed a basis.

$$(\mathbf{c-g}) \quad \mathbf{c}_\lambda = \sum_{\phi \geq \mu \geq \lambda} \binom{\mu}{\lambda} \mathbf{g}_\mu$$

Here $\binom{\mu}{\lambda}$ is the number of submodules of type λ contained in a module of type μ . For each $\lambda \leq \phi$ set:

$$\mathcal{H}_\phi^\lambda = \text{Span}_{\mathbb{C}}\{\mathbf{c}_{\lambda'} | \lambda' \leq \lambda\} \quad \mathcal{H}_\phi^{\lambda^-} = \text{Span}_{\mathbb{C}}\{\mathbf{c}_{\lambda'} | \lambda' < \lambda\}$$

The following proposition is proved in [Hil94] and in a more general setup in [BO04]. It reveals the importance of the cellular base.

Proposition 3.1. \mathcal{H}_ϕ^λ and $\mathcal{H}_\phi^{\lambda^-}$ are ideals $\forall \lambda \in [0, \phi]$.

Corollary 3.

- (1) $\left\{ \mathcal{K}_\lambda = \mathcal{H}_\phi^\lambda / \mathcal{H}_\phi^{\lambda^-} \right\}_{\lambda \leq \phi}$ is a complete set of irreducible representations of \mathcal{H}_ϕ .
- (2) If \mathbf{e}_λ is the idempotent in \mathcal{H}_ϕ corresponding to \mathcal{K}_λ for all $\lambda \leq \phi$ then there is a triangular matrix such that:

$$(\mathbf{c-e}) \quad \mathbf{c}_\lambda = \sum_{\kappa \leq \lambda} A_{\lambda \kappa} \mathbf{e}_\kappa$$

The cellular basis appears as a bridge between the geometric and algebraic bases. It is upper triangular with respect to the former and lower triangular with respect to the latter. In the next section we shall use it to compute the idempotents explicitly.

Remark: Corollary 3 can be used to label the irreducible representations of \mathcal{H}_ϕ . \mathcal{K}_λ is the *unique* representation which is **annihilated** by all $\{\mathcal{H}_\phi^\mu\}_{\mu < \lambda}$ and **not annihilated** by \mathcal{H}_ϕ^λ . In view of the well known dictionary between representations of G which occur in \mathcal{F}_ϕ and representations of the Hecke algebra \mathcal{H}_ϕ , we can now label the irreducibles in \mathcal{F}_ϕ by: $\mathcal{U}_\lambda \leftrightarrow \mathcal{K}_\lambda$.

Moreover, by the definition of \mathbf{c}_λ as the composition $\mathbf{T}_{\phi > \lambda} \mathbf{T}_{\lambda < \phi}$, the annihilation criterion above translates to the fact that \mathcal{U}_λ occurs in \mathcal{F}_λ and does not occur in \mathcal{F}_μ for $\mu < \lambda$.

3.3. Inverting the matrix (c-g). The discussions in [BO04] and appendix A.1 here enable the following calculation:

Proposition 3.2.

$$\mathbf{g}_\lambda = \sum_{\{\nu | \phi \geq \nu \geq \lambda \geq \phi \nu\}} (-1)^{|\nu| - |\lambda|} q^{n(\nu) - n(\lambda)} \prod_{i \geq 1} \begin{bmatrix} \nu'_i - \nu'_{i+1} \\ \nu'_i - \lambda'_i \end{bmatrix} \cdot \mathbf{c}_\nu$$

Recall that for every partition ξ , $n(\xi) = \sum (i-1)\xi_i$, and that $|\xi| = \sum \xi_i$.

Proof. By relation (c-g) in §2.2,

$$\mathbf{c}_\lambda = \sum_{\phi \geq \nu \geq \lambda} \binom{\nu}{\lambda} \mathbf{g}_\nu$$

By [BO04, Corollary 2.10 and Example 2.11], the inverse of the matrix $\binom{\nu}{\lambda}$ is given by

$$\hat{\mu}(\lambda, \nu) = \sum_{\lambda \xleftrightarrow{i} \nu} \binom{\nu = \nu}{\lambda \xleftrightarrow{i} \nu} \chi(\text{coker}(i))$$

The function χ is defined and computed in A.1 below, and for every embedding type $\lambda \xleftrightarrow{i} \nu$ it is given by

$$\chi(\text{coker}(i)) = \begin{cases} 0 & \wp \cdot \text{coker}(i) \neq 0 \\ (-1)^{|\nu| - |\lambda|} q^{\binom{|\nu| - |\lambda|}{2}} & \wp \cdot \text{coker}(i) = 0 \end{cases}$$

whereas the number

$$\sum_{\{\lambda \xleftrightarrow{i} \nu | \wp \cdot \text{coker}(i) = 0\}} \binom{\nu = \nu}{\lambda \xleftrightarrow{i} \nu}$$

is exactly the Hall coefficient $G_{\lambda, 1^{(|\nu| - |\lambda|)}}^\nu$. The latter is explicitly computed in [Mac95, Chapter 4]. \square

3.4. Transition matrix: cellular to idempotents. Let F be a fixed \mathcal{O} -module of type $\Phi = k^n$. We say that two submodules are *transversal* if their intersection is zero. Let κ , λ and μ be types of modules and let x_μ and x_κ be two transversal submodules of F of types μ and κ respectively. Define:

- $[\mu \prec \dot{\lambda} \cap \kappa]_\Phi =$ The number of submodules of type λ in F which contain x_μ and are transversal to x_κ .
- $[\mu \prec \dot{\lambda}]_\Phi =$ the number of submodules of F of type λ which contain a given submodule of type μ (in the above notation this is $[\mu \prec \dot{\lambda} \cap 0]_\Phi$).

We wish to compute the idempotents of \mathcal{H}_ϕ explicitly. Since the cellular structure must agree with the idempotent decomposition, we already know that there exist a lower triangular matrix $A_{\lambda\kappa}$ such that the relation **c-e** above holds. We have already seen that the transition matrix from the geometric basis to the cellular basis depends only on geometric invariants of the lattice of submodules (relation **c-g** above) in a very simple way. We were happy to discover that this also holds for the cellular-idempotents matrix. The main result in this section is:

Theorem 4. $A_{\lambda\kappa} = [\kappa \prec \dot{\lambda}]_\phi [\lambda \prec \dot{\phi} \cap \kappa]$.

Our strategy is to analyze the multiplication in the algebra with respect to the cellular basis. Let $B_{\lambda\mu}^\nu$ be multiplication coefficients with respect to the cellular basis:

$$(1) \quad \mathbf{c}_\lambda \cdot \mathbf{c}_\kappa = \sum_{\nu \leq \lambda \wedge \kappa} B_{\lambda\kappa}^\nu \mathbf{c}_\nu$$

Observe that $B_{\lambda\kappa}^\kappa = A_{\lambda\kappa}$ for $\kappa \leq \lambda$.

Lemma 3.3.

- (1) $\mathbf{T}_{\phi \succ \kappa} \mathbf{T}_{\kappa \succ \nu} = [\nu \prec \dot{\kappa}]_\phi \mathbf{T}_{\phi \succ \nu}$.
- (2) $\mathbf{T}_{\nu \prec \kappa} \mathbf{T}_{\kappa \prec \phi} = [\nu \prec \dot{\kappa}]_\phi \mathbf{T}_{\nu \prec \phi}$.

Proof. Follows immediately from lemma 3.4 in [BO04]. \square

Substituting $\mathbf{c}_\eta = \mathbf{T}_{\phi \succ \eta} \mathbf{T}_{\eta \prec \phi}$ in equation (1) and using lemma 3.3 gives (assume $\kappa \leq \lambda$):

$$(2) \quad \begin{aligned} & \mathbf{T}_{\phi \succ \lambda} \left(\mathbf{T}_{\lambda \prec \phi} \mathbf{T}_{\phi \succ \kappa} \right) \mathbf{T}_{\kappa \prec \phi} \\ &= \mathbf{T}_{\phi \succ \lambda} \left(\sum_{\nu \leq \kappa} \frac{B_{\lambda\kappa}^\nu}{[\nu \prec \dot{\lambda}]_\phi [\nu \prec \dot{\kappa}]_\phi} \mathbf{T}_{\lambda \succ \nu} \mathbf{T}_{\nu \prec \kappa} \right) \mathbf{T}_{\kappa \prec \phi} \end{aligned}$$

In the previous section we focused on the subspace of \mathcal{F}_λ which consisted of irreducible representations which occur in \mathcal{F}_ϕ and was denoted $\mathcal{F}_\lambda^\circ = \mathcal{F}_{\lambda|\phi}$. We now restrict attention to a yet smaller subspace of \mathcal{F}_λ :

$$\mathcal{F}_\lambda^\bullet = \text{Im}(\mathbf{T}_{\lambda \prec \phi})$$

This subspace is a copy of \mathcal{H}_ϕ^λ which sits canonically in the ϕ -part of \mathcal{F}_λ . Observe that the set of maps $\Delta = \Delta_{\lambda\kappa} = \{\mathbf{T}_{\lambda \succ \nu} \mathbf{T}_{\nu \prec \kappa}\}_{\nu \leq \lambda \wedge \kappa}$ when restricted to $\mathcal{F}_\kappa^\bullet$, forms a basis for G -maps $\mathcal{F}_\kappa^\bullet \rightarrow \mathcal{F}_\lambda^\bullet$. Indeed, these maps are independent after being composed with $\mathbf{T}_{\phi \succ \lambda}$ on the left and $\mathbf{T}_{\kappa \prec \phi}$ on the right and using lemma 3.3. We get the following identity on $\mathcal{F}_\lambda^\bullet$:

$$\begin{aligned} \mathbf{T}_{\lambda \prec \phi} \mathbf{T}_{\phi \succ \kappa} &= \sum_{\nu \leq \kappa} \frac{B_{\lambda\kappa}^\nu}{[\nu \prec \lambda]_\phi [\nu \prec \kappa]_\phi} \mathbf{T}_{\lambda \succ \nu} \mathbf{T}_{\nu \prec \kappa} \\ (3) \qquad &= \frac{A_{\lambda\kappa}}{[\kappa \prec \lambda]_\phi} \mathbf{T}_{\lambda \succ \kappa} + \{\text{terms with } \nu < \kappa\} \end{aligned}$$

Denote the coefficient of an operator S w.r.t to a basis element $D \in \Delta$ by $\langle S, D \rangle_\Delta$.

Definition 3.4. *A triple $\kappa \leq \lambda \leq \eta$ is called good if*

$$\langle \mathbf{T}_{\lambda \prec \eta} \mathbf{T}_{\eta \succ \kappa}, \mathbf{T}_{\lambda \succ \kappa} \rangle_\Delta = [\lambda \prec \eta \upharpoonright \kappa]_\Phi$$

Combining definition 3.4 with (3) we see that Theorem 4 is equivalent to:

Theorem 5. *$\kappa \leq \lambda \leq \phi$ is a good triple.*

This theorem, the heart of the matter, will keep us busy for the rest of the section. We would like to take a small pause and explain the strategy which we undertake. The idea is to show that it is enough to find a path connecting λ and ϕ in the segment $[\lambda, \phi]$ which can be paved with good triples, and then exhibit such path. More precisely, we follow three steps:

- (1) Given $\lambda_0 \leq \lambda_1 \leq \dots \leq \lambda_r \leq \phi$ such that $\kappa \leq \lambda_{i-1} \leq \lambda_i$ is good for all $1 \leq i \leq r$ and also $\kappa \leq \lambda_r \leq \phi$ is good, implies that so is $\kappa \leq \lambda_0 \leq \phi$.
- (2) $\kappa \leq \lambda_1 \leq \lambda_2$ is good whenever λ_2 covers λ_1 (i.e. $[\lambda_1, \lambda_2] = \{\lambda_1, \lambda_2\}$) and has the same rank.
- (3) $\kappa \leq \epsilon \leq \phi$ is good whenever ϵ and ϕ are symmetric.

Note that a Jordan-Hölder sequence of types from λ to a symmetric type ϵ of the same rank gives an appropriate path: $\lambda = \lambda_0 \leq \dots \leq \lambda_r = \epsilon \leq \phi$.

Step 1

Lemma 3.5. *If $\kappa \leq \lambda \leq \theta$ and $\kappa \leq \theta \leq \phi$ are good so is $\kappa \leq \lambda \leq \phi$.*

Proof.

$$\begin{aligned}
\langle \mathbf{T}_{\lambda \prec \phi} \mathbf{T}_{\phi \succ \kappa}, \mathbf{T}_{\lambda \succ \kappa} \rangle_{\Delta} &= \frac{1}{[\lambda \prec \theta]_{\phi}} \langle (\mathbf{T}_{\lambda \prec \theta} \mathbf{T}_{\theta \prec \phi}) \mathbf{T}_{\phi \succ \kappa}, \mathbf{T}_{\lambda \succ \kappa} \rangle_{\Delta} \\
&= \frac{[\theta \prec \dot{\phi} \cap \kappa]}{[\lambda \prec \dot{\theta}]_{\phi}} \langle \mathbf{T}_{\lambda \prec \theta} (\mathbf{T}_{\theta \prec \kappa} + \text{lower terms}), \mathbf{T}_{\lambda \succ \kappa} \rangle_{\Delta} \\
&= \frac{[\theta \prec \dot{\phi} \cap \kappa]}{[\lambda \prec \dot{\theta}]_{\phi}} \langle \mathbf{T}_{\lambda \prec \theta} \mathbf{T}_{\theta \prec \kappa}, \mathbf{T}_{\lambda \succ \kappa} \rangle_{\Delta} \\
&= \frac{[\theta \prec \dot{\phi} \cap \kappa][\lambda \prec \dot{\theta}]_{\kappa}}{[\lambda \prec \dot{\theta}]_{\phi}} = \frac{[\lambda \prec \dot{\theta} \prec \dot{\phi} \cap \mu]}{[\lambda \prec \dot{\theta}]_{\phi}} = [\lambda \prec \dot{\phi} \cap \kappa]_{\phi}
\end{aligned}$$

□


Using this lemma r times completes the proof of step 1. Note that the lemma can be used successively only from the 'top' to the 'bottom'.

Step 2

Let λ_1 and λ_2 be types of same rank and assume that λ_2 covers λ_1 . Let $\kappa \leq \lambda_1$. We want to show that $\langle \mathbf{T}_{\lambda_1 \prec \lambda_2} \mathbf{T}_{\lambda_2 \succ \kappa}, \mathbf{T}_{\lambda_1 \succ \kappa} \rangle_{\Theta} = [\lambda_1 \prec \lambda_2 \cap \kappa]$. However, the assumption that λ_1 and λ_2 have the same rank guarantees that any module of type λ_2 containing a module of type λ_1 which is transversal w.r.t. a module of type κ inherits this transversality. Hence, the requirement to avoid κ is redundant and $[\lambda_1 \prec \lambda_2 \cap \kappa] = [\lambda_1 \prec \lambda_2]$. We start by expanding the product $\mathbf{T}_{\lambda_1 \prec \lambda_2} \mathbf{T}_{\lambda_2 \succ \kappa}$:

$$(4) \quad \mathbf{T}_{\lambda_1 \prec \lambda_2} \mathbf{T}_{\lambda_2 \succ \kappa} = a_{\lambda_1} \mathbf{T}_{\lambda_1 \leftrightarrow \lambda_1 \leftrightarrow \kappa} + a_{\lambda_2} \mathbf{T}_{\lambda_1 \leftrightarrow \lambda_2 \leftrightarrow \kappa}$$

The assumptions on λ_1 and λ_2 assures that no other terms appear in (4). Evidently $\mathbf{T}_{\lambda_1 \succ \kappa} = \mathbf{T}_{\lambda_1 \leftrightarrow \lambda_1 \leftrightarrow \kappa}$ and $a_{\lambda_1} = [\lambda_1 \prec \lambda_2]$. We are therefore reduced to showing that the Δ expansion of the second term in (4) does not contain a multiple of $\mathbf{T}_{\lambda_1 \succ \kappa}$. This is accomplished by claim 3.6. Let κ_1 be the unique type which can (possibly) complete a cartesian diagram (see the first part of claim A.4 in appendix A):



Claim 3.6. $(\mathbf{T}_{\lambda_1 \leftrightarrow \lambda_2 \leftrightarrow \kappa})_{|\mathcal{F}_{\kappa}^{\bullet}} \in \mathbb{C} \cdot (\mathbf{T}_{\lambda_1 \succ \kappa_1} \mathbf{T}_{\kappa_1 \prec \kappa})_{|\mathcal{F}_{\kappa}^{\bullet}}$.

Proof. We shall prove the equivalent statement:

$$\mathbf{T}_{\phi \succ \lambda_1} \mathbf{T}_{\lambda_1 \leftrightarrow \lambda_2 \leftrightarrow \kappa} \in \mathbb{C} \cdot \mathbf{T}_{\phi \succ \lambda_1} \mathbf{T}_{\lambda_1 \succ \kappa_1} \mathbf{T}_{\kappa_1 \prec \kappa} \quad (= \mathbb{C} \cdot \mathbf{T}_{\phi \succ \kappa_1} \mathbf{T}_{\kappa_1 \prec \kappa} \text{ by lemma 3.3})$$

We begin by applying the r.h.s. to a cyclic element $\delta_{x_0} \in \mathcal{F}_\kappa$:

$$\begin{aligned} [\mathbf{T}_{\phi \succ \kappa_1} \mathbf{T}_{\kappa_1 \prec \kappa} \delta_{x_0}](y_0) &= \sum_{\substack{z \subset y_0 \\ \tau(z) = \kappa_1}} [\mathbf{T}_{\kappa_1 \prec \kappa} \delta_{x_0}](z) = \sum_{\substack{z \subset y_0 \\ \tau(z) = \kappa_1}} \sum_{\substack{x \supset z \\ \tau(x) = \phi}} \delta_{x_0}(x) \\ &= \sum_{\substack{z \subset y_0 \\ \tau(z) = \kappa_1}} \mathbf{1}_{\{z | z \subset x_0, \tau(z) = \kappa_1\}} = |\{z | z \subset y_0 \cap x_0, \tau(z) = \kappa_1\}| \\ &= \begin{cases} \binom{\kappa}{\kappa_1} & \text{if } y_0 \supset x_0 \\ 1 & \text{if } \tau(y_0 \cap x_0) = \kappa_1 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

Let w_0 , x_0 and y_0 be submodules of F such that $w_0 \subset x_0 \cap y_0$ with types $\tau(w_0) = \kappa_1$, $\tau(x_0) = \kappa$ and $\tau(y_0) = \phi$. Define:

$$B = B_{x_0, y_0, w_0} = \{z | z \subset y_0, \tau(z) = \lambda_1, \tau(z + x_0) = \lambda_2, z \cap x_0 = w_0\}$$

By the second part of claim A.4:

$$B = \{z | w_0 \subset z \subset y_0, \tau(z) = \lambda_1, \tau(\wp z + \wp x_0) = \wp \lambda_2, \wp z \cap \wp x_0 = \wp w_0\}$$

This description of B together with the symmetricity of y_0 implies that $|B|$ depends only on the types of $\wp x_0$ and w_0 , denote it by $b_{\wp \kappa, \kappa_1}$.

Applying the l.h.s. to δ_{x_0} yields:

$$\begin{aligned} [\mathbf{T}_{f \succ \lambda_1} \mathbf{T}_{\lambda_1 \leftrightarrow \lambda_2 \leftrightarrow \kappa} \delta_{x_0}](y_0) &= \sum_{\substack{z \subset y_0 \\ \tau(z) = \lambda_1}} [\mathbf{T}_{\lambda_1 \leftrightarrow \lambda_2 \leftrightarrow \kappa} \delta_{x_0}](z) = \sum_{\substack{z \subset y_0 \\ \tau(z) = \lambda_1}} \sum_{\substack{x \\ \tau(z+x) = \lambda_2}} \delta_{x_0}(x) \\ &= |\{z | z \subset y_0, \tau(z) = \lambda_1, \tau(z + x_0) = \lambda_2\}| \\ &= \begin{cases} \binom{\kappa}{\kappa_1} b_{\wp \kappa, \kappa_1} & \text{if } y_0 \supset x_0 \\ b_{\wp \kappa, \kappa_1} & \text{if } \tau(y_0 \cap x_0) = \kappa_1 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

Hence, the two operators differ by a constant. \square

Step 3

We should prove that the triple $\kappa \leq \epsilon \leq \phi$ is good when both ϵ and ϕ are symmetric types. This is precisely the assertion of theorem 5 in [BO04]. In particular see relation (8) in the proof. The only delicate point which deserves a remark, is the duality axiom which is used in the proof and should be justified. Indeed, in a module of type κ , the number of submodules of type α equals to the number of submodules of co-type α (cf. [Mac95, Chapter 4]).

4. FOURIER DECOMPOSITION OF THE HECKE ALGEBRA

In this section we collect the results obtained so far and shift them to the limit space, which was our original problem.

4.1. **Explicit Formulas.** Let $q = |\mathcal{O}/\wp|$. For $m, n \in \mathbb{N}$ Set

$$[n]_q = 1 - q^{-n}, \quad [0]_q = 1,$$

$$[n]_q! = [n]_q [n-1]_q \cdots [1]_q, \quad \begin{bmatrix} m \\ n \end{bmatrix}_q = \frac{[n]_q!}{[m]_q! [n-m]_q!}$$

The subscript q will be occasionally omitted from the notation. Using the embedding of Λ_m in \mathbb{Z}^∞ via the transposed coordinates, and using the standard bilinear form there we will write

$$\langle \lambda', \nu' \rangle = \sum \lambda'_i \nu'_i$$

Claim 4.1.

$$(1) \quad \begin{pmatrix} \nu \\ \lambda \end{pmatrix} = q^{\langle \nu' - \lambda', \lambda' \rangle} \prod_{i \geq 1} \begin{bmatrix} \nu'_i - \lambda'_{i+1} \\ \nu'_i - \lambda'_i \end{bmatrix}$$

$$(2) \quad [\lambda \prec \dot{\nu}]_\phi = q^{\langle \phi' - \nu', \nu' - \lambda' \rangle} \begin{bmatrix} \phi'_1 - \lambda'_1 \\ \phi'_1 - \nu'_1 \end{bmatrix} \prod_{i \geq 1} \begin{bmatrix} \nu'_i - \lambda'_{i+1} \\ \nu'_i - \nu'_{i+1} \end{bmatrix}$$

$$(3) \quad [\nu \prec \dot{\phi} \cap \lambda]_\Phi = q^{\langle \Phi' - \phi', \phi' - \nu' \rangle} \begin{bmatrix} \Phi'_1 - \nu'_1 - \lambda'_1 \\ \phi'_1 - \nu'_1 \end{bmatrix}$$

Proof. A basic quantity, which all other quantities are scaled to, is the size of $\text{Hom}(\oplus \mathcal{O}/\wp^{\lambda_i}, \oplus \mathcal{O}/\wp^{\nu_i})$ which is denoted and computed by

$$\text{hom}(\lambda, \nu) = q^{\langle \lambda, \nu \rangle}$$

The subset of all injective morphisms will be denoted $\text{Hom}^{1-1}(\oplus \mathcal{O}/\wp^{\lambda_i}, \oplus \mathcal{O}/\wp^{\nu_i})$ and we will use

$$\text{hom}^{1-1}(\lambda, \nu) = |\text{Hom}^{1-1}(\oplus \mathcal{O}/\wp^{\lambda_i}, \oplus \mathcal{O}/\wp^{\nu_i})|$$

For a given type λ , $\text{hom}^{1-1}(\lambda, \lambda)$ is computed in [Mac95, Chapter 4].

A similar computation shows that

$$\text{hom}^{1-1}(\lambda, \nu) = \prod_{i \geq 1} \frac{[\nu'_i - \lambda'_{i+1}]!}{[\nu'_i - \lambda'_i]!} \text{hom}(\lambda, \nu)$$

Observe that the map

$$\text{Hom}^{1-1}(\oplus \mathcal{O}/\wp^{\lambda_i}, \oplus \mathcal{O}/\wp^{\nu_i}) \rightarrow \text{Gr}(\lambda, \oplus \mathcal{O}/\wp^{\nu_i}), \quad \psi \mapsto \text{Im}(\psi)$$

is $\text{hom}^{1-1}(\lambda, \lambda)$ to one, thus

$$\begin{pmatrix} \nu \\ \lambda \end{pmatrix} = \frac{\text{hom}^{1-1}(\lambda, \nu)}{\text{hom}^{1-1}(\lambda, \lambda)}$$

and (1) follows. Given an \mathcal{O} -module f of type ϕ , counting in two ways the size of the set

$$\{x, y < f \mid x < y, \tau(x) = \lambda, \tau(y) = \nu\}$$

gives

$$\binom{\phi}{\lambda} [\lambda \prec \dot{\nu}]_{\phi} = \binom{\phi}{\nu} \binom{\nu}{\lambda}$$

which (given (1)) readily proves (2).

In order to prove (3) we need a little preparation. Let F be a module of type Φ . Let $z < F$ be a fixed module of type λ . Denote by F_1 a module of type k^{λ_1} containing z . Observe that F_1 is a direct summand of F . Fix a complimentary direct summand to F_1 in F and denote it by F_2 , thus

$$F = F_1 \oplus F_2$$

Denote the corresponding projections by p_1 and p_2 . Assume that a type ν is given. We denote

$$X = \{x < F \mid \tau(x) = \nu, x \cap z = 0\}, \quad [\dot{\nu} \pitchfork \lambda] = |X|$$

Observe that the map

$$\mathrm{Hom}(\oplus \mathcal{O}/\wp^{\nu_i}, F_1) \oplus \mathrm{Hom}^{1-1}(\oplus \mathcal{O}/\wp^{\nu_i}, F_2) \hookrightarrow \mathrm{Hom}(\oplus \mathcal{O}/\wp^{\nu_i}, F) \rightarrow X,$$

$$(\psi_1, \psi_2) \mapsto \mathrm{Im}(\psi_1 + \psi_2)$$

is $\mathrm{hom}^{1-1}(\nu, \nu)$ to one. This map is indeed into X as

$$\mathrm{Im}(\psi_1 \oplus \psi_2) \cap z \subset \mathrm{Im}(\psi_1 \oplus \psi_2) \cap F_1 \simeq \mathrm{Ker}(\psi_2) = 0$$

It is onto X as for a given $\psi \in \mathrm{Hom}^{1-1}(\oplus \mathcal{O}/\wp^{\nu_i}, F)$, with $\mathrm{Im}(\psi) \in X$ we have $\mathrm{Ker}(p_2 \circ \psi) \simeq \mathrm{Im}(\psi) \cap F_1$ which is 0, as the \wp -torsion of F_1 is equal the \wp -torsion of z . We get that

$$[\dot{\nu} \pitchfork \lambda] = \binom{k^{n-\lambda_1}}{\nu} \mathrm{hom}(\nu, k^{\lambda_1})$$

Counting in two ways the size of the set

$$\{x, y < F \mid x < y, y \cap z = 0, \tau(x) = \nu, \tau(y) = \phi\}$$

gives

$$[\dot{\phi} \pitchfork \lambda] \binom{\phi}{\nu} = [\dot{\nu} \pitchfork \lambda] [\nu \prec \dot{\phi} \pitchfork \lambda]$$

and (3) follows. □

4.2. Transition matrices. We are finally in a position to collect the pieces, and write down explicitly the transition matrix between the delta functions basis of \mathcal{H}_m and the idempotents of \mathcal{H}_m (the spherical functions). In order to do that we introduce an intermediate basis, which is the limit of the (normalized image in \mathcal{H}_m) of the finite levels cellular bases. We define

$$\bar{\mathbf{c}}_\lambda = \lim_k \frac{\mathbf{c}_\lambda}{\binom{\Phi}{\phi}}$$

Thus obtaining

$$\begin{aligned} \delta_\lambda &= \sum_{\kappa \geq \lambda \geq \varnothing \kappa} \hat{\mu}(\lambda, \kappa) \bar{\mathbf{c}}_\kappa \\ &= \sum_{\kappa \geq \lambda \geq \varnothing \kappa} (-1)^{|\nu| - |\lambda|} q^{n(\nu) - n(\lambda)} \prod \left[\begin{matrix} \nu'_i - \nu'_{i+1} \\ \nu'_i - \lambda'_i \end{matrix} \right] \bar{\mathbf{c}}_\kappa \\ \bar{\mathbf{c}}_\kappa &= \sum_{\nu \leq \kappa} \frac{[\nu \prec \dot{\kappa}]_\phi [\kappa \prec \dot{\phi} \uparrow \nu]_\Phi}{\binom{\Phi}{\phi}} \mathbf{e}_\nu \\ &= \sum_{\nu \leq \kappa} q^{-(n-2m)|\kappa| - m|\nu| - \langle \kappa', \kappa' - \nu' \rangle} \frac{\begin{bmatrix} m - \nu'_1 \\ m - \kappa'_1 \end{bmatrix} \begin{bmatrix} n - \kappa'_1 - \nu'_1 \\ m - \kappa'_1 \end{bmatrix}}{\begin{bmatrix} n \\ m \end{bmatrix}} \prod \left[\begin{matrix} \kappa'_i - \nu'_{i+1} \\ \kappa'_i - \nu'_i \end{matrix} \right] \mathbf{e}_\nu \end{aligned}$$

Which combined together give the desired $(\delta - \mathbf{e})$ transition matrix.

Remark 4.2. *The transition matrices $\delta \leftrightarrow \bar{\mathbf{c}} \leftarrow \mathbf{e}$ are given explicitly by a combinatorial data. Examples imply that this should also be the case for the transition matrix $\bar{\mathbf{c}} \rightarrow \mathbf{e}$, and it would be interesting to find such interpretation.*

5. FINAL REMARKS

5.1. Grassmannians over archimedean local fields. The Grassmann representation over an archimedean local field (i.e \mathbb{R} or \mathbb{C}) was considered by James and Constantine in [JC74]. The decomposition to irreducible representations consists of the same indexing set Λ_m as the non-archimedean case discussed here. An interesting question is whether there exist a natural correspondence between the irreducible indexed by λ going over all the places of a global field. This question is addressed in [Onn04]. It turns out that the quantum Grassmannian interpolates between the Grassmannians over local fields.

5.2. Nilpotent representations of $\mathrm{GL}_n(\mathcal{O})$. The irreducible representations considered here were studied by Gregory Hill in [Hil94]. They seem to play an important role in the representation theory of $\mathrm{GL}_n(\mathcal{O})$. Proposition 3.1 was proved in [Hil94] using quite involved calculations. We refer the reader to [BO04] for a general discussion on Hecke algebras arising from Grassmannians in which a generalization of the latter proposition is given in a natural way.

APPENDIX A. LOCAL RINGS AND DVR'S

In this appendix we prove some claims regarding local rings. All modules under consideration are assumed to be of finite rank.

A.1. The Euler characteristic of a module. Let M be a module over a ring R . Denote by $\mathcal{L}(M)$ the lattice of submodules of M . One naturally associates a simplicial complex to M , denoted $C(M)$, with vertices consisting of the non-trivial submodules of M (all but 0 and M). The simplices of $C(M)$ are given by

$$\{x_1, x_2, \dots, x_m \mid 0 \subset x_1 \subset x_2 \subset \dots \subset x_m \subset M\}$$

We denote the Euler characteristic of $C(M)$ by $\chi(M)$. In case M is a finite module, we denote the incidence algebra associated with $\mathcal{L}(M)$ by $I(M)$. $\chi(M)$ is related to the Möbius function, $\mu \in I(M)$, by (cf. [Rot64, Proposition 6])

$$\mu(0, M) = \chi(M)$$

The following Proposition is probably well known. We give here a direct proof.

Proposition A.1. *Let R be a local ring with maximal ideal \wp , and let M be a finite R -module with $\wp M \neq 0$. Then $C(M)$ is contractible. In particular, $\chi(M) = 0$.*

Proof. Denote $C = C(M)$, and $D \subseteq C$ the subcomplex spanned by the subset of vertices $\{x \mid \wp M \subseteq x \subset M\} \subseteq C_0$. D is a cone over the vertex $\wp M$, hence contractible. The function

$$\varphi : \mathcal{L}(M) \rightarrow \mathcal{L}(M), \quad \varphi(x) = x + \wp M$$

extends to a retraction $\varphi_* : C \rightarrow D$. The function

$$\Psi : C \times [0, 1] \rightarrow C \quad \Psi(c, t) = (1 - t)c + t\varphi_*(c)$$

establishes a deformation retract from C to D . Therefore C is contractible. \square

Assume that $R/\wp \simeq \mathbb{F}_q$. Proposition A.1 combined with a well known formula giving the Euler characteristic of the Tits building associated to $GL_d(\mathbb{F}_q)$, gives For every module type λ ,

$$\chi(\lambda) = \begin{cases} 0 & \wp \cdot \lambda \neq 0 \\ (-1)^{\dim_{\mathbb{F}_q}(\lambda)} q^{\binom{\dim_{\mathbb{F}_q}(\lambda)}{2}} & \wp \cdot \lambda = 0 \end{cases}$$

A.2. Let R be a local ring with a maximal ideal \wp and let x, y be R -modules. Let $\bar{x} = x/\wp x$. By Nakayama's lemma $x \rightarrow y$ is onto if and only if the induced map $\bar{x} \rightarrow \bar{y}$ is onto. Equivalently $\text{rk}(x) = \dim x/\wp x$.

Claim A.2. *Let R be a local ring with maximal ideal \wp . Let z be an R -module and x, y two submodules of z . Then:*

$$\text{rk}(x + y) = \text{rk}(x) + \text{rk}(y) - \text{rk}(x \cap y) + \dim \left(\frac{\wp x \cap \wp y}{\wp(x \cap y)} \right)$$

Proof. Using the equalities $\wp(x + y) = \wp x + \wp y$ and $\wp(x \oplus y) = \wp x \oplus \wp y$, we obtain a commutative diagram with exact columns and rows (5) and exact sequence (6):

$$(5) \quad \begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & \wp x \cap \wp y & \rightarrow & \wp x \oplus \wp y & \rightarrow & \wp x + \wp y \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & x \cap y & \rightarrow & x \oplus y & \rightarrow & x + y \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & \frac{x \cap y}{\wp x \cap \wp y} & \dashrightarrow & \frac{x \oplus y}{\wp x \oplus \wp y} & \dashrightarrow & \frac{x + y}{\wp(x + y)} \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

(the exactness of the dashed row follows from the obvious exactness of the other rows and columns)

$$(6) \quad 0 \rightarrow \frac{\wp x \cap \wp y}{\wp(x \cap y)} \rightarrow \frac{x \cap y}{\wp(x \cap y)} \rightarrow \frac{x \cap y}{\wp x \cap \wp y} \rightarrow 0$$

and obtain:

$$\begin{aligned} \text{rk}(x) + \text{rk}(y) &= \text{rk}(x \oplus y) \stackrel{(5)}{=} \text{rk}(x + y) + \dim \frac{x \cap y}{\wp x \cap \wp y} \\ &\stackrel{(6)}{=} \text{rk}(x + y) + \text{rk}(x \cap y) - \dim \left(\frac{\wp x \cap \wp y}{\wp(x \cap y)} \right) \end{aligned}$$

□

Claim A.3. *Let R be a local ring with maximal ideal \wp . Let z be an R -module and x, y two submodules of z . Then:*

$$\text{rk}(x) = \text{rk}(x + y) \iff \text{rk}(x \cap y) = \text{rk}(y) \text{ and } \wp x \cap \wp y = \wp(x \cap y)$$

Proof. (\Leftarrow) clear. (\Rightarrow) if $\text{rk}(x) = \text{rk}(x + y)$ we have that the two non-negative terms $\text{rk}(y) - \text{rk}(x \cap y)$ and $\dim\left(\frac{\wp x \cap \wp y}{\wp(x \cap y)}\right)$ must sum up to zero. \square

We now specialize to the situation to which we apply these assertions.

Claim A.4. *Let \mathcal{O} be a discrete valuation ring with maximal ideal \wp . Let z be a finite \mathcal{O} -module and x, y two submodules of z . Assume that $\text{rk}(x + y) = \text{rk}(x)$ and $x + y$ covers x .*

- (1) $\tau(x \cap y)$ depends only on $\tau(x)$, $\tau(y)$ and $\tau(x + y)$.
- (2) $\tau(x + y) = \lambda \iff \tau(\wp x + \wp y) = \wp \lambda$, $\wp x \cap \wp y = \wp(x \cap y)$ and $\wp(x \cap y) \subsetneq \wp y$.

Proof. One immediately verifies that for a module w , $\tau(w) = \lambda \iff \dim \bar{w} = \lambda_1 = \text{rk}(\lambda)$ and $\tau(\wp w) = \wp \lambda$.

- (1) There exist a unique i such that $\wp^i(x + y)/\wp^i x \neq (0)$. By the isomorphism $\wp^i(x + y)/\wp^i x \simeq \wp^i y/\wp^i(x \cap y)$ it is also the unique i for which $\wp^i y/\wp^i(x \cap y) \neq (0)$.
- (2) (\Rightarrow) Assume $\tau(x + y) = \lambda$. Clearly $\tau(\wp x + \wp y) = \wp \lambda$, and by Claim A.3 also $\wp x \cap \wp y = \wp(x \cap y)$. Since $x \cap y \subsetneq y$ and have the same rank, the dimensions of $\overline{x \cap y}$ and \bar{y} must be the same and $\wp(x \cap y) \subsetneq \wp y$.
 (\Leftarrow) The data on the right together with claim A.3 implies that $\tau(\wp(x + y)) = \wp \lambda$ and $\text{rk}(x + y) = \text{rk}(x)$. This implies that $\tau(x + y) = \lambda$. \square

REFERENCES

- [BO04] Uri Bader and Uri Onn, *On cellular Hecke algebras associated with generalized Grassmannians*, preprint (2004).
- [Hil94] Gregory Hill, *On the nilpotent representations of $GL_n(\mathcal{O})$* , *Manuscripta Math.* **82** (1994), no. 3-4, 293–311.
- [JC74] A. T. James and A. G. Constantine, *Generalized Jacobi polynomials as spherical functions of the Grassmann manifold*, *Proc. London Math. Soc.* (3) **29** (1974), 174–192.
- [Mac95] I. G. Macdonald, *Symmetric functions and Hall polynomials*, second ed., Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1995, With contributions by A. Zelevinsky, Oxford Science Publications.
- [Onn03] Uri Onn, *On the Grassmann Representation of $GL(n)$* , Thesis (2003), 44.

- [Onn04] _____, *From p-adic to real Grassmannians via the quantum*, preprint (2004).
- [Rot64] Gian-Carlo Rota, *On the foundations of combinatorial theory. I. Theory of Möbius functions*, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **2** (1964), 340–368.

TECHNION, HAIFA 32000, ISRAEL

E-mail address: bader@tx.technion.ac.il uonn@science.uva.nl