

PROPERTY (T) AND RIGIDITY FOR ACTIONS ON BANACH SPACES

URI BADER, ALEX FURMAN, TSACHIK GELANDER, AND NICOLAS MONOD

ABSTRACT. We study property (T) and the fixed point property for actions on L^p and other Banach spaces. We show that property (T) holds when L^2 is replaced by L^p (and even a subspace/quotient of L^p), and that in fact it is independent of $1 \leq p < \infty$. We show that the fixed point property for L^p follows from property (T) when $1 < p < 2 + \varepsilon$. For simple Lie groups and their lattices, we prove that the fixed point property for L^p holds for any $1 < p < \infty$ if and only if the rank is at least two. Finally, we obtain a superrigidity result for actions of irreducible lattices in products of general groups on superreflexive Banach spaces.

1. INTRODUCTION AND THE MAIN RESULTS

1.1. Since its introduction by Kazhdan in [Ka], property (T) became a fundamental concept in mathematics with a wide range of applications to such areas as: Structure of infinite groups – finite generation of higher rank lattices [Ka], obstruction to free or amalgamated splittings [Wa], [A], [M4] structure of normal subgroups [M2] etc.; combinatorics – construction of expanders [M1] (see [Lu]); operator algebras – factors with trivial fundamental group [C], rigidity theorems related to Orbit Equivalence [Po],[Hj]; the Banach-Ruziewicz problem [M3],[Su]; rigidity in dynamics [FM1],[FM2],[N]. It has also been an important tool in providing interesting (counter) examples – for the von Neumann conjecture [Gr1, 5.6] and in the context of Baum-Connes conjecture [Gr2].

Property (T) , initially defined in terms of unitary representations, turned out to be equivalent to Serre's property (FH) – a fixed point property for affine isometric actions on Hilbert spaces (the equivalence was proved by Delorme [D] and Guichardet [Gu]). Some of the above applications use this latter characterization. Quite recently Shalom [Sh] studied such affine isometric actions on Hilbert spaces in the context of lattices in general products of locally compact groups, deducing new rigidity results. For further details and more references on these topics we suggest the monograph [HV] and the forthcoming [BHV].

1.2. Motivated by these broad themes: property (T) , property (FH) , lattices in semisimple groups and in general products, we study similar notions in the broader

Date: 9th June 2005.

U.B. partially supported by ISF grant 100146, A.F. partially supported by NSF grant DMS-0094245, T.G. partially supported by NSF grants DMS-0404557, N.M. partially supported by FNS grant 8220-067641 and NSF grant DMS 0204601.

framework of Banach spaces rather than Hilbert spaces. Some of the results below apply to general *superreflexive* Banach spaces, whilst some are specific to the subclass of $L^p(\mu)$ -spaces with $1 < p < \infty$. (A Banach space is superreflexive if it admits an equivalent uniformly convex norm, see Proposition 2.3 below.)

One of the motivations to consider such questions came from the work of Fisher and Margulis [FM1], [FM2], in which an L^p analogue of property (T) with $p \gg 2$ allowed them to weaken smoothness assumptions in their results.

1.3. Let G be a topological group and B a Banach space. By a *linear isometric G -representation* on B , we shall mean a continuous homomorphism $\varrho : G \rightarrow \mathbf{O}(B)$ where $\mathbf{O}(B)$ denotes the (“orthogonal”) group of all invertible linear isometries $B \rightarrow B$ (see Lemma 2.4 for a clarification of the continuity assumption). We say that such a representation *almost has invariant vectors* if

$$(1.1) \quad \forall \text{ compact subset } K \subseteq G, \quad \inf_{\|v\|=1} \text{diam}(\varrho(K)v) = 0.$$

Denote by $B^{\varrho(G)}$ the closed subspace of G -fixed vectors; the G -representation ϱ descends to a linear isometric G -representation ϱ' on $B' = B/B^{\varrho(G)}$ (see Remark 2.11 for more details in the case of superreflexive spaces). We shall use the following as a Banach space analogue of Kazhdan’s property (T):

Definition 1.1. Let B be a Banach space. A topological group G is said to have property (T_B) if for any continuous linear isometric G -representation $\varrho : G \rightarrow \mathbf{O}(B)$ the quotient G -representation $\varrho' : G \rightarrow \mathbf{O}(B/B^{\varrho(G)})$ does not almost have G -invariant vectors.

Note that if B is a Hilbert space, ϱ' is isomorphic to the restriction of ϱ to the orthogonal complement $(B^{\varrho(G)})^\perp$ of the subspace of $\varrho(G)$ -invariants. Thus for Hilbert spaces the above definition agrees with Kazhdan’s property (T).

Let μ be a σ -finite measure on a standard Borel space (X, \mathcal{B}) . We are most interested in the family $L^p(\mu)$, $1 < p < \infty$, of Banach spaces, which are close relatives of Hilbert spaces. They also possess a rich group of linear isometries $\mathbf{O}(L^p(\mu))$.

Theorem A. *Let G be a locally compact second countable group. If G has Kazhdan’s property (T) then G has property (T_B) for Banach spaces B of the following types:*

- (i) $L^p(\mu)$ for any σ -finite measure μ and any $1 \leq p < \infty$.
- (ii) A closed subspace of $L^p(\mu)$ for any $1 < p < \infty$, $p \neq 4, 6, 8, \dots$
- (iii) A quotient space of $L^p(\mu)$ for any $1 < p < \infty$, $p \neq \frac{4}{3}, \frac{6}{5}, \frac{8}{7}, \dots$

If G has $(T_{L^p([0,1])})$ for some $1 < p < \infty$ then G has Kazhdan’s property (T).

1.4. Next we consider group actions by isometries on Banach spaces. Such actions are always affine, with the linear part being isometric as well.

Definition 1.2. We say that G has property (F_B) if any continuous action of G on B by isometries has a G -fixed point.

When B is a Hilbert space this is precisely Serre's property (FH) . Delorme [D] and Guichardet [Gu] proved that properties (T) and (FH) are equivalent. Below we summarize the relations between properties (T) and (F_B) which hold for general groups.

Theorem 1.3. *For a locally compact second countable group G we have*

- (1) (F_B) implies (T_B) for any Banach space B .
- (2) (T) implies (F_B) for closed subspaces B of $L^p(\mu)$ where $0 < p \leq 2$.[†]
- (3) (T) also implies (F_B) for closed subspaces of $L^p(\mu)$ for $2 \leq p < 2 + \varepsilon$, where $\varepsilon = \varepsilon(G) > 0$ might depend on the Kazhdan group G .

Remarks 1.4. (1) is essentially due to Guichardet [Gu] as his proof of $(FH) \Rightarrow (T)$ applies to all Banach spaces. We give two proofs for (2) reducing the problem, in both, to one of the proofs of $(T) \Rightarrow (FH)$. We note that the particular case of $p = 1$ in (2) is one of the results of [RS]. Statement (3) is due to Fisher and Margulis (unpublished). With their kind permission we have included their argument here (see Section 3.3).

The above results imply that any locally compact group G with Kazhdan's property (T) has property (T_{L^p}) for all $1 < p < \infty$, and has the fixed point property (F_{L^p}) for $1 < p < 2 + \varepsilon(G)$. It turns out, however, that many Kazhdan groups (e.g. hyperbolic ones) do not have property (F_{L^p}) for large values of p :

Indeed, in his study of L^p -homology Pansu [Pa] proved that $\mathbf{Sp}_{n,1}(\mathbf{R})$ and cocompact lattices in these groups have a non-trivial first L^p -cohomology $L^p H^1$ for all $p > 4n + 1$. This is equivalent to asserting that for $p > 4n + 1$ these groups admit fixed-point-free affine isometric actions on $L^p(G)$ with linear part being the regular representation. Hence these groups do not have property (F_{L^p}) for $p > 4n + 1$, whilst enjoying (T) .

More generally, $L^p H^1(\Gamma)$ and hence $H^1(\Gamma, \ell^p \Gamma)$ is non-zero for any non-elementary hyperbolic group when p is large enough. Indeed, Bourdon and Pajot identify this cohomology with a Besov space of functions on the boundary, which they prove to be non-trivial as soon as p is larger than the Hausdorff dimension of an Ahlfors-regular metric on the boundary, see Corollaire 6.2 in [BP]. Again, this contradicts (F_{L^p}) for large p .

More recently, using Mineyev's homological bicomings [Mi], Yu [Y] gave a very short proof that any hyperbolic group Γ admits a *proper* action by isometries on $\ell^p(\Gamma \times \Gamma)$ if p is large enough. This is a strong negation of (F_{L^p}) for hyperbolic groups and all their infinite subgroups.

1.5. It is often remarked that property (T) for (simple) higher rank Lie groups and their lattices is more robust than property (T) enjoyed by the rank one groups $\mathbf{Sp}_{n,1}(\mathbf{R})$ and many other Gromov hyperbolic groups. In view of the preceding discussion of hyperbolic groups, the following result might be viewed as yet another evidence supporting this view.

[†]For subspaces B of $L^1(\mu)$ and of pseudo-normed spaces $L^r(\mu)$, $0 < r < 1$, property (F_B) should read: any isometric G -action on B has bounded orbits.

Theorem B. *Let $G = \prod_{i=1}^m \mathbf{G}_i(k_i)$, where k_i are local fields (of any characteristic), $\mathbf{G}_i(k_i)$ are k_i -points of Zariski connected isotropic simple k_i -algebraic groups \mathbf{G}_i . Assume that each simple factor $\mathbf{G}_i(k_i)$ has k_i -rank ≥ 2 .*

Then G and the lattices in G have property (F_B) for all $L^p(\mu)$ -related spaces B as in (i)–(iii) in Theorem A, assuming $1 < p < \infty$.

1.6. A broader class of spaces in which we propose to study properties (T_B) and (F_B) consists of *superreflexive* spaces, which can be defined as topological vector spaces isomorphic to uniformly convex Banach spaces. In this context we consider linear representations (resp. affine actions) which are *uniformly equicontinuous*; more concretely, for any given norm compatible with the topology, the class of all such linear representations (resp. affine actions) is that of *uniformly bounded* linear representations (resp. *uniformly Lipschitz* affine actions). It turns out that such representations (resp. actions) can always be viewed as isometric with respect to some equivalent norm that is *simultaneously* uniformly convex and uniformly smooth (Proposition 2.13).

Note that whether a given linear G -representation almost contains invariant vectors or not, in the sense of (1.1), does not depend on a particular norm among all mutually equivalent norms. Hence we can make the following

Definition 1.5. Let B be a superreflexive topological vector space and G a locally compact second countable group. We say that G has property (\bar{T}_B) if for every uniformly equicontinuous linear representation ϱ of G on B the quotient G -representation on $B/B^{\varrho(G)}$ does not almost have invariant vectors.

Likewise, G has (\bar{F}_B) if every uniformly equicontinuous affine G -action on B has a fixed point.

Conjecture 1.6. *Higher rank groups $G = \prod \mathbf{G}_i(k_i)$ as in Theorem B and their lattices have property (\bar{F}_B) , and hence (\bar{T}_B) , for all superreflexive B .*

Remark 1.7. To support this conjecture let us point out the following:

(1) Much of our proof of Theorem B is done in the broad context of uniformly equicontinuous affine actions on general superreflexive spaces except for one argument – a version of relative property (T_B) , whose proof is special to L^p -related spaces.

(2) V. Lafforgue proved [Lg] that the group $\mathbf{PGL}_3(\mathbf{Q}_p)$ has property (\bar{T}_B) for all superreflexive B (his result is actually stronger, in that he allows linear representations with *slowly growing*, rather *uniformly bounded* Lipschitz norms). Combined with our proof of Theorem B it implies for example that $\mathbf{SL}_n(\mathbf{Q}_p)$, $n \geq 4$, has property (\bar{F}_B) .

(3) Y. Shalom has proved (unpublished) that for Hilbert spaces \mathcal{H} higher rank groups (and their lattices) have property $(\bar{F}_{\mathcal{H}})$, and hence $(\bar{T}_{\mathcal{H}})$, whilst rank one groups have neither $(\bar{F}_{\mathcal{H}})$ nor $(\bar{T}_{\mathcal{H}})$.

1.7. One way to generalize the context of semisimple (non-simple) Lie/algebraic groups is simply to consider general products $G = G_1 \times \cdots \times G_n$ of $n \geq 2$ arbitrary topological groups. In the absence of any assumption on the factors G_i , one can

still establish splitting results for uniformly equicontinuous affine G -actions on superreflexive spaces.

Theorem C. *Let $G = G_1 \times \cdots \times G_n$ be a product of topological groups with a continuous action by uniformly equicontinuous affine maps on a superreflexive topological vector space B without G -fixed point. Assume that the associated linear G -representation ρ does not almost have non-zero invariant vectors.*

Then there is a G -invariant closed complemented affine subspace $\underline{B} \subseteq B$ and an affine equicontinuous G -equivariant isomorphism $\underline{B} \cong B_1 \oplus \cdots \oplus B_n$, where each B_i is a superreflexive Banach space with an equicontinuous affine G -action factoring through $G \rightarrow G_i$.

Remarks 1.8. (1) If G has property (\bar{T}_B) then the assumption that ρ does not almost have invariant vectors is redundant.

(2) In the particular case where B is a Hilbert space and G locally compact acting by affine isometries, a stronger result was established by Shalom in [Sh]: One assumes only that the *affine* G -action does not almost have fixed points. We replace Shalom's Hilbertian approach with an analogue of the geometric method used in the splitting theorem of [Mo].

(3) This result can be reformulated in terms of the cohomology of the associated linear G -representation ρ on B as

$$H^1(G, B) \cong \bigoplus_{i=1}^n H^1(G_i, B^{\otimes (\prod_{j \neq i} G_j)}).$$

It should be stressed that no such product formula holds in general. Not only does it fail for more general Banach spaces (Example 2.26), but even for Hilbert space one needs at least Shalom's assumption mentioned above. Compare the similar situation for the cohomological product formulas of [Sh] and [BMd].

1.8. When G is locally compact, we can as in the Lie case consider its lattices. One then calls a lattice $\Gamma < G$ *irreducible* if its projections to all G_i are dense. The above Theorem C can be used to establish a superrigidity result for irreducible lattices much in the way of [Sh]. (The general idea to use irreducibility in order to transfer results from $G_1 \times \cdots \times G_n$ to Γ was also illustrated in [BMz], [BMd], [MS]; it seems to originate from the work of Margulis and [BK].)

Theorem D. *Let Γ be an irreducible uniform lattice in a locally compact σ -compact group $G = G_1 \times \cdots \times G_n$. Let B be a superreflexive space with a uniformly equicontinuous affine Γ -action on it. Assume that either:*

- *The group G has compact Abelianization and property (\bar{T}_B) , or:*
- *The associated linear Γ -representation does not almost have invariant vectors.*

Then there is a Γ -closed complemented affine subspace of B on which the Γ -action is a sum of actions extending continuously to G and factoring through $G \rightarrow G_i$.

Remark 1.9. More precisely, the conclusion means that there are superreflexive spaces E_i endowed each with a continuous uniformly equicontinuous affine

G -action factoring through $G \rightarrow G_i$ and a Γ -equivariant affine continuous map $\bigoplus_{i=1}^n E_i \rightarrow B$. Equivalently, the cocycle $b : \Gamma \rightarrow B$ of the original Γ -action is cohomologous to a sum $b_1 + \dots + b_n$ of cocycles b_i ranging in a subspace $B_i \subseteq B$ on which the *linear* Γ -representation extends continuously to a G -representation factoring through G_i and such that b_i extends continuously to a cocycle $G \rightarrow G_i \rightarrow B_i$ (with respect to the corresponding linear G -representation). Moreover, $B_i \cong E_i$ as G -spaces.

If one disregards a component of B where the linear Γ -representation ranges in a compact group of operators, this *sum of actions* is actually just a direct sum $\bigoplus B_i \subseteq B$ (see Remark 8.9).

Remark 1.10. A *uniform lattice* (in a locally compact group) is just a discrete cocompact subgroup; the theorem however also holds for certain non-uniform lattices, see Section 8 where we state a more general form of the result (Theorem 8.3).

Organization of the Paper. In Section 2 we collect preliminary facts and lemmas on uniformly convex/smooth and superreflexive Banach spaces, linear representations and affine isometric on such spaces, special properties of L^p -spaces, and some general remarks and basic counter-examples. In Section 3 Theorem 1.3 is proved. Equivalence of properties (T) and (T_{L^p}) (Theorem A) is proved in Section 4. In Section 5 we discuss higher rank groups and prove Theorem B. Section 6 studies minimal convex sets. Section 7 addresses product groups and proves the splitting theorem (Theorem C); it also proposes a proof of Theorem B that provides some evidence for Conjecture 1.6. In Section 8, we prove Theorem D. Appendix 9 describes Shalom's proof of a generalized Howe-Moore theorem.

Acknowledgments. We would like to thank D. Fisher and G.A. Margulis for their interest in this work and for letting us include their argument for $2 < p < 2 + \varepsilon$ in Section 3.3. We are indebted to A. Naor for several helpful conversations and to A. Nevo for his remarks on the first manuscript. We are grateful to Y. Shalom for letting us give his proof of a ucus version of Howe-Moore (Theorem 9.1).

2. PRELIMINARIES

This section contains basic definitions, background facts and some preliminary lemmas to be used in the proofs of our main results.

2.1. Banach Spaces. Let B be a Banach space. We denote by $S(B) = \{x \in B : \|x\| = 1\}$ its unit sphere. For $x \in B$ and $r > 0$ we denote by $B(x, r)$ and $\bar{B}(x, r)$ the open, respectively closed, ball of radius r around x .

A Banach space B is said to be *strictly convex* if its unit sphere does not contain straight segments, or equivalently if $\|(x + y)/2\| < 1$ whenever $x \neq y \in S(B)$. A Banach space B is called *uniformly convex* if the *convexity modulus* function

$$(2.1) \quad \delta(\varepsilon) = \inf \{1 - \|x + y\|/2 : \|x\|, \|y\| \leq 1, \|x - y\| \geq \varepsilon\}$$

is positive $\delta(\varepsilon) > 0$ whenever $\varepsilon > 0$.

We shall also use the notion of *uniform smoothness* of Banach spaces, which is easiest to define as the uniform convexity of the dual space B^* (see [BL, App. A]).

Hence a Banach space B is *uniformly convex and uniformly smooth* (hereafter abbreviated *ucus*) if both B and its dual B^* are uniformly convex.

Facts 2.1. We refer to [BL] for the following:

- (1) The function $\delta(\varepsilon)$ is non-decreasing and tends to 0 when ε tends to 0. If B is uniformly convex then $\delta(\varepsilon) \rightarrow 0 \Leftrightarrow \varepsilon \rightarrow 0$.
- (2) Uniformly convex Banach spaces are reflexive. Hence the class of *ucus* Banach spaces is closed under taking duals. This class is also closed under the operations of taking closed subspaces and quotients.
- (3) If B^* is strictly convex, in particular if B is uniformly smooth, then every $x \in S(B)$ has a unique supporting functional $x^* \in S(B^*)$, i.e. a unit functional with $\langle x, x^* \rangle = 1$.
- (4) If B is *ucus* then the *duality map* $*$: $S(B) \rightarrow S(B^*)$, $x \mapsto x^*$, is a *uniformly continuous* homeomorphism with a uniformly continuous inverse.
- (5) To any non empty bounded subset $E \subseteq B$ of a reflexive strictly convex Banach space B , one can associate a unique point $C(E) \in B$, the *circumcentre* of E (a.k.a. the Chebyshev centre), defined as the unique $x \in B$ minimizing $\inf\{r > 0 : E \subseteq \overline{B}(x, r)\}$.

The existence of $x = C(E)$ in (5) follows from weak compactness of closed bounded convex sets (i.e. from reflexivity), whilst the uniqueness follows from convexity. Note that somewhat contrary to the intuition, it was shown by V. Klee [Kl] that if $\dim(B) \geq 3$ and B is not a Hilbert space, then there exist a bounded subset $E \subseteq B$ for which $C(E)$ does not belong to the closed convex hull of E . The notion of circumcentre is also used in CAT(0) geometry. For CAT(0) spaces, the circumcentre $C(E)$ always lies in the closed convex hull of E^\ddagger .

The following can be found e.g. in [BL, A.6, A.8]:

Theorem 2.2. *The following conditions on a topological vector space V are equivalent:*

- (1) V is isomorphic to a uniformly convex Banach space.
- (2) V is isomorphic to a uniformly smooth Banach space.
- (3) V is isomorphic to a *ucus* Banach space.

The space V is called *superreflexive* if these equivalent condition hold. The class of superreflexive spaces is closed under taking duals, closed subspaces and quotients of topological vector spaces.

2.2. Linear Representations. Let V be a topological vector space. We denote by $\mathbf{GL}(V)$ the group of invertible linear transformations of V which are continuous together with their inverses.

Following the standard terminology [B1, Def. 2 of §2 n° 1], a group G of transformations of V is *uniformly equicontinuous* (with respect to the uniform structure

[‡]Note that Hilbert spaces are, in a sense, the most convex Banach spaces – they have the largest possible modulus of continuity $\delta(\varepsilon)$ among Banach spaces. On the other hand, Hilbert spaces have the smallest possible modulus of continuity among CAT(0) spaces. Thus, in a sense, CAT(0) spaces are more convex than (non-Hilbertian) Banach spaces

deduced from the topological vector space structure) if for any neighbourhood U of $0 \in V$ there exists a neighbourhood W of 0 such that

$$(2.2) \quad x - y \in W \implies \forall g \in G : g(x) - g(y) \in U.$$

This definition will be applied to both linear groups, or affine groups.

For a topological vector space V , we denote by $N(V)$ the (a priori possibly empty) set of norms on V defining the given topology. Elements of $N(V)$ will be called *compatible* norms and are pairwise equivalent.

The following key proposition is an equivariant version of Theorem 2.2. It enables us to reduce questions about uniformly equicontinuous linear representations on superreflexive spaces to isometric linear representations on ucus Banach spaces.

Proposition 2.3 (Invariant ucus norm). *For a superreflexive topological vector space V and a group of linear transformations G of V , the following conditions are equivalent:*

- (1) G is uniformly equicontinuous group of linear transformations of V .
- (2) G acts by uniformly bounded linear transformations with respect to any/all compatible norm on V .
- (3) G acts by linear isometries with respect to some uniformly convex compatible norm on V .
- (4) G acts by linear isometries with respect to some uniformly smooth compatible norm on V .
- (5) G acts by linear isometries with respect to some uniformly convex and uniformly smooth compatible norm on V .

Proof. The main part of the proof is the implication “[3) and (4)] \Rightarrow (5)”; we begin by establishing this.

Let $N(V)$ denote the set of all compatible norms on V equipped with the metric

$$d(\|\cdot\|_1, \|\cdot\|_2) = \sup_{x \neq 0} \left| \log \frac{\|x\|_1}{\|x\|_2} \right|.$$

This is a complete metric space. Let $N(V)^G$ stand for the closed subspace of G -invariant norms in $N(V)$. Denoting by $\delta_{\|\cdot\|}$ the convexity modulus of $\|\cdot\| \in N(V)^G$, the subset $N_{uc}(V)^G$ of uniformly convex G -invariant norms on V is given by the countable intersection

$$N_{uc}(V)^G = \bigcap_{n=1}^{\infty} O_n, \quad \text{where} \quad O_n = \{ \|\cdot\| \in N(V)^G : \delta_{\|\cdot\|}(1/n) > 0 \}.$$

Observe that the sets O_n are open. If $\|\cdot\|_0$ is some fixed G -invariant compatible uniformly convex norm (given in (3)) then any $\|\cdot\| \in N(V)^G$ can be viewed as a limit of uniformly convex norms $\|\cdot\| + \varepsilon\|\cdot\|_0$ as $\varepsilon \searrow 0$. Hence $N_{uc}(V)^G$ is a dense G_δ set in $N(V)^G$.

By duality between $N_{uc}(V^*)^G$ and the set $N_{us}(V)^G$ of uniformly smooth norms in $N(V)^G$, the latter is also a dense G_δ set in the Baire space $N(V)^G$. In particular $N_{uc}(V)^G \cap N_{us}(V)^G$ is not empty, as claimed.

Now we observe that “(1) \Leftrightarrow (2)” follows from the definitions and that “(5) \Rightarrow [(3) and (4)]” as well as “[(3) or (4) or (5)] \Rightarrow (2)” are trivial. Moreover, proving “(2) \Rightarrow (3)” will also yield “(2) \Rightarrow (4)” by duality, using the fact that the dual to a superreflexive space is superreflexive. Therefore it remains only to justify “(2) \Rightarrow (3)”:

Let $\|\cdot\|$ be a compatible uniformly convex norm on V . The corresponding operator norms $\|g\| = \sup_{x \neq 0} \|gx\|/\|x\|$ are uniformly bounded by some $C < \infty$. Hence

$$\|x\|' = \sup_{g \in G} \|gx\|$$

defines a norm, equivalent to $\|\cdot\|$, and G -invariant. It is also uniformly convex. Indeed, if $\|x\|' = \|y\|' = 1$ and $\|(x+y)/2\|' > 1 - \alpha$ then for some $g \in G$

$$\|(gx + gy)/2\| > 1 - \alpha \quad \text{whilst} \quad \|gx\| \leq \|x\|' = 1, \quad \|gy\| \leq \|y\|' = 1.$$

Thus $\alpha \geq \delta_{\|\cdot\|}(\|gx - gy\|) \geq \delta_{\|\cdot\|}(\|x - y\|'/C)$. Hence the convexity moduli satisfy

$$\delta_{\|\cdot\|'}(\varepsilon) \geq \delta_{\|\cdot\|}(\varepsilon/C) > 0 \quad \text{for all } \varepsilon > 0.$$

□

If G is a topological group, one should impose a continuity assumption on linear G -representations on V , that is on homomorphisms $\varrho : G \rightarrow \mathbf{GL}(V)$. $\mathbf{GL}(V)$ is naturally equipped with the operator norm (which is too strong for representation theory), and with the *weak* and the *strong* operator topologies. For uniformly equicontinuous representations the latter two topologies impose the same continuity assumption:

Lemma 2.4. *Let G be a topological group, V a superreflexive topological vector space, and $\varrho : G \rightarrow \mathbf{GL}(V)$ a homomorphism. Then the following are equivalent.*

- (1) ϱ is weakly continuous.
- (2) ϱ is strongly continuous.
- (3) The orbit maps $g \mapsto \varrho(g)u$ are continuous.
- (4) The action map $G \times B \rightarrow B$ is jointly continuous.

Proof. Clearly it is enough to prove (1) \Rightarrow (4). Let $\|\cdot\|$ be a $\varrho(G)$ -invariant ucus norm on V . Assume $g_n \rightarrow e \in G$ and $u_n \rightarrow u \in S(B)$. Then

$$\begin{aligned} |\langle \varrho(g_n)u_n, u^* \rangle - 1| &\leq |\langle \varrho(g_n)u_n, u^* \rangle - \langle \varrho(g_n)u, u^* \rangle| + |\langle \varrho(g_n)u, u^* \rangle - 1| \\ &\leq \|u_n - u\| + |\langle \varrho(g_n)u, u^* \rangle - 1| \rightarrow 0 \end{aligned}$$

It follows that $\varrho(g_n)u_n \rightarrow u$ because

$$1 \leftarrow \left\langle \frac{\varrho(g_n)u + u}{2}, u^* \right\rangle \leq \left\| \frac{\varrho(g_n)u + u}{2} \right\| \leq 1 - \delta(\|\varrho(g_n)u - u\|).$$

□

2.3. Invariant complements. One of the convenient properties of Hilbert spaces is the existence of a canonical complement M^\perp to any closed subspace M . Recall that a closed subspace X of a Banach space B is called complemented if there is another closed subspace $Y \leq B$ such that $B = X \oplus Y$ algebraically and topologically. This is equivalent to each of the following:

- There is a continuous linear projection from B to X .
- There is a closed subspace Y and a continuous linear projection $p : B \rightarrow Y$ with $\ker(p) = X$.

A classical theorem of Lindenstrauss and Tzafriri says that every infinite dimensional Banach space which is not isomorphic to a Hilbert space, admits a non-complemented closed subspace. However, for any uniformly equicontinuous linear representation ϱ of a group G on a superreflexive space B , the subspace of invariant vectors $B^{\varrho(G)}$ admits a canonical complement, described below.

In view of Proposition 2.3 we may assume that the representation is linear isometric with respect to a ucus norm on B , which allows to use the duality map of the unit spheres $* : S(B) \rightarrow S(B^*)$.

Given any linear representation $\varrho : G \rightarrow \mathbf{GL}(V)$ there is an associated dual (or contragradient) linear G -representation $\varrho^* : G \rightarrow \mathbf{GL}(V^*)$ defined by

$$\langle x, \varrho^*(g)y \rangle = \langle \varrho(g^{-1})x, y \rangle \quad (g \in G, x \in V, y \in V^*).$$

If B is a Banach space and $\varrho : G \rightarrow \mathbf{O}(B)$ is linear isometric representation, then so is its dual $\varrho^* : G \rightarrow \mathbf{O}(B^*)$, where B^* is equipped with the dual norm. Hence the dual to a uniformly equicontinuous representation on a superreflexive space is also of the same type.

Observation 2.5. If B is a ucus Banach space and $\varrho : G \rightarrow \mathbf{O}(B)$, then the duality map $* : S(B) \rightarrow S(B^*)$ between the unit spheres intertwines the actions of $\varrho(G)$ and $\varrho^*(G)$. In particular it maps the set of $\varrho(G)$ -fixed unit vectors to the set of $\varrho^*(G)$ -fixed unit vectors.

Proposition 2.6. *Let ϱ be a uniformly equicontinuous linear representation of G on a superreflexive space B , let $B^{\varrho(G)}$ denote the subspace of $\varrho(G)$ -fixed vectors in B , and let $B' = B'(\varrho)$ be the annihilator of $(B^*)^{\varrho^*(G)}$ in B . Then*

$$B = B^{\varrho(G)} \oplus B'(\varrho).$$

Furthermore, the decomposition is canonical in the following sense: If we denote by $p(\varrho)$ and $p'(\varrho)$ the associated projections, then for every morphism of uniformly equicontinuous linear representations $\phi : (B_1, \varrho_1) \rightarrow (B_2, \varrho_2)$, the following diagrams are commutative:

$$(2.3) \quad \begin{array}{ccc} B_1 & \xrightarrow{\phi} & B_2 \\ p(\varrho_1) \downarrow & & p(\varrho_2) \downarrow \\ B_1 & \xrightarrow{\phi} & B_2 \end{array} \quad \begin{array}{ccc} B_1 & \xrightarrow{\phi} & B_2 \\ p'(\varrho_1) \downarrow & & p'(\varrho_2) \downarrow \\ B_1 & \xrightarrow{\phi} & B_2 \end{array}$$

Remark 2.7. The conclusion fails if we drop the superreflexivity assumption, see Example 2.28.

Proof of the proposition. Choose a G -invariant uniformly convex and uniformly smooth norm on B , and the dual one on B^* (Proposition 2.3). For any unit vector $x \in B^{\varrho(G)}$ and arbitrary $y \in B'$

$$1 = \langle x, x^* \rangle = \langle x - y, x^* \rangle \leq \|x - y\| \cdot \|x^*\| = \|x - y\|.$$

Thus $B^{\varrho(G)} \cap B' = \{0\}$ and $B^{\varrho(G)} \oplus B'$ is a closed subspace in B . It is also dense in B . Indeed if $\lambda \in B^*$ is a unit vector vanishing on B' it can not vanish on $B^{\varrho(G)}$, because $\lambda \in (B^*)^{\varrho^*(G)}$ implies $\lambda^* \in B^{\varrho(G)}$ and $\langle \lambda^*, \lambda \rangle = 1$. Thus $B^{\varrho(G)} \oplus B' = B$.

The last assertion follows from the fact that $\phi(B_1^{\varrho_1}) \subseteq B_2^{\varrho_2}$, and $\phi^*((B_2^*)^{\varrho_2}) \subseteq (B_1^*)^{\varrho_1}$ yields $\phi(B_2') \subseteq B_1'$. \square

Corollary 2.8. *The decomposition $B = B^{\varrho(G)} \oplus B'$ is preserved by the normalizer of $\varrho(G)$ in $\mathbf{GL}(B)$.*

Corollary 2.9. *Let $G = G_1 \times G_2$ be any product of two groups and B a super-reflexive space with a uniformly equicontinuous linear G -representation ϱ . Then there is a canonical G -invariant decomposition*

$$B = B^{\varrho(G)} \oplus B_0 \oplus B_1 \oplus B_2$$

such that $B^{\varrho(G_i)} = B^{\varrho(G)} \oplus B_i$ for $i = 1, 2$.

Proposition 2.10. *Let ϱ be a uniformly equicontinuous linear G -representation on a superreflexive space B . Then*

- (1) $B^{\varrho(G)}$ is isomorphic to B/B' as topological vector spaces.
- (2) B' is isomorphic to $B/B^{\varrho(G)}$ as G -representations.
- (3) $(B^{\varrho(G)})^*$ is isomorphic to $(B^*)^{\varrho^*(G)}$ as topological vector spaces.
- (4) $(B')^*$ is isomorphic to $(B^*)'$ as G -representations.
- (5) B' almost has invariants if and only if $(B^*)'$ almost has invariants.
- (6) If $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is an exact sequence of uniformly equicontinuous linear G -representations on superreflexive spaces, then B' almost has invariant vectors if and only if A' or C' do.

If B is equipped with a compatible uniformly convex and uniformly smooth G -invariant norm, then the natural isomorphisms in (1) and (3) are isometric.

Proof. Equip B with a G -invariant ucus norm (Proposition 2.3).

By the open mapping theorem the maps $p : B \rightarrow B^{\varrho(G)}$ and $p' : B \rightarrow B'$ induce isomorphisms of topological vector spaces

$$(1) \quad \tilde{p} : B/B' \rightarrow B^{\varrho(G)}, \quad (2) \quad \tilde{p}' : B/B^{\varrho(G)} \rightarrow B'.$$

By 2.5 $(B^{\varrho(G)})^*$ is $(B^*)^{\varrho^*(G)}$ and the latter is isomorphic to $B^*/(B^*)'$. This proves (3).

To see that (1) and (3) are isometric (with respect to the norms corresponding to any ucus G -invariant norm on B) we note that the isomorphisms above satisfy $\|\tilde{p}^{-1}\|, \|\tilde{p}'^{-1}\| \leq 1$ by the definition of the norm on a quotient space. Furthermore,

for $v \in S(B^{\varrho(G)})$, we have $v^* \in S((B^*)^{\varrho^*(G)})$, hence

$$\begin{aligned} \|\tilde{p}^{-1}(v)\|_{B/B'} &= \inf \{ \|v + v'\|_B : v' \in B' \} \\ &\geq \inf \{ \langle v + v', v^* \rangle : v' \in B' \} = \langle v, v^* \rangle = 1. \end{aligned}$$

Hence \tilde{p} is an isometry $B/B' \cong B^{\varrho(G)}$. Similarly, $(B^{\varrho(G)})^* = (B^*)^{\varrho^*(G)} \cong B^*/(B^*)'$.

In general Banach spaces the dual E^* of a subspace $E < F$ is isometric to the quotient F^*/E^\perp by the annihilator $E^\perp < F^*$ of E . Thus with respect to a ucus norm on B and the above spaces, $(B')^*$ is isometric to $B^*/(B^*)^{\varrho^*(G)}$ as Banach spaces, while the latter is isomorphic to $(B^*)'$ as a topological vector space by (2). Whence (4).

(5) Assume that there exist $x_n \in S(B')$ with $\text{diam}(\varrho(K) \cdot x_n) \rightarrow 0$. The uniformly continuous map $*$: $S(B) \rightarrow S(B^*)$ takes vectors $x_n \in S(B')$ to vectors $x_n^* \in S(B^*)$ with $\text{diam}(\varrho^*(K) \cdot x_n^*) \rightarrow 0$. Since $\{x_n^*\}$ are uniformly separated from $(B^*)^{\varrho^*(G)}$ their normalized projection y_n^* to $(B^*)'$ still satisfy $\text{diam}(\varrho^*(K) \cdot y_n^*) \rightarrow 0$.

(6) As A' maps into B' , if A' almost has invariants, then so does B' . If C' almost has invariants, then so does $(C^*)'$, hence $(B^*)'$, hence B' . On the other hand, assume B' almost has invariant unit vectors b_n . Assume for simplicity that $A = A'$, $B = B'$ and $C = C'$. Note that C is isomorphic to B/A , and denote by $\pi : B \rightarrow C$ the projection. Then either $\pi(b_n)$ converges to $0 \in C$, then there exist a_n such that $b_n - a_n$ converges to $0 \in B$, and the normalized sequence $(\frac{a_n}{\|a_n\|})$ is almost invariant in A , or there exist a subsequence b_{n_k} with $\inf_k \|\pi(b_{n_k})\| > 0$, and then the normalized sequence $(\frac{\pi(b_{n_k})}{\|\pi(b_{n_k})\|})$ is almost invariant in C . \square

Remark 2.11. For ucus Banach space B , Definition 1.1 of property (T_B) can be rephrased as follows: For any representation $\varrho : G \rightarrow \mathbf{O}(B)$, the restriction $\varrho' : G \rightarrow \mathbf{O}(B')$ of ϱ to the invariant subspace B' complement to $B^{\varrho(G)}$ does not almost have invariant vectors, i.e. for some compact $K \subseteq G$ and $\varepsilon > 0$

$$\forall v \in S(B') \quad \exists g \in K \text{ s.t. } \|\varrho(g)v - v\| \geq \varepsilon.$$

Hence item (4) gives:

Corollary 2.12. *Let B be a ucus Banach space, and G be a locally compact group. Then G has property (T_B) iff it has (T_{B^*}) .*

2.4. Affine Actions. The affine group $\text{Aff}(V)$ of a real affine space V (a vector space who forgot its origin) consists of invertible maps satisfying:

$$T(t \cdot x + (1-t) \cdot y) = t \cdot T(x) + (1-t) \cdot T(y), \quad (t \in \mathbf{R}, x, y \in V)$$

The group $\text{Aff}(V)$ is a semi-direct product $\text{Aff}(V) = \mathbf{GL}(V) \ltimes V$, i.e. an invertible affine map T has the form $T(x) = Lx + b$ where $L \in \mathbf{GL}(V)$ is linear invertible.

An affine action of a group G on V , i.e. a homomorphism $G \rightarrow \text{Aff}(V)$, has the form

$$g \cdot x = \varrho(g)x + c(g),$$

where $\varrho : G \rightarrow \mathbf{GL}(V)$ is a linear G -representation (we call it the *linear part* of the action) and $c : G \rightarrow V$ is a ϱ -cocycle, namely an element of the Abelian group

$$(2.4) \quad Z^1(\varrho) = \{c : G \rightarrow V : c(gh) = \varrho(g)c(h) + c(g), \forall g, h \in G\}.$$

The group $Z^1(\varrho)$ of ϱ -cocycles contains the subgroup of ϱ -coboundaries

$$(2.5) \quad B^1(\varrho) = \{c(g) = v - \varrho(g)v : v \in V\}.$$

$Z^1(\varrho)$ describes all affine G -actions on V with linear part ϱ , and $B^1(\varrho)$ corresponds to those affine actions which have a G -fixed point (namely v in (2.5)). This description involves the choice of reference point – the origin – in the space. Two cocycles differing by a coboundary can be thought of defining the same affine action viewed from different reference points. The first cohomology of G with ϱ -coefficients is the Abelian group

$$H^1(G, \varrho) = Z^1(\varrho)/B^1(\varrho).$$

It describes different types of actions in the above sense. $H^1(G, \varrho) = 0$ iff any affine G -action on V with linear part ϱ has a fixed point.

For a Banach space B denote by $\text{Isom}(B)$ the group of isometries of B as a metric space. It is a classical theorem of Mazur-Ulam that any surjective isometry T of a Banach space B is necessarily affine $T(x) = Lx + c$ with linear part $L \in \mathbf{O}(B)$ being isometric. (This theorem is actually easy when B is strictly convex; see Lemma 6.1). Hence $\text{Isom}(B) = \mathbf{O}(B) \ltimes B$.

Now suppose that V is a superreflexive topological vector space. We shall say that a group G of affine self maps is *uniformly equicontinuous* if it satisfies (2.2). This condition is equivalent to uniform equicontinuity of the linear part $\varrho : G \rightarrow \mathbf{GL}(V)$.

Proposition 2.13. *For a superreflexive topological vector space V and a group of transformations G of V the following conditions are equivalent:*

- (1) G is uniformly equicontinuous group of affine transformations of V .
- (2) G acts by uniformly Lipschitz affine transformations with respect to any/all compatible norms on V .
- (3) G acts by isometries with respect to some compatible norm on V .
- (4) G acts by affine isometries with respect to some uniformly convex and uniformly smooth compatible norm on V .

Proof. Apply Proposition 2.3 to the linear part of the affine action, using Ulam-Mazur to deduce in (3) that the action is affine. \square

If G is a topological group acting by affine transformations on a topological vector space V , continuity of the action

$$G \times V \rightarrow V, \quad g \cdot x = \varrho(g)x + c(g)$$

is equivalent to continuity of the linear part $G \times V \rightarrow V$ and the continuity of the cocycle $c : G \rightarrow V$. Indeed $c(g) = g \cdot 0$, and $\varrho(g)x = g \cdot x - c(g)$.

Hence in the context of topological groups, affine actions should be assumed continuous, and $Z^1(G, \varrho)$ will include only continuous cocycles $c : G \rightarrow V$ (we

assume that the linear part ϱ is continuous as well). If G is a locally compact σ -compact group, then $Z^1(\varrho)$ has a natural structure of a Fréchet space with respect to the family of semi-norms

$$\|c\|_K = \sup_{g \in K} \|c(g)\|$$

where $K \subseteq G$ runs over a countable family of compact subsets which cover G . Moreover, if G is compactly generated (e.g. if G has property (T)) say by K_0 , then $\|c\|_{K_0}$ is a norm on $Z^1(\varrho)$ (note that any cocycle $c \in Z^1(\varrho)$ is completely determined by its values on a generating set), and $Z^1(\varrho)$ is a Banach space with respect to this norm. We remark that in general $B^1(\varrho)$ is not closed in $Z^1(\varrho)$ (this is the idea behind the $(F_B) \Rightarrow (T_B)$ argument of Guichardet – see Section 3).

Lemma 2.14. *For a uniform equicontinuous affine action of a group G on a superreflexive space B , the following are equivalent:*

- (1) *There exists a bounded G -orbit.*
- (2) *All G -orbits are bounded*
- (3) *G fixes a point in B .*
- (4) *G preserves a (Borel regular) probability measure on B .*

Note that the notion of a subset $E \subseteq V$ being bounded, means that for any open neighbourhood U of $0 \in V$ there is some $t \in \mathbf{R}$ so that $E \subseteq t \cdot U$. This notion agrees with the notion of being bounded with respect to any compatible norm on V .

Proof. Introduce a G -invariant uniformly convex norm on V (Proposition 2.13). The only non-trivial implications are $(4) \Rightarrow (1) \Rightarrow (3)$. For the first, let μ be a G -invariant probability on B . Since B is a countable union of closed bounded sets, there is a closed bounded set $A \subseteq B$ with $\mu(A) > 1/2$. For all $g \in G$ we have $\mu(gA) > 1/2$ hence $gA \cap A \neq \emptyset$. It follows that the G -orbit of every point of A is bounded.

The latter implication follows by considering the circumcentre (compare Section 2.1) of the given bounded G -orbit. \square

Proposition 2.15. *Let B be a ucus Banach space. Then*

- (1) *Any finite (or compact) group has properties (T_B) and (F_B) .*
- (2) *Properties (T_B) and (F_B) pass to quotient groups.*
- (3) *If $G = G_1 \times \cdots \times G_n$ is a finite product of topological groups then G has property (T_B) (resp. (F_B)) iff all G_i have this property.*

Proof. (1) and (2) are straightforward, (3) follows from Corollary 2.9. \square

2.5. Special Properties of $L^p(\mu)$ -Spaces. In this section we collect some special properties of the Banach spaces $L^p(\mu)$ which will be used in the proofs.

By an $L^p(\mu)$, or $L^p(X, \mu)$ space we mean the usual space of equivalence classes (modulo null sets) of measurable p -integrable functions $f : X \rightarrow \mathbf{R}$, where μ is a positive σ -finite measure defined on a standard Borel space (X, \mathcal{B}) . If $1 < p < \infty$ then $L^p(\mu)$ is ucus, whilst $L^1(\mu)$ and $L^\infty(\mu)$ are not (they are not even strictly

convex). For $1 \leq p < \infty$ the dual to $L^p(\mu)$ is $L^q(\mu)$ where $1 < q \leq \infty$ is determined by $q = p/(p-1)$.

The space $L^p([0, 1], \text{Lebesgue})$ is usually denoted by L^p . Any $L^p(\mu)$ -space with *non-atomic* finite or σ -finite measure μ is isometrically isomorphic to L^p . Indeed let $\varphi \in L^1(\mu)$ be a strictly positive measurable function with integral one and let μ_1 be given by $d\mu_1 = \varphi d\mu$. Then

$$f \in L^p(\mu) \mapsto f \cdot \varphi^{-1/p} \in L^p(\mu_1)$$

is a surjective isometry. Since any non-atomic standard probability spaces is isomorphic to $[0, 1]$ as a measure space, $L^p(\mu_1) \cong L^p$. If μ is purely atomic then a similar argument gives an isomorphism of $L^p(\mu)$ with finite or infinite dimensional ℓ^p space. A general $L^p(\mu)$ space is therefore isometrically isomorphic to a direct sum of L^p and ℓ^p components.

More generally, for another Banach space B , one defines the spaces $L^p(\mu, B)$ of B -valued function classes by means of the Bochner integral. We refer the reader to [DU] for details; we recall here that the dual of $L^p(\mu, B)$ is $L^q(\mu, B^*)$ through the natural pairing for all $1 \leq p < \infty$, but only when B has the Radon-Nikodým property – this includes all ucus spaces (see again [DU]). These spaces will be used in Section 8.2 in order to *induce* isometric (linear or affine) actions.

Banach [Ba] and Lamperti [Li] (see also [FJ, Theorem 3.25]) classified the linear isometries of $L^p(\mu)$ as follows.

Theorem 2.16 (Banach, Lamperti). *For any $1 < p < \infty$ where $p \neq 2$, any linear isometry U of $L^p(X, \mathcal{B}, \mu)$ has the form*

$$Uf(x) = f(T(x))h(x) \left(\frac{dT_*\mu}{d\mu}(x) \right)^{\frac{1}{p}}$$

where T is a measurable, measure class preserving map of (X, μ) , and h is a measurable function with $|h(x)| = 1$ a.e.

Let $\mu = \mu_a + \mu_c$ be the decomposition of μ into its atomic and continuous parts ($\mu_a = \mu|_A$ where $A \subseteq X$ is the (at most countable) set of atoms of μ). Then

$$L^p(\mu) = L^p(\mu_c) \oplus L^p(\mu_a) \cong L^p \oplus \ell^p(A) \quad \text{or just} \quad \ell^p(A),$$

the latter case occurs if $\mu = \mu_a$ is a purely atomic measure. Note that it follows from Banach-Lamperti theorem that this decomposition is preserved by any linear isometry of $L^p(\mu)$. As $\ell^p(A)$ has a much smaller group of linear (or affine) isometries than L^p we could restrict our attention only to the latter. However we shall not make use of this “simplification”.

Another useful tool in the study of L^p -spaces is the Mazur map.

Theorem 2.17 ([BL, Theorem 9.1]). *Let μ be a σ -finite measure. For any $1 \leq p, q < \infty$ the Mazur map $M_{p,q} : L^p(\mu) \rightarrow L^q(\mu)$ defined by*

$$M_{p,q}(f) = \text{sign}(f) \cdot |f|^{\frac{p}{q}}$$

is a (non-linear) map which induces a uniformly continuous homeomorphism between the unit spheres $M_{p,q} : S(L^p(\mu)) \rightarrow S(L^q(\mu))$.

(Note that if $p, q \neq 1$ and $p^{-1} + q^{-1} = 1$ then the restriction of $M_{p,q}$ to the unit spheres is just the duality map $*$: $S(L^p(\mu)) \rightarrow S(L^p(\mu)^*)$).

In the proofs of Theorems A and B, the results for subspaces and quotients are deduced from the $L^p(\mu)$ case using the following theorem of Hardin about extension of isometries defined on subspaces of $L^p(\mu)$. The formulation we give here is not quite identical to the original, but it easily follows from it and from its proof (see [Ha, Theorem 4.2] or [FJ, Theorem 3.3.14]).

Theorem 2.18 (Hardin). *Let (X, \mathcal{B}, μ) be a measure space. For every closed subspace $F \subseteq L^p(X, \mu)$, there is a canonical extension $F \subseteq \tilde{F} \subseteq L^p(\mu)$ which is isometric to $L^p(X', \mu')$ for some other measure space (X', μ') . Furthermore, if $1 < p \notin 2\mathbf{Z}$, then every linear isometry $U : F \rightarrow L^p(Y, \nu)$ extends uniquely to a surjective linear isometry $\tilde{U} : \tilde{F} \rightarrow \tilde{U}\tilde{F} \subseteq L^p(Y, \nu)$.*

Remark 2.19. If $\mathcal{B}' \leq \mathcal{B}$ is the minimal sub σ -algebra with respect to which all the functions in F are measurable, then $\tilde{F} = L^p(X, \mathcal{B}', \mu)$.

A straightforward consequence is the following:

Corollary 2.20. *Let $1 < p \notin 2\mathbf{Z}$, and let $F \subseteq L^p(X, \mu)$ be a closed subspace. Let ρ be a linear isometric representation of the group G on F . Then there is some linear isometric G -representation ρ' of G on some other space $L^p(X', \mu')$, and a linear G -equivariant isometric embedding $F \hookrightarrow L^p(X', \mu')$.*

Another important fact about $L^p(\mu)$ -spaces, this time for $p \in (0, 2]$, is that $B = L^p(\mu)$ has an embedding $j : B \rightarrow \mathcal{H}$ into the unit sphere of a Hilbert space so that $\langle j(x), j(y) \rangle = \|x - y\|^p$. Having such an embedding is equivalent (via the classical result of I.J. Schoenberg, see [BHV]) to the following:

Proposition 2.21. *For $0 < p < 2$ and any $s > 0$ the function $f \mapsto e^{-s\|f\|^p}$ is positive definite on $L^p(\mu)$, i.e. for any finite collection $f_i \in L^p(\mu)$ and any $\lambda_i \in \mathbf{C}$*

$$\sum_{i,j} e^{-s\|f_i - f_j\|^p} \lambda_i \overline{\lambda_j} \geq 0.$$

In fact, more is known: It was shown by Bretagnolle, Dacunha-Castelle and Krivine [BDCK] (c.f. [WW, 5.1]) that, for $1 \leq p \leq 2$, a Banach space X is isometric to a closed subspace of $L^p(\mu)$ iff $e^{-s\|\cdot\|^p}$ is a positive definite function on X for any $s > 0$.

2.6. Some Easy Counterexamples and Remarks.

Example 2.22 ($(T_B) \not\Rightarrow (F_B)$). Let B be a Banach space with $\mathbf{O}(B) \cong \mathbf{Z}/2\mathbf{Z}$, i.e. a space where the only isometries are the identity and the antipodal map $x \mapsto -x$. A trivial example of such a space is the line $B = \mathbf{R}$, but there are such spaces of arbitrary dimensions even within the class of ucus Banach spaces. Clearly for such a space any group has property (T_B) . However the groups \mathbf{Z} or \mathbf{R} or any group G with sufficiently large Abelianization $G/[G, G]$ would fail to have property (F_B) for it would admit an isometric action by translations: $n \cdot x := x + nx_0$ where $0 \neq x_0 \in B$ is arbitrary. However groups with trivial Abelianization would also have property (F_B) on such an asymmetric Banach space B .

Example 2.23 ($(T) \not\Rightarrow (F_B)$). Suppose G is locally compact non-compact (e.g. $G = \mathbf{SL}_3(\mathbf{R})$ or $G = \mathbf{SL}_3(\mathbf{Z})$). Fix a Haar measure on G and let

$$B = L_0^1(G) = \left\{ f \in L^1(G) : \int f dg = 0 \right\}$$

be the subspace of functions with 0 mean. Then B is isometric to the affine subspace $\{f \in L^1(G) : \int f = 1\}$ on which G acts isometrically by translations without fixed points. Hence G does not have property (F_B) . This Banach space is not ucus.

Example 2.24 ($(T) \not\Rightarrow (T_B)$). Let G be as in Example 2.23. Consider the space $B = C_0(G)$ of continuous real valued functions on G which tend to 0 at ∞ with the sup (L^∞ -) norm. The action of G on B by translations is a linear isometric action. A function $f \in B$ which decays very slowly forms an “almost invariant vector”. On the other hand there are no non-zero invariant vectors. Hence G does not have property (T_B) .

Remark 2.25. Since any separable Banach space is a quotient of ℓ^1 , Example 2.24 shows that case (iii) of Theorem A cannot be extended to $p = 1$.

Example 2.26 (Remarks 1.8(2)). Let $G = G_1 \times G_2$ be any product of non-compact locally compact groups (e.g. $G = \mathbf{Z} \times \mathbf{Z}$). Let $B = L_0^1(G)$ as in Example 2.23. Then $H^1(G, B) \neq 0$, but there are no non-zero G_i -fixed vectors in the associated linear representation. Thus the product formula of Remarks 1.8(2) cannot hold for B .

Let us make some remarks about Kazhdan’s property (T) and property (T_B) as in 1.1 and 2.11. Given a unitary representation (ϱ, \mathcal{H}) of a locally compact group G , a compact subset $K \subseteq G$ and $\varepsilon > 0$, one says that a vector $0 \neq v \in \mathcal{H}$ is (K, ε) -almost invariant if

$$\sup_{g \in K} \|\varrho(g)v - v\| < \varepsilon \cdot \|v\|.$$

A locally compact group G has Kazhdan’s property (T) if and only if it satisfies the following equivalent conditions:

- (1) For any unitary G -representation (ϱ, \mathcal{H}) there exists a compact $K \subseteq G$ and an $\varepsilon > 0$ so that the G -representation ϱ' on $(\mathcal{H}^{\varrho(G)})^\perp \cong \mathcal{H}/\mathcal{H}^{\varrho(G)}$ has no (K, ε) -almost invariant vectors.
- (2) There exist a compact $K \subseteq G$ and an $\varepsilon > 0$ so that all non-trivial irreducible unitary G -representations (ϱ, \mathcal{H}) have no (K, ε) -almost invariant vectors.
- (3) There exist a compact $K \subseteq G$ and an $\varepsilon > 0$ so that for all unitary G -representations (ϱ, \mathcal{H}) the G -representation ϱ' on $(\mathcal{H}^{\varrho(G)})^\perp \cong \mathcal{H}/\mathcal{H}^{\varrho(G)}$ has no (K, ε) -almost invariant vectors.

In the above, (3) clearly implies both (1) and (2). In showing $(1) \Rightarrow (3)$ one uses the fact that the category of Hilbert spaces and unitary representations is closed under ℓ^2 sums and L^2 -integration. The fact that any unitary representation decomposes as an L^2 -integral of irreducible ones gives $(2) \Rightarrow (3)$.

Remark 2.27. Definition 1.1 (Remark 2.11) of property (T_B) is modeled on (1) above. There does not seem to be any reasonable theory of irreducible representations (and decomposition into irreducibles) for Banach spaces other than Hilbert ones. Hence form (2) of property (T) does not seem to have a Banach space generalization. As for (3), for any given $1 < p < \infty$ the class of $L^p(\mu)$ -spaces is closed under taking ℓ^p -sums (and L^p -integrals) and hence for groups with property (T_{L^p}) an analogue of (3) holds, namely there exist $K \subseteq G$ and $\varepsilon > 0$ which are good for all $\varrho : G \rightarrow \mathbf{O}(L^p)$. Also, if a group G has property (T_B) for *all* ucus Banach spaces B (conjecturally all higher rank groups and their lattices) then for every ucus Banach space B there is (K, ε) which is good for all linear isometric representations $G \rightarrow \mathbf{O}(B)$. This uses the fact that $L^2(\mu, B)$ is a ucus if B is (see Lemma 8.5 below).

Finally, we justify Remark 2.7:

Example 2.28. Let G be a discrete group and consider the Banach space $B = \ell^\infty(G)$ with the (linear isometric) regular G -representation ϱ . Then one shows that the space $B^{\varrho(G)}$ (which consists of the constant functions) admits a G -invariant complement (if and) *only if G is amenable*. Indeed, the Riesz space (or Banach lattice) structure of B allows to take the “absolute value” of any linear functional on B ; renormalizing the absolute value of any non-zero invariant functional would yield an invariant mean on G . Alternatively, one can argue similarly on the Banach space of continuous function on any compact topological G -space.

We point out that nevertheless the space B' is well-defined for any topological vector G -space B ; in the case at hand, we have $B' = B$ which shows why it cannot be a complement for $B^{\varrho(G)} \neq 0$.

3. PROOF OF THEOREM 1.3

3.1. Guichardet: $(F_B) \implies (T_B)$.

Proof. Assume G does not have (T_E) , where E is a Banach space, and let $\varrho : G \rightarrow \mathbf{O}(E)$ be a representation such that $E/E^{\varrho(G)}$ admits almost invariant vectors. In order to show that $H^1(G, \varrho) \neq \{0\}$ it suffices to prove that $B^1(G, \varrho) \subseteq Z^1(G, \varrho)$ is not closed.

As was mentioned in Section 2 the space of ϱ -cocycles $Z^1(G, \varrho)$ is always a Fréchet space (and even a Banach space if G is compactly generated). Note that $B^1(G, \varrho)$ is the image of the bounded linear map

$$\tau : E \rightarrow Z^1(G, \varrho), \quad (\tau(v))(g) = v - \varrho'(g)v.$$

If $\tau(E)$ were closed, and hence a Fréchet space, the open mapping theorem would imply that $\tau^{-1} : B^1(G, \varrho) \rightarrow E/E^{\varrho(G)}$ is a bounded map. That would mean that for some $M < \infty$ and a compact $K \subseteq G$

$$\|v\| \leq M \cdot \|\tau(v)\|_K = M \cdot \sup_{g \in K} \|\varrho(g)v - v\|, \quad v \in E/E^{\varrho(G)}$$

contrary to the assumption that ϱ almost contains invariant vectors. \square

3.2. $(T) \implies (F_{L^p}), 0 < p \leq 2$.

Proof. Let G be a locally compact group with Kazhdan's property (T) acting by affine isometries on a closed subspace $B \subseteq L^p(\mu)$ with $0 < p \leq 2$. Using Proposition 2.21 and a slight modification of a Delorme-Guichardet argument for $(T) \implies (FH)$ we shall prove that such an action has bounded orbits. For $1 < p \leq 2$ uniform convexity of $B \subseteq L^p(\mu)$ yields a G fixed point using Lemma 2.14.

Proposition 2.21 allows to define a family, indexed by $s > 0$, of Hilbert space \mathcal{H}_s , embeddings $U_s : B \rightarrow \mathcal{S}(\mathcal{H}_s)$ and unitary representations $\pi_s : G \rightarrow \mathbf{O}(\mathcal{H}_s)$ with the following properties:

- (1) The image $U_s(B)$ spans a dense subspace of \mathcal{H}_s ;
- (2) $\langle U_s(x), U_s(y) \rangle = e^{-s \cdot \|x-y\|^p}$ for all $x, y \in B$;
- (3) $U_s(gx) = \pi_s(g)U_s(x)$ for all $x \in B, g \in G$.

Indeed, one constructs \mathcal{H}_s as the completion of the pre-Hilbert space whose vectors are finite linear combinations $\sum a_i x_i$ of points $x_i \in B$, and the inner product is given by

$$\langle \sum a_i x_i, \sum b_j y_j \rangle = \sum_{i,j} a_i \bar{b}_j e^{-s \cdot \|x_i - y_j\|^p}.$$

The representation π_s can be constructed (and is uniquely determined) by property (3).

Since G is assumed to have Kazhdan's property (T), for some compact subset $K \subseteq G$ and $\varepsilon > 0$, any unitary G -representation with (K, ε) -almost invariant vectors has a non-trivial invariant vector.

Let $x_0 \in B$ be fixed. The isometric G -action is continuous, so Kx_0 is a compact and hence bounded subset of B , hence:

$$R_0 = \sup_{g \in K} \|gx_0 - x_0\| < \infty.$$

For the unit vectors $u_s = U_s(x_0) \in \mathcal{S}(\mathcal{H}_s)$ we have

$$\min_{g \in K} |\langle \pi_s(g)u_s, u_s \rangle| \geq e^{-sR_0^p} \rightarrow 1 \quad \text{as } s \rightarrow 0.$$

In particular for a sufficiently small $s > 0$, $\max_{g \in K} \|\pi_s(g)u_s - u_s\| < \varepsilon$. Let us fix such an s , and rely on property (T) to deduce that π_s has an invariant vector $v \in \mathcal{S}(\mathcal{H}_s)$.

We claim that G must have bounded orbits for its affine isometric action on B . Indeed, otherwise there would exist a sequence $g_n \in G$ so that

$$\|g_n x - y\| \rightarrow \infty \quad \text{and hence} \quad \langle \pi_s(g_n)U_s(x), U_s(y) \rangle \rightarrow 0$$

for all $x, y \in B$. This implies that $\langle \pi_s(g_n)w, u \rangle \rightarrow 0$ for any $w, u \in \text{span}(U_s(B))$, and since $\text{span}(U_s(B))$ is dense in \mathcal{H}_s , for any $w, u \in \mathcal{H}_s$. Taking $w = u = v$, we get a contradiction. Therefore the isometric G -action on B has bounded orbits, and hence fixes a point in case of $1 < p \leq 2$. \square

3.3. Fisher-Margulis: $(T) \implies (F_{L^p})$, $p < 2 + \varepsilon(G)$. Let G have Kazhdan's property (T) . Fix a compact a compact generating subset K of G .

Lemma 3.1. *There exists a constant $C < \infty$ and $\varepsilon > 0$ such that for any G -action by isometries on a closed subspace $B \subseteq L^p(\mu)$ with $p \in (2 - \varepsilon, 2 + \varepsilon)$ and any $x \in B$ there exists a point $y \in B$ with*

$$\|x - y\| \leq C \cdot \text{diam}(K \cdot x), \quad \text{diam}(K \cdot y) < \frac{\text{diam}(K \cdot x)}{2}.$$

Proof. By contradiction there exists a sequence of subspaces $B_n \subseteq L^{p_n}$ with $p_n \rightarrow 2$, isometric G -actions on B_n and points $x_n \in B_n$ so that, after a rescaling to achieve $\text{diam}(K \cdot x_n) = 1$, we have

$$(3.1) \quad \text{diam}(K \cdot y) \geq \frac{1}{2} \quad \forall y \in B(x_n, n).$$

Passing to an ultralimit of the spaces B_n with the marked points x_n and the corresponding G -actions, one obtains an isometric G -action on a Hilbert space \mathcal{H} , because the limit of L^p -parallelogram as $p \rightarrow 2$ is the parallelogram identity, which characterizes Hilbert spaces. If G is a topological group, one needs to "smoothen" the construction to ensure continuity of the limit action (see Shalom [Sh] for details). Due to (3.1) this G -action has no fixed points, contradicting property (FH) and hence (T) of G . \square

Proof of (F_B) for $B \subseteq L^p(\mu)$, $2 \leq p < 2 + \varepsilon(G)$. Now consider an arbitrary isometric G -action on a closed subspace $B \subseteq L^p$ with $|p - 2| < \varepsilon$ where $\varepsilon = \varepsilon(G) > 0$ is as in the lemma. Define a sequence $x_n \in B$ inductively, starting from an arbitrary x_0 . Given x_n , let $R_n = \text{diam}(K \cdot x_n)$. Then applying the lemma there exists x_{n+1} within the ball $B(x_n, C \cdot R_n)$ so that

$$\text{diam}(K \cdot x_{n+1}) < R_n/2.$$

We get $R_n < R_0/2^n$ and $\sum \|x_{n+1} - x_n\| < \infty$. The limit of the Cauchy sequence $\{x_n\}$ is a G -fixed point. \square

Question 3.2. For a given group with property (T) , e.g. for $G = \text{Sp}(n, 1)$, what is $\inf p$ over all p for which G does not have property (F_B) for some closed subspace $B \subseteq L^p$?

4. PROOF OF THEOREM A

We start with the first assertion of the theorem: $(T) \implies (T_B)$ for B being an L^p -related space as in (i), (ii) or (iii) in the theorem. We first reduce to the case (i) where $B = L^p(\mu)$ with $1 \leq p < \infty$. Then using Corollary 2.20 of Hardin's extension theorem, $(T_{L^p(\mu)})$ implies (T_B) for subspaces $B \subseteq L^p(\mu)$ where $p \neq 4, 6, \dots$ as in (ii), and the duality argument (Corollary 2.12) gives the result for quotients of $L^q(\mu)$ with $q \neq 4/3, 6/5, \dots$ as in (iii). Hence it suffices to prove $(T) \implies (T_{L^p(\mu)})$ for $1 \leq p < \infty$. We give two proofs for this implication.

Let us note that our restriction on p and q when taking subspaces/quotients comes from our use of Hardin's theorem.

Question 4.1. Does property (T) implies property (T_B) for and closed subspace and any quotient B of $L^p(\mu)$ for any $1 < p < \infty$?

4.1. Property (T) Implies $(T_{L^p(\mu)})$ – First Proof.

Proof. Assuming that a locally compact group G fails to have property $(T_{L^p(\mu)})$ for some $1 \leq p < \infty$, we are going to show that G does not have (T). We may and will assume $p \neq 2$. Using Remark 2.11 there is a representation $\varrho : G \rightarrow \mathbf{O}(B)$ so that for the canonical complement B' of $B^{\varrho(G)}$ the restriction $\varrho' : G \rightarrow \mathbf{O}(B')$ almost has invariant vectors, i.e. there exist unit vectors $v_n \in S(B')$ so that

$$f_n(g) = \|\varrho(g)v_n - v_n\|$$

converges to 0 uniformly on compact subsets of G .

We shall construct a related unitary, or orthogonal, representation $\pi : G \rightarrow \mathbf{O}(\mathcal{H})$ where $\mathcal{H} = L^2(\mu)$, using the following:

Lemma 4.2. *For $p \neq 2$, the conjugation $U \mapsto M_{p,2} \circ U \circ M_{2,p}$ by the non-linear Mazur map sends $\mathbf{O}(L^p(\mu))$ to $\mathbf{O}(L^2(\mu))$.*

Proof. Follows from Banach-Lamperti description of $\mathbf{O}(L^p(\mu))$ (Theorem 2.16) by calculation. \square

Let us then define $\pi : G \rightarrow \mathbf{O}(\mathcal{H})$ by $\pi(g) = M_{p,2} \circ \varrho(g) \circ M_{2,p}$. Note that $M_{p,2}$ maps $B^{\varrho(G)}$ onto $\mathcal{H}^{\pi(G)}$.

As $S(B')$ is uniformly separated (in fact is at distance 1) from $B^{\varrho(G)}$, the uniform continuity of the Mazur map (Theorem 2.17) implies that $u_n = M_{p,2}(v_n)$ is a sequence in $S(\mathcal{H})$ such that $\text{dist}(u_n, \mathcal{H}^{\pi(G)}) \geq \delta > 0$ and $\varphi_n(g) = \|\pi(g)u_n - u_n\| \rightarrow 0$ uniformly on compact subsets of G . Let w_n denote the projections of u_n to $\mathcal{H} = (\mathcal{H}^{\pi(G)})^\perp$. Then

$$\|w_n\| \geq \delta > 0 \quad \text{and} \quad \|\pi(g)w_n - w_n\| \leq \varphi_n(g) \rightarrow 0$$

uniformly on compacta. Thus the restriction π' of π to \mathcal{H}' does not have G -invariant vectors, but almost has ones. Hence G does not have Kazhdan's property (T). \square

4.2. Property (T) Implies $(T_{L^p(\mu)})$ – Second Proof.

Proof. For $1 < p \leq 2$ we have $(T) \Rightarrow (F_{L^p(\mu)}) \Rightarrow (T_{L^p(\mu)})$ by Theorem 1.3 (1) and (2). Using duality (Corollary 2.12) this implication extends to $L^p(\mu)$ with $2 < p < \infty$. \square

4.3. Property (T_{L^p}) Implies (T).

Proof. Assume that G is not Kazhdan, i.e. G admits a unitary representation without non-trivial invariant vectors, but almost containing (non-trivial) invariant ones. Connes and Weiss [CW] showed how to find such a representation of the form $L_0^2(\mu)$. More precisely, given a unitary G -representation π with almost invariant unit vectors but no invariant ones, they construct a measure-preserving,

ergodic G -action on a probability space (X, μ) which admits a sequence $\{E_n\}$ of asymptotically invariant measurable subsets, namely for every $g \in G$,

$$(4.1) \quad \forall g \in G \quad \mu(gE_n \Delta E_n) \rightarrow 0 \quad \text{whilst} \quad \mu(E_n) = 1/2.$$

Then the unitary G -representation π' on $L_0^2(\mu)$ – the space of zero mean square integrable functions, which is the orthogonal complement of the constants, does not have non-trivial invariant vectors (due to ergodicity) but almost has ones (namely normalized $v_n = \mathbf{1}_{E_n} - 1/2$).

For a given $1 \leq p < \infty$, consider the linear isometric G -representation ϱ on $B = L^p(\mu)$, $\varrho(g)f(x) = f(g^{-1}x)$. Then $B^{\varrho(G)} = \mathbf{R}\mathbf{1}$ – the constants, and its canonical complement is

$$B' = L_0^p(\mu) = \left\{ f \in L^p(\mu) : \int f d\mu = 0 \right\}.$$

The above sequence $\{f_n\}$ lies in $L_0^p(\mu)$ and still satisfies $\|\varrho(g)f_n - f_n\|_p \rightarrow 0$ whilst $\|f_n\|_p = (1/2)^{\frac{1}{p}}$. Hence failing to have Kazhdan's property (T) a group G does not have $(T_{L^p(\mu)})$ either.

In the original paper [CW], Connes and Weiss considered discrete groups. In a similar context the case of locally compact groups was also considered by Glasner and Weiss (see [GW, Section 3] and references therein). One way to treat the non-discrete case, is the following: start from a unitary representation π of a given lcsc G which has almost invariant vectors but no invariant ones, and apply the original Connes-Weiss Gaussian construction to the restriction $\pi|_\Gamma$ of π to some dense countable subgroup $\Gamma \subseteq G$. This gives an ergodic measure-preserving Γ -action on a probability space (X, μ) with an asymptotically invariant sequence $\{E_n\}$ on X . The fact that the representation $\pi|_\Gamma$ came from G is manifested by the fact that it is continuous in the topology on Γ induced from G . It can be shown to imply that the Γ -representation on $L_0^2(X, \mu)$ is also continuous, hence extends to G , and thus the Γ -action on (X, μ) extends to a measurable G -action. This construction gives a uniform convergence in (4.1) on compact subsets of G . \square

5. FIXED POINT PROPERTY FOR HIGHER RANK GROUPS

5.1. The objective of this section is to prove Theorem B; we start with some preliminaries for the *linear* part.

The first ingredient needed for the proof is an analogue of Howe-Moore's theorem on vanishing of matrix coefficients, or rather its corollary analogous to Moore's ergodicity theorem, extended to the framework of uniformly equicontinuous representations on superreflexive Banach spaces. The ucus Banach space version of Howe-Moore is due to Yehuda Shalom (unpublished). With his kind permission we have included the argument in Appendix 9. Here we shall use the following corollary, which we formulate for the case of simple groups.

Corollary 5.1 (Banach space analogue of Moore's theorem). *Let k be a local field and let $G = \mathbf{G}(k)$ be the k -points of a Zariski connected isotropic simple*

k -algebraic group \mathbf{G} . Let G^+ be the image of the simply connected form \tilde{G} in G under the cover map. Let $H \subseteq G^+$ be a closed non-compact subgroup.

Then for any superreflexive space B and any continuous uniformly equicontinuous linear G -representation $\varrho : G^+ \rightarrow \mathbf{GL}(B)$, $B^{\varrho(H)} = B^{\varrho(G^+)}$ and the canonical complements with respect to both $\varrho(G^+)$ and $\varrho(H)$ coincide, and can be denoted just by B' .

Proof. By Proposition 2.3, we may assume that B is a ucus Banach space and ϱ is a linear isometric representation $\varrho : G \rightarrow \mathbf{O}(B)$. Now the statement follows readily from Theorem 9.1. \square

5.2. The second ingredient is *strong relative property (T)*. It will be used to prove Claim 5.5 below which is the only part which is specific to L^p -like spaces. The rest of the argument applies to all isometric actions on ucus Banach spaces, or all uniformly equicontinuous affine actions on a superreflexive space.

Definition 5.2. Let $H \times U$ be a semi-direct product of locally compact groups. We shall say that it has

strong relative property (T): if for any unitary representation π of $H \times U$ for which H almost has non-trivial invariant vectors, U has invariant vectors.

strong relative property (T_B): where B is a Banach space, if for any linear isometric representation $\varrho : H \times U \rightarrow \mathbf{O}(B)$ the linear isometric H -representation $\varrho' : H \rightarrow \mathbf{O}(B/B^{\varrho(U)})$ does not almost have non-trivial invariant vectors.

Remarks 5.3. (1) The first definition is a variant of “relative property (T)”. The latter usually refers to a pair of groups $G_0 \subseteq G$ and requires that any unitary G -representation with G -almost invariant vectors, has non-trivial G_0 -invariant vectors. Strong relative property (T) for $H \times U$ implies, but is not equivalent to, relative property (T) for $(H \times U, U)$. In fact $\mathbf{SL}_2(\mathbf{R}) \times \mathbf{R}^2$ has the strong relative (T) and so relative (T) as well, whilst its lattice $\mathbf{SL}_2(\mathbf{Z}) \times \mathbf{Z}^2$ does not have strong relative (T) even though the pair $(\mathbf{SL}_2(\mathbf{Z}) \times \mathbf{Z}^2, \mathbf{Z}^2)$ has relative property (T) (cf. M. Burger’s appendix in [HV]).

- (2) If B is a ucus Banach space, then the canonical splitting with respect to $\varrho(U)$, namely $B = B^{\varrho(U)} \oplus B'$ is preserved by $\varrho(H)$ which normalizes U (Corollary 2.8). Hence, as in Remark 2.11, for ucus space B strong relative property (T_B) requires that the *restriction* of $\varrho(H)$ to B' does not almost have invariant vectors. Strong relative ($T_{\mathcal{H}}$) for a Hilbert space \mathcal{H} is equivalent to the strong relative (T).

Lemma 5.4. *A semi-direct product $H \times U$ with strong relative property (T) has strong relative property (T_B) for all L^p -related Banach spaces B of types (i), (ii), (iii) as in Theorem A.*

Proof. This is analogous to the proof of $(T) \Rightarrow (T_B)$ given in Section 4.1. First observe that the extension Theorem 2.20 and a duality argument (based on Proposition 2.10) reduce the statement to the case (i) of $B = L^p(\mu)$.

Thus we assume that $B = L^p(\mu)$ with $p \neq 2$, and $\varrho : H \ltimes U \rightarrow \mathbf{O}(B)$ is a linear isometric representation. Let $B = B^{\varrho(U)} \oplus B'$ be the canonical splitting with respect to U . It is preserved by $\varrho(H)$ because H normalizes U . Now let $\pi = M_{p,2} \circ \varrho \circ M_{2,p}$ be the conjugate of ϱ by the Mazur map. Then π is an orthogonal representation $\pi : H \ltimes U \rightarrow \mathbf{O}(\mathcal{H})$ where $\mathcal{H} = L^2(\mu)$ (Lemma 4.2).

If $H \ltimes U$ fails to have strong relative (T_B) , then there exist $x_n \in S(B')$ so that $\|\varrho(h)x_n - x_n\| \rightarrow 0$ uniformly on compact subsets of H . Uniform continuity of $M_{p,2}$ and the fact that $\text{dist}(S(B'), S(B^{\varrho(U)})) = 1$, imply that for $v_n = M_{p,2}(x_n)$

$$\text{dist}(v_n, \mathcal{H}^{\varrho(U)}) \geq \delta > 0 \quad \|\pi(h)v_n - v_n\| \rightarrow 0$$

uniformly on compact subsets of H . Taking projections of v_n to \mathcal{H}' we show that in this case $H \ltimes U$ does not have strong relative property (T) . \square

5.3. Proof of Theorem B. In view of (the independent) Sections 8.1 and 8.2, more specifically Proposition 8.7(2) and the discussion following Definition 8.2, property (F_B) for $G = \prod G_i$ is inherited by its lattices. Thus it suffices to consider the ambient group $G = \prod G_i$ only. By Proposition 2.15(3) the statement reduces to that about almost-simple factors G_i .

Next we will show that we can assume that G is simply connected. Assume that Theorem B is indeed known for \tilde{G} , we will prove it for G . For any affine isometric action of G on B there is an associated action of \tilde{G} , inflated via the covering map $\tilde{G} \rightarrow G$. \tilde{G} has a fixed point by assumption, hence G has a compact orbit, as the cokernel of the covering map is compact [M5, Theorem 2.3.1(b)]. It follows that G has a fixed point as well.

So we are left proving the theorem for $G = \mathbf{G}(k)$, a higher rank connected, simply-connected, almost-simple group. Using Proposition 2.13, we assume that B is a ucus Banach space and we consider a G -action on B by affine isometries, with $\varrho : G \rightarrow \mathbf{O}(B)$ denoting the linear part of the action. Let $B = B^{\varrho(G)} \oplus B'$ be the canonical decomposition and $\varrho' : G \rightarrow \mathbf{O}(B')$ denote the corresponding sub-representation.

Claim 5.5 (For L^p -like spaces). G contains a direct product $A \times H$ so that

- (1) The restriction $\varrho'|_H : H \rightarrow \mathbf{O}(B')$ does not almost contain invariant vectors.
- (2) A contains a non-trivial semisimple element, and in particular it is not compact.

Proof. Any higher rank almost-simple group $G = \mathbf{G}(k)$ is known to contain a subgroup whose simply-connected cover is isomorphic to either $G_0 = \mathbf{SL}_3(k)$ or $G_0 = \mathbf{Sp}_4(k)$ [M5, Theorem 1.6.2]. In the first case $G_0 = \mathbf{SL}_3(k)$ contains the

semi-direct product $H_0 \times U_0 = \mathbf{SL}_2(k) \times k^2$ embedded in $\mathbf{SL}_3(k)$ as

$$\left\{ \left(\begin{array}{ccc} a & b & x \\ c & d & y \\ 0 & 0 & 1 \end{array} \right) \middle| ad - bc = 1 \right\}$$

where $U_0 \cong k^2$ is the subgroup given by $a = d = 1, b = c = 0$. It is normalized by the copy H_0 of $\mathbf{SL}_2(k)$ embedded in the upper left corner. Let $A_0 \subseteq \mathbf{SL}_3(k)$ be the subgroup $\text{diag}[\lambda, \lambda, \lambda^{-2}]$, $\lambda \in k^*$, which centralizes H_0 in G_0 , and let A and $H \times U$ denote the corresponding subgroups in G .

The semi-direct product $\mathbf{SL}_2(k) \times k^2$ is known to have strong relative property (T). Hence it has strong relative property (T_B) for L^p -related spaces B (Lemma 5.4). By 5.1 we have $B^{\varrho(G)} = B^{\varrho(U)}$ and we have denoted by B' the common canonical complement. Then (1) follows from the strong relative property (T_B) for $H \times U$, while (2) is clear from the construction.

In the second case G contains a copy of $G_0 = \mathbf{Sp}_4(k)$ which is usually defined as a subgroup of $\mathbf{SL}_4(k)$ by

$$\mathbf{Sp}_4(k) = \{g \in \mathbf{SL}_4(k) \mid {}^t g J g = J\}, \quad \text{where} \quad J = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}.$$

The semi-direct product $H_0 \times U_0$ embedded in $\mathbf{SL}_4(k)$ is

$$\left\{ \left(\begin{array}{cc} A & B \\ 0 & {}^t A^{-1} \end{array} \right) \middle| A \in \mathbf{SL}_2(k), {}^t B = A^{-1} B ({}^t A) \right\}$$

with H_0 denoting the image $A \mapsto \text{diag}[A, {}^t A^{-1}]$ of $\mathbf{SL}_2(k)$, and U_0 the normal Abelian subgroup

$$\left\{ \left(\begin{array}{cc} I & B \\ 0 & I \end{array} \right) \middle| {}^t B = B \right\}.$$

The semi-direct product $H_0 \times U_0$ actually lies in $\mathbf{Sp}_4(k)$, it is isomorphic to $\mathbf{SL}_2(k) \times S^2(k)$, where $S^2(k)$ is the space of symmetric bilinear forms on k^2 with the natural $\mathbf{SL}_2(k)$ action. This semi-direct product is also known to have strong relative property (T), and therefore strong relative (T_B) . H_0 is centralized by $A_0 = \{\text{diag}[\lambda, \lambda, \lambda^{-1}, \lambda^{-1}] : \lambda \in k^*\}$. As in the $G_0 = \mathbf{SL}_3(k)$ case, we conclude that the corresponding product $A \times H \subseteq G$ satisfies (1) and (2). The claim is proved. \square

We now turn to the affine isometric G -action defined by a ϱ -cocycle $c \in Z^1(\varrho)$. We shall prove that $c \in B^1(\varrho)$ i.e. that G has a global fixed point. Write $c(g) = c_0(g) + c'(g)$ with $c_0(g) \in B^{\varrho(G)}$ and $c'(g) \in B'$ where $B = B^{\varrho(G)} \oplus B'$ is the canonical splitting. Then $c_0 : G \rightarrow B$ is a homomorphism into the (additive) Abelian group. As G has compact Abelianization, $c_0(g) \equiv 0$, which means that the affine G -action preserves each affine subspace $p + B'$. Hence both the affine G -action and the representation can be restricted to B' .

Claim 5.5 provides an input for the following general lemma:

Lemma 5.6. *Let a direct product of topological groups $A \times H$ act by affine isometries on a Banach space B . Suppose that the associated linear isometric representation ϱ restricted to H does not almost have invariant vectors. Then the affine action of A has bounded orbits in B . In particular, if B is uniformly convex, then A has a fixed point in B .*

Remark 5.7. In the uniformly convex case, this follows of course from the stronger splitting theorem (Theorem C); compare also with Theorem 7.1 below for the weaker assumption that the *product* does not almost have invariant vectors.

Proof of the lemma. Let $\varrho : A \times H \rightarrow \mathbf{O}(B)$ and $c \in Z^1(\varrho)$ denote the associated linear isometric representation and the translation cocycle. The commutation relation between any $h \in H$ and $a \in A$ gives

$$c(h) + \varrho(h)c(a) = c(ha) = c(ah) = c(a) + \varrho(a)c(h)$$

which can be rewritten as

$$(I - \varrho(h))c(a) = (I - \varrho(a))c(h).$$

By the assumption on $\varrho(H)$, there exists a compact subset $K \subseteq H$ and an $\varepsilon > 0$ so that $\max_{h \in K} \|\varrho(h)v - v\| \geq \varepsilon \cdot \|v\|$ for all $v \in B$. Let $R = \max_{h \in K} \|c(h)\| < \infty$. Then for $a \in A$

$$\varepsilon \cdot \|c(a)\| \leq \max_{h \in K} \|(I - \varrho(h))c(a)\| \leq 2R.$$

Hence $\sup_{a \in A} \|c(a)\| \leq 2R/\varepsilon$, i.e. the A -orbit of 0 is bounded. If B is uniformly convex then the circumcentre of this orbit is an A -fixed point as in Lemma 2.14. \square

Let us restrict the G -action to B' . For $g \in G$ let $\text{Fix}(g)$ denote the set of g -fixed points in B' . It follows from Claim 5.5 and Lemma 5.6 that for some non-elliptic semisimple element $a \in A \subseteq G$, $\text{Fix}(a) \subseteq B'$ is non-empty.

Note that $\text{Fix}(a)$ cannot contain more than one point. Indeed, if $x, y \in \text{Fix}(a)$ then $x - y \in B'$ is a fixed vector for the linear isometry $\varrho(a)$. Since the cyclic group $\langle a \rangle$ is unbounded, the ucus analogue of Moore's ergodicity 5.1 implies that $B^{\varrho(\langle a \rangle)} = B^{\varrho(G)}$. Hence $x - y = 0$. Thus $\text{Fix}(a) = \{x_0\}$.

If $g, h \in G$ commute then $\text{Fix}(g)$ is an h -invariant set. So if $\langle g \rangle$ is unbounded and $\text{Fix}(g) \neq \emptyset$ then $\text{Fix}(g)$ is a single point fixed by h . If $\langle h \rangle$ is also unbounded then $\text{Fix}(g) = \text{Fix}(h)$. Hence the following lemma implies that $G \cdot x_0 = x_0$:

Lemma 5.8. *For any two non-elliptic semisimple elements $g, h \in G$ there is a chain $g = g_1, g_2, \dots, g_n = h$ of non-elliptic semisimple elements, each commuting with its successor. Furthermore, the set of non-elliptic semisimple elements is dense in G .*

Proof. The first claim is equivalent to the connectivity of the Tits boundary. Both claims are well-known. \square

This concludes the proof of Theorem B. \square

Remark 5.9. Observe that the somewhat restrictive assumption that B is an L^p -related space is used only in the proof of Claim 5.5, the rest of the argument being in the context of general ucus Banach spaces.

6. MINIMAL SETS

Let B be a strictly convex reflexive Banach space and G a group acting on B by isometries. Consider the ordered category \mathcal{C} of non-empty closed convex G -invariant subsets of B endowed with G -equivariant isometric maps and inclusion order. The goal of this section is to study minimal elements of \mathcal{C} (regardless of whether they exist). In Section 7 we shall prove their existence, under certain conditions (see Corollary 7.6).

The Mazur-Ulam theorem states that a surjective isometry between Banach spaces is affine. It is not known (and probably not true under no further assumptions) whether the analogous of the Mazur-Ulam theorem holds in the general context of convex subsets of Banach spaces. However, for subsets of strictly convex spaces it is obviously true:

Lemma 6.1. *Let $C \subseteq B$ be a convex subset. Then every isometric map $C \rightarrow B$ is affine.*

Proof. It is enough to show that for all $x, y \in C$ and every $0 < t < 1$ the point $p = tx + (1 - t)y$ is determined metrically. This is true since by strict convexity

$$\overline{B}(x, (1 - t)\|x - y\|) \cap \overline{B}(y, t\|x - y\|) = \{p\}.$$

□

In particular the morphisms of \mathcal{C} are affine. Another useful geometric property of closed convex sets in B is the existence of a nearest point projection.

Lemma 6.2. *Let C be a non-empty closed convex subset of B . Then for every $x \in B$ there exist a unique point $\pi_C(x) \in C$ such that $\|x - \pi_C(x)\| = d(x, C)$.*

Proof. The uniqueness follows from strict convexity. By the Hahn-Banach theorem C is weakly closed since it is closed and convex; therefore, by reflexivity and the Banach-Alaoglu theorem we have a nested family $C \cap \overline{B}(x, d)$ of weakly compact sets as $d \searrow d(x, C)$; its intersection yields existence. □

The map $\pi_C : B \rightarrow C$ is called the *nearest point projection* on C . We remark that it is not continuous in general. It is continuous for uniformly convex Banach spaces and non-expanding for Hilbert spaces. Still, the distance between a point and its projection is always a 1-Lipschitz function:

Lemma 6.3. *Let C be a non-empty closed convex subset of B . Then the function $x \mapsto \|\pi_C(x) - x\|$ from C to \mathbf{R} is 1-Lipschitz.*

Proof. For any $x, y \in C$

$$\|\pi_C(x) - x\| \leq \|\pi_C(y) - x\| \leq \|\pi_C(y) - y\| + \|y - x\|.$$

□

Lemma 6.4. *If $C \in \mathcal{C}$ is a minimal element, then any convex G -invariant continuous (or lower semi-continuous) function $\varphi : C \rightarrow \mathbf{R}$ is constant.*

Proof. If φ were to assume two distinct values $s < t$, then $\varphi^{-1}((-\infty, s])$ would be a strictly smaller element of \mathcal{C} . \square

Lemma 6.5. *Let $C, C' \in \mathcal{C}$ with C minimal. Then the nearest point projection $\pi = \pi_{C'}|_C : C \rightarrow C'$ is affine.*

Proof. For every $x, y \in C$ and $t \in [0, 1]$, the definition of π implies

$$(6.1) \quad \begin{aligned} \|\pi(tx + (1-t)y) - (tx + (1-t)y)\| \\ \leq \|(t\pi(x) + (1-t)\pi(y)) - (tx + (1-t)y)\| \\ \leq t\|\pi(x) - x\| + (1-t)\|\pi(y) - y\|. \end{aligned}$$

It follows that the function $C \rightarrow \mathbf{R}$, $x \mapsto \|\pi(x) - x\|$ is convex. Clearly it is G -invariant, and by Lemma 6.3 it is continuous, hence Lemma 6.4 implies that $\|\pi(x) - x\|$ is constant on C . This constant must be $d(C, C')$; as both the right-hand side and the left-hand side in (6.1) equal $d(C, C')$, it follows that

$$\|(t\pi(x) + (1-t)\pi(y)) - (tx + (1-t)y)\| = \|\pi(tx + (1-t)y) - (tx + (1-t)y)\|.$$

Therefore, by the uniqueness part of Lemma 6.2, $t\pi(x) + (1-t)\pi(y)$ must be $\pi(tx + (1-t)y)$. \square

Lemma 6.6. *If $C \in \mathcal{C}$ is minimal and $T : C \rightarrow B$ is a G -equivariant affine map, then there exist a $\varrho(G)$ -invariant vector $b \in B$ such that $T(c) = c + b$ for all $c \in C$.*

Proof. The map $C \rightarrow \mathbf{R}$, $x \mapsto \|Tx - x\|$ is G -invariant, continuous and convex, hence by Lemma 6.4 it has a constant value $d \geq 0$. Since B is strictly convex and C is convex, the affine map $\sigma(x) = Tx - x$ from C to the sphere of radius d in B must be constant. Its value $b = \sigma(C)$ is the desired ($\varrho(G)$ -invariant) translation vector. \square

Corollary 6.7. *The map $\pi_C : C \rightarrow C'$ from Lemma 6.5 is in fact a translation.*

Corollary 6.8. *If $C, C' \in \mathcal{C}$ are minimal, then they are equivariantly isometric. Moreover, any equivariant isometry $C \rightarrow C'$ is a translation by a $\varrho(G)$ -invariant vector.*

Proof. By Corollary 6.7, $\pi_{C'}|_C : C \rightarrow C'$ is an isometry; it is G -equivariant and hence onto by minimality of C' . The second claim follows from Lemma 6.1 and Lemma 6.6. \square

7. ACTIONS OF PRODUCT GROUPS AND SPLITTING

7.1. The main goal of this section is to prove Theorem C. By Proposition 2.13 we may assume the affine action to be isometric with respect to a ucus norm on a Banach space B . The main step is the following theorem.

Theorem 7.1. *Let $G = G_1 \times G_2$ be a product of topological groups with a continuous action by isometries on a uniformly convex Banach space B without G -fixed point. Assume that the associated linear G -representation ϱ does not almost have non-zero invariant vectors. Then there exists a non-zero $\varrho(G_i)$ -invariant vector for some $i \in \{1, 2\}$.*

The proof of Theorem 7.1 uses minimal sets (in analogy to [Mo]); notice that we are in the setting of Section 6 since uniformly convex spaces are reflexive and strictly convex [BL, App. A]. More precisely, we show:

Proposition 7.2. *Let G and B be as above. Then there exists a minimal non-empty closed convex G_1 -invariant subset in B . In fact, any non-empty closed convex G_1 -invariant subset contains a minimal such subset.*

Remark 7.3. In view of the additional statement of Proposition 7.2, the set C_0 of Corollary 7.6 is contained in every non-empty closed convex G -invariant subset. Thus it is indeed the (non-empty) intersection of all those subsets.

Proof of Theorem 7.1. Proposition 7.2 provides a minimal non-empty closed convex G_1 -invariant set $C \subseteq B$. If there is no non-zero $\varrho(G_1)$ -invariant vector, Lemma 6.6 (applied to G_1) shows that G_2 fixes every point of C . Since G_1 preserves C and G has no fixed point, C cannot consist of a single point. Picking two distinct points $x, y \in C$ yields the non-zero $\varrho(G_2)$ -invariant vector $x - y$. \square

Recall that uniform convexity is characterized by the positivity of the convexity modulus δ defined in Section 2.1. Moreover, δ is a positive, non-decreasing function which tends to zero at zero. Defining

$$\delta^{-1}(t) = \sup\{\varepsilon : \delta(\varepsilon) \leq t\},$$

δ^{-1} is easily seen to share the same properties. Furthermore, for every $\varepsilon > 0$, $\delta^{-1} \circ \delta(\varepsilon) \geq \varepsilon$.

Proof of Proposition 7.2. Let $C_0 \subseteq B$ be any non-empty closed convex G_1 -invariant subset; we will show that C_0 contains a minimal subset (if no initial C_0 was prescribed, one may choose $C_0 = B$).

Pick any $p \in C_0$ and let $C_1 \subseteq C_0$ be the closed convex hull of the G_1 -orbit of p . By Hausdorff's maximal principle, we can choose a maximal chain \mathcal{D} of non-empty closed convex G_1 -invariant subsets of C_1 . If $b_C := \pi_C(0)$ is bounded as C ranges over \mathcal{D} , then for some $R > 0$ we have a nested family of non-empty sets $\overline{B}(0, R) \cap C$ which are weakly compact by reflexivity, Hahn-Banach theorem and Banach-Alaoglu theorem. In particular the intersection $\bigcap \mathcal{D}$ is non-empty, thus providing a minimal set for G_1 . Therefore, we may from now on assume for a contradiction that the (non-decreasing) net $R_C := \|b_C\|$ is unbounded over $C \in \mathcal{D}$. Let $\mathcal{D}' \subseteq \mathcal{D}$ be the cofinal segment defined by $R_C > 0$. We will obtain a contradiction by showing that for every compact $K \subseteq G$, $\text{diam}(\varrho(K)\hat{b}_C)$ tends to zero along $C \in \mathcal{D}'$, where $\hat{b}_C = \frac{b_C}{R_C}$.

Indeed, choose $K_i \subseteq G_i$ compact with $K \subseteq K_1 \times K_2$ and let $L = \max_{g \in K_1 \times K_2} \|g \cdot 0\|$. The choice of b_C implies $g \cdot b_C \neq 0$ and $R_C \leq \|\frac{b_C + g \cdot b_C}{2}\|$ for all $g \in G_1$.

Therefore, setting $x = \frac{b_C}{\|g \cdot b_C\|}$, $y = \frac{g \cdot b_C}{\|g \cdot b_C\|}$, the convexity modulus $\delta_{C,g} := \delta(\|x - y\|)$ gives

$$\begin{aligned} R_C &\leq \left\| \frac{1}{2} b_C + \frac{1}{2} g \cdot b_C \right\| \leq \left\| \frac{1}{2} (x + y) \right\| \cdot \|g \cdot b_C\| \leq (1 - \delta_{C,g}) \cdot (\|g \cdot b_C - g \cdot 0\| + \|g \cdot 0\|) \\ &\leq (1 - \delta_{C,g})(R_C + L) \leq R_C \left(1 + \frac{L}{R_C} - \delta_{C,g}\right) \quad \forall g \in K_1. \end{aligned}$$

Therefore $\delta_C := \sup_{g \in K_1} \delta_{C,g} \leq \frac{L}{R_C} \rightarrow 0$ along $C \in \mathcal{D}'$ and hence

$$\sup_{g \in K_1} \frac{\|g \cdot b_C - b_C\|}{\|g \cdot b_C\|} \leq \delta^{-1}(\delta_C) \rightarrow 0.$$

Using $\|g \cdot b_C\| \leq \|g \cdot b_C - g \cdot 0\| + L \leq R_C + L$, it follows that

$$(7.1) \quad \sup_{g \in K_1} \frac{\|g \cdot b_C - b_C\|}{R_C} \rightarrow 0 \quad \text{along } C \in \mathcal{D}'.$$

On the other hand, for every $g \in G_2$, the function $z \mapsto \|g \cdot z - z\|$ is continuous, convex and G_1 -invariant; therefore, it is bounded by $\|g \cdot p - p\|$ on C_1 . Setting $L' = \max_{g \in K_2} \|g \cdot p - p\|$, it follows now that for all $k = (g_1, g_2) \in K$ we have

$$\begin{aligned} R_C \cdot \|\varrho(k) \hat{b}_C - \hat{b}_C\| &= \|k b_C - b_C - k 0\| \leq \|g_1 b_C - b_C\| + \|g_2 g_1 b_C - g_1 b_C\| + L \\ &\leq \|g_1 b_C - b_C\| + L' + L. \end{aligned}$$

Thus in view of (7.1) $\text{diam}(\varrho(K) \hat{b}_C)$ goes to zero as claimed. \square

Proof of Theorem C. We adopt the notation and assumptions of that theorem; let ϱ be the linear part of the action. Assume first $n = 2$. Since we have in particular $B^{\varrho(G)} = 0$, Corollary 2.9 yields a canonical splitting $B = B^{\varrho(G_1)} \oplus B^{\varrho(G_2)} \oplus B_0$ invariant under $\varrho(G)$. Decomposing the cocycle $G \rightarrow B$ along this splitting shows that up to affine isometry we may assume that the affine G -space B splits likewise as affine product of affine spaces with corresponding linear parts. However, Theorem 7.1 shows that the resulting affine G -action on B_0 must have a fixed point since $B_0^{\varrho(G_i)} = 0$. Therefore we obtain a G -invariant affine subspace G -isometric to $B^{\varrho(G_1)} \oplus B^{\varrho(G_2)}$ in B , as claimed.

In order to obtain the general case $n \geq 2$, we only need to observe that Corollary 2.9 applied to the product $G_1 \times \prod_{i \geq 2} G_i$ allows us to apply induction on n . \square

Remark 7.4. The above proof characterizes as follows the subspaces $B_i \subseteq B$ appearing in the statement of Theorem C: Upon possibly replacing the B_i with the corresponding linear subspace (which corresponds to replacing the cocycles with cohomologous cocycles), we have $B_i = B^{\varrho(G'_i)}$ for $G'_i = \prod_{j \neq i} G_j$.

7.2. A More Geometric Approach to Theorem B and a Step Towards Conjecture 1.6. Before going on towards the superrigidity theorem, let us explain a more geometric, and seemingly more general, approach to prove $(T_B) \Rightarrow (F_B)$, which is based on minimal sets. First we shall formulate a very general statement in the vein of Conjecture 1.6:

Theorem 7.5. *Let B be a ucus Banach space and G a topological group with property (T_B) and compact Abelianization. Then for any continuous affine isometric action of G on B there is a minimal non-empty closed convex subset $C \subseteq B$. Moreover $\text{Aut}_G(C)$ is trivial, $C \subseteq B'$ and C is unique up to translations by a $\varrho(G)$ -invariant vector.*

The proof of Theorem 7.5 relies on the following consequence of our discussion of minimal sets:

Corollary 7.6. *Let G be a topological group with a continuous action by isometries on a uniformly convex Banach space B . Assume that the associated linear representation does not almost have non-zero invariant vectors. Then there exists a unique minimal non-empty closed convex G -invariant subset $C_0 \subseteq B$. Moreover, there are no non-trivial G -equivariant isometries of C_0 .*

Proof of Corollary 7.6. For the existence of C_0 , we may apply Proposition 7.2 if $G = G_1 \times 1$ has no fixed point, or otherwise take such a fixed point for C_0 . Both uniqueness and the additional statement follow now from Corollary 6.8. \square

Proof of Theorem 7.5. Since G has compact Abelianization, the $\varrho(G)$ invariant subspace B' is in fact G -invariant as an affine space, as the projection of the cocycle to $B^{\varrho(G)}$ must be a homomorphism. It follows that every minimal non-empty closed convex G -invariant set is contained in some coset of B' . The existence and uniqueness of such subset C inside B' follows from Corollary 7.6. The fact that any two such sets are different by a $\varrho(G)$ -invariant vector is a consequence of Corollary 6.8. \square

Let us now describe an alternative proof for Theorem B. Let B be an L^p -related Banach space as in Theorem B. Most of the proof is carried out along the same lines as the first proof given in Section 4. We reduce to the case where G is connected, simply-connected and almost-simple. As explained in Section 4, it is enough to show that for some non-elliptic semisimple element $a \in G$ the set $\text{Fix}(a) \in B$ is a singleton. Now G either contains a copy of $\mathbf{SL}_3(k)$ or of $\mathbf{Sp}(4, k)$ that, in each case, contains a semidirect product $H \rtimes U$ with the strong relative property (T_B) (see Lemma 5.4 and the proof of Claim 5.5). We decompose $B = B^{\varrho(U)} \oplus B'$ according to that U action (note that by Howe-Moore the $B^{\varrho(U)} = B^{\varrho(G)}$). Then B' is invariant under the affine action of G and H does not almost have invariant vectors in B' . Hence, by Corollary 7.6 there is a unique minimal non-empty closed convex H -invariant subset $C \subseteq B'$ and it has no non-trivial automorphisms which commute with the H action. Since, by Claim 5.5, the centralizer of the H is non-compact it follows by Howe-Moore that $|C| = 1$. We can then take a to be any non-elliptic semisimple in the H copy, or in its centralizer.

8. INDUCTION AND SUPERRIGIDITY

Let $\Gamma < G = G_1 \times \cdots \times G_n$ be a lattice in a product of $n \geq 2$ locally compact groups. Under an irreducibility assumption, the splitting theorem (Theorem C)

implies a superrigidity result for uniformly equicontinuous affine Γ -actions on superreflexive spaces B . As before such an action can be viewed as an affine isometric Γ -action on a ucus Banach space B . It therefore suffices to apply the splitting theorem to the *induced* G -action on an *induced space* $L^p(G/\Gamma, B)$ (compare [Sh] for the Hilbertian case).

The goal of this section is to address the various (mostly technical) issues that arise when carrying out this programme. We begin by preparing for a statement (Theorem 8.3 below) that will then imply a more general form of Theorem D.

8.1. Let G be a locally compact group and $\Gamma < G$ a lattice. The induction procedure will work smoothly if Γ is uniform (i.e. cocompact); in order to treat some non-uniform cases, one introduces the following.

Definition 8.1 ([M5, III.1.8]). The lattice Γ is *weakly cocompact* if the G -representation $L_0^2(G/\Gamma)$, i.e. the orthogonal complement of the trivial representation in $L^2(G/\Gamma)$, does not almost have non-zero invariant vectors.

One verifies that any cocompact lattice is weakly cocompact. If G has property (T), then all its lattices are weakly cocompact. This also holds if G is any connected semisimple Lie group ([Be], compare also [M5, II.1.12]).

Definition 8.2 (See [Sh, 1.II]). Let $p > 0$. The lattice Γ is *p-integrable* if either (i) it is uniform; or (ii) it is finitely generated and for some (or equivalently any) finite generating set $S \subseteq \Gamma$, there is a Borel fundamental domain $\mathcal{D} \subseteq G$ (with null boundary) such that

$$\int_{\mathcal{D}} \|\chi(g^{-1}h)\|_S^p dh < \infty \quad \forall g \in G,$$

where $\|\cdot\|_S$ is the word-length associated to S and $\chi : G \rightarrow \Gamma$ is defined by $\chi^{-1}(e) = \mathcal{D}$, $\chi(g\gamma^{-1}) = \gamma\chi(g)$.

This formulation is a bit awkward so as to include all uniform lattices since (ii) would otherwise fail when G is not compactly generated. Condition (ii) holds (with any $p \geq 1$) for all irreducible lattices in higher rank semisimple Lie/algebraic groups, see [Sh, §2]; it holds likewise for Rémy's Kac-Moody lattices [R].

Finally, given a product structure $G = G_1 \times \cdots \times G_n$, we say that a lattice $\Gamma < G$ is *irreducible* if its projection to each G_i is dense.

Theorem 8.3. *Let Γ be an irreducible lattice in a locally compact σ -compact group $G = G_1 \times \cdots \times G_n$. Let B be a ucus Banach space with a Γ -action by isometries. Assume that Γ is weakly cocompact and p-integrable for some $p > 1$.*

If the associated linear Γ -representation does not almost have invariant vectors, then there is a Γ -closed complemented affine subspace of B on which the Γ -action is a sum of actions extending continuously to G and factoring through $G \rightarrow G_i$. (Compare Remark 1.9.)

In order to justify that Theorem 8.3 implies Theorem D, there are two points to address: First, Proposition 2.13 allows us to assume indeed that the topological vector space of Theorem D is in fact a ucus Banach space with a Γ -action by

isometries. Secondly we need to consider the case where the associated linear Γ -representation ϱ does almost have invariant vectors. We then have the assumption that G has property (T_B) and compact Abelianization. Let $B = B^e \oplus B'$ be the decomposition of Proposition 2.6 and decompose the cocycle $b : \Gamma \rightarrow B$ of the Γ -action accordingly as $b = b_0 + b'$. Then $b_0 = 0$ since G has compact Abelianization and property (T_B) ensures that we can apply Theorem 8.3 to the action on B' determined by b' .

8.2. Induction. Throughout this section, G is a locally compact second countable group and $\Gamma < G$ a lattice. In particular, the Haar measure induces a standard Lebesgue space structure on G/Γ .

Remark 8.4. Even though Theorem 8.3 and Theorem D was stated in the more general setting of σ -compact groups, it is indeed enough to treat the second countable case: one can reduce to the latter by a structural result of Kakutani-Kodaira [KK] (the details of the straightforward reduction are expounded at length in [Mo]).

Let B be any Banach space and $1 < p < \infty$. We consider the Banach space $E = L^p(G/\Gamma, B)$ as in Section 2.5.

Lemma 8.5. *If B is uniformly convex or ucus, then so is E .*

Proof. This follows from a result of Figiel and Pisier, see Theorem 1.e.9 point (i) in [LT], Volume II. \square

Suppose now that B is endowed with a linear isometric Γ -representation ϱ . Then E can be canonically isometrically identified

$$(8.1) \quad E \cong L^{[p]}(G, B)^{\varrho(\Gamma)}$$

with the space of those Bochner-measurable Γ -equivariant function classes $f : G \rightarrow B$ such that $\|f\|_B : G/\Gamma \rightarrow \mathbf{R}$ is p -integrable (the latter condition is symbolized by the notation $L^{[p]}$). Here, we choose to interpret Γ -equivariance as $f(g\gamma) = \varrho(\gamma)^{-1}f(g)$. The isomorphism (8.1) can be e.g. realized by restricting equivariant maps to any Borel fundamental domain $\mathcal{D} \subseteq G$ for Γ since $\mathcal{D} \cong G/\Gamma$ as Lebesgue spaces. This identification allows us to endow E with a continuous linear isometric G -representation by left multiplication. This G -representation is called the *induced representation*. If we choose a fundamental domain $\mathcal{D} \subseteq G$ and consider the corresponding map χ as in Definition 8.2, then this G -representation reads as follows for $f \in E = L^p(G/\Gamma, B)$:

$$(8.2) \quad (hf)(g\Gamma) = \varrho(\chi(g)^{-1}\chi(h^{-1}g))f(h^{-1}g\chi(h^{-1}g)\Gamma)$$

(a good indication that the model (8.1) is more natural!).

Lemma 8.6. *Assume Γ weakly cocompact in G . If the linear Γ -representation does not almost have invariant vectors, then the induced linear G -representation does not either.*

Proof. The proof given by Margulis in the unitary case [M5, III.1.11] holds without changes. \square

Suppose now that B is endowed with an isometric Γ -action – not necessarily linear anymore. We want to endow E with a continuous affine isometric G -action by identifying E with a space of Γ -equivariant function classes $G \rightarrow B$ as before, except that equivariance is now understood with respect to the affine Γ -action. Formally, there is nothing to change to the special case of linear action considered above; the action is defined by left G -translation of equivariant maps, so that via the natural identification we get for $f \in E = L^p(G/\Gamma, B)$ the action

$$(8.3) \quad (hf)(g\Gamma) = \chi(g)^{-1}\chi(h^{-1}g)f(h^{-1}g\chi(h^{-1}g)\Gamma)$$

in complete analogy with (8.2). However, the L^p integrability property might be lost. The condition (ii) of Definition 8.2 is a straightforward sufficient condition to retain integrability; cocompactness of Γ is also enough, because it ensures that one can choose \mathcal{D} in such a way that for any compact $C \subseteq G$ the set $\{\eta \in \Gamma : \mathcal{D}\eta \cap C \neq \emptyset\}$ is finite [B2, VII §2 Ex. 12]. Compare [Sh, §2] (and [Mo, App. B]).

In conclusion, we may always consider the continuous *induced (affine) isometric G -action* on E when Γ is p -integrable.

By construction, the linear part of the induced affine action coincides with the induced linear G -representation on E considered earlier. If we denote by $b : \Gamma \rightarrow B$ the cocycle of the original affine Γ -action, then comparing (8.2) with (8.3) shows that the cocycle $\tilde{b} : G \rightarrow E$ of the induced affine action is given by

$$(8.4) \quad \tilde{b}(h)(g\Gamma) = b\left(\chi(g)^{-1}\chi(h^{-1}g)\right).$$

Moreover, the correspondence $b \mapsto \tilde{b}$ induces a (topological) isomorphism $H^1(\Gamma, B) \rightarrow H^1(G, E)$.

At this point, we record the following.

Proposition 8.7. *Keep the notation of this section.*

- (1) *If Γ has property (F_B) then so does G .*
- (2) *If G has property (F_E) and Γ is p -integrable, then Γ has property (F_B) .*

Proof. For (1), consider any continuous isometric G -action on B ; then there is a Γ -fixed point $b \in B$. The corresponding orbit map $G \rightarrow B$ descends to a continuous map $G/\Gamma \rightarrow B$. The image of the normalized invariant measure on G/Γ in B being preserved by G , it follows from Lemma 2.14 that there is a G -fixed point.

For (2), consider an isometric Γ -action on B and endow E with the induced affine action as in the discussion above. Then there is a G -fixed point $f \in E$. It follows from the description of E as space of equivariant maps that f is essentially constant and that its essential value is a Γ -fixed point of B . \square

8.3. Superrigidity. In order to prove Theorem 8.3, we now analyse the interplay between the induction constructions and the setting of irreducible lattices $\Gamma < G = G_1 \times \cdots \times G_n$ as in the beginning of this Section 8. We will roughly imitate the arguments given by Shalom in [Sh] when he deduces Corollary 4.2 *ibid*.

Keep all the above notations and write $G'_i = \prod_{j \neq i} G_j$. First we observe that the irreducibility of Γ implies that for each i it is a well-posed definition to consider

the maximal (possibly zero) linear subspace $B_i \subseteq B$ on which the linear Γ -representation ϱ extends to a continuous G -representation $\varrho_i : G \rightarrow G_i \rightarrow \mathbf{O}(B_i)$ factoring through G_i ; moreover B_i is automatically closed by maximality.

The induced space E is ucus by Lemma 8.5. The isometric (affine) G -action on E has no fixed point by the very same argument given to prove Proposition 8.7(2). On the other hand, the linear part does not have almost invariant vectors by Lemma 8.6. Thus Theorem C applies: There is a G -invariant closed complemented affine subspace $\underline{E} \subseteq E$ and an affine isometric G -equivariant isomorphism $\underline{E} \cong E_1 \oplus \cdots \oplus E_n$, where each E_i is a ucus space with an affine isometric G -action factoring through $G \rightarrow G_i$. In view of Remark 7.4, there is no loss of generality in assuming that E_i is the space of G'_i -fixed under the induced linear representation. One verifies readily the following:

Lemma 8.8. *The map $B_i \rightarrow E \cong L^{[p]}(G, B)^{\varrho(\Gamma)}$ that to $v \in B_i$ associates the function $G \rightarrow B$ defined by $g \mapsto \varrho_i(g^{-1})v$ yields an isometric isomorphism of (linear) G -spaces $B_i \cong E_i$. \square*

Indeed, since the image of Γ in G_i is dense, the Fubini-Lebesgue theorem implies that any map $f : G \rightarrow B$ in E that is G'_i -invariant in the linear representation on E is an orbit map as in the lemma.

At this point we observe that if the subspaces B_i had trivial intersection, we would indeed have found a subspace $\bigoplus B_i \cong \bigoplus E_i$ of B on which the affine Γ -action extends continuously to G as requested. In general, we have a Γ -equivariant affine map

$$\bigoplus E_i \longrightarrow \sum B_i \subseteq B$$

induced by the maps of Lemma 8.8. Alternatively, we can think of this map as follows: The cocycle induced as in (8.4) decomposes as a sum of cocycles $\tilde{b} = \bigoplus \tilde{b}_i : G \rightarrow E$, $\tilde{b}_i : G \rightarrow G_i \rightarrow E_i$, and in turn by Lemma 8.8 each \tilde{b}_i is the cocycle induced under the correspondence (8.4) from a cocycle $b_i : \Gamma \rightarrow B_i$; the affine Γ -action on $\sum B_i$ is determined by the cocycle $\sum b_i$. This completes the proof of Theorem 8.3. \square

Remark 8.9. As mentioned in Remark 1.9, the obstruction to extending the affine Γ -action on some subspace of B is confined within a compact group. Indeed, the only reason we might end up with a *sum* of action extending to G through various G_i rather than with a direct sum (which then extends globally to G) is the possibility that $B_i \cap B_j \neq 0$ for some $i \neq j$. But then the linear representation of Γ on $B_i \cap B_j$ extends continuously to G in two different ways, *both* through G_i and through G_j . This may indeed happen but forces the image of Γ in $\mathbf{O}(B_i \cap B_j)$ to be compact, see examples and discussion in [Mo].

Let us only mention the most basic example: $\Gamma < G = G_1 \times G_2$ with $G_i = \mathbf{Z} \rtimes \{\pm 1\}$ and $\Gamma = \mathbf{Z}^2 \rtimes \{\pm 1\}$. Then Γ acts affinely isometrically without fixed point on $B = \mathbf{R}$ (by $(n, m; \varepsilon).x = \varepsilon x + n + m$) and the associated linear representation does not almost have invariant vectors. However, it is easy to check that this action does not extend to G . Instead, it is a sum of actions extending to G_i with sum map $\mathbf{R} \oplus \mathbf{R} \rightarrow B = \mathbf{R}$. Here $B_1 = B_2 = B$.

9. APPENDIX: HOWE-MOORE THEOREM ON BANACH SPACES

In this appendix we sketch the proof of a version of the well known Howe-Moore's theorem on vanishing of matrix coefficients for unitary representations, extended to the framework of uc Banach spaces. This generalization is due to Yehuda Shalom (unpublished) and we state it here with a sketch of the proof for reader's convenience.

Theorem 9.1. *Let I be a finite set, k_i , $i \in I$ be local fields, \mathbf{G}_i a connected isotropic semisimple and simply-connected k_i -groups, $G_i = \mathbf{G}_i(k_i)$ the locally compact group of k_i points, and $G = \prod_{i \in I} G_i$.*

Let B be a uc Banach space and $\varrho : G \rightarrow \mathbf{O}(B)$ a continuous isometric linear representation, such that $B^{\varrho(G_i)} = \{0\}$ for each $i \in I$. Then all matrix coefficients $c_{x,\lambda}(g) = \langle \varrho(g)x, \lambda \rangle$, $x \in B$, $\lambda \in B^$, vanishes at infinity, i.e. $c_{x,\lambda} \in C_0(G)$.*

Proof of Theorem 9.1. In a way of contradiction, assume that for some $g_n \rightarrow \infty$ in G , $v \in S(B)$, $\lambda \in S(B^*)$ one has

$$\inf |\langle \varrho(g_n)x, \lambda \rangle| = \varepsilon > 0.$$

We shall prove that at least one simple factor G_i of G has a non-trivial $\varrho(G_i)$ invariant vector.

Let $G = KAK$ be a Cartan decomposition of G (here $K = \prod K_i$ and $A = \prod A_i$ where $G_i = K_i A_i K_i$ is the Cartan decomposition for G_i). We first show that without loss of generality one may assume $g_n \in A$.

Lemma 9.2 (KAK Reduction). *There exists a sequence $a_n \rightarrow \infty$ in the Cartan subgroup $A \subseteq G$ and non-zero vectors $y, z \in B$ so that*

$$\varrho(a_n)y \xrightarrow{w} z \neq 0.$$

where \xrightarrow{w} denotes the weak convergence.

Proof. Write $g_n = k_n a_n k'_n$ where $k_n, k'_n \in K$ and $a_n \in A$. Then $a_n \rightarrow \infty$ because $g_n \rightarrow \infty$. Upon passing to a subsequence, $k'_n \rightarrow k' \in K$ and $k_n \rightarrow k \in K$. Denote

$$y_n = \varrho(k'_n)x, \quad y = \varrho(k')x, \quad \mu_n = \varrho^*(k_n^{-1})\lambda, \quad \mu = \varrho^*(k^{-1})\lambda$$

where ϱ^* is the dual (contragradient) G -representation on B^* . Using the weak-compactness of the unit ball of B we may also assume that

$$\varrho(a_n)y \xrightarrow{w} z.$$

We shall show that $\langle z, \mu \rangle = \lim \langle \varrho(g_n)x, \lambda \rangle$ which is bounded away from zero, hence implying $z \neq 0$.

Recall that in a uc Banach space B the weak and the strong topologies agree on the unit sphere $S(B)$: indeed if $y_n \xrightarrow{w} y$ are unit vectors, then

$$1 \leftarrow \langle (y_n + y)/2, y^* \rangle \leq \|y_n + y\|/2 \leq 1 - \delta(\|y_n - y\|).$$

Hence $\delta(\|y_n - y\|) \rightarrow 0$ and $\|y_n - y\| \rightarrow 0$. For the same reason we also have $\|\mu_n - \mu\| \rightarrow 0$ in $S(B^*)$. For an arbitrary $\xi \in B^*$

$$|\langle \varrho(a_n)y_n, \xi \rangle - \langle \varrho(a_n)y, \xi \rangle| \leq \|y_n - y\| \cdot \|\xi\| \rightarrow 0.$$

Hence $\varrho(a_n)y_n \xrightarrow{w} z$. In general, if $z_n \xrightarrow{w} z$ in B and $\mu_n \xrightarrow{w} \mu$ in B^* then $\langle z_n, \mu_n \rangle \rightarrow \langle z, \mu \rangle$ because weakly convergent sequences are bounded in norm and

$$\begin{aligned} |\langle z_n, \mu_n \rangle - \langle z, \mu \rangle| &\leq |\langle z_n, \mu_n - \mu \rangle| + |\langle z_n - z, \mu \rangle| \\ &\leq (\sup \|z_n\|) \cdot \|\mu_n - \mu\|^* + |\langle z_n - z, \mu \rangle| \rightarrow 0. \end{aligned}$$

Therefore

$$\langle \varrho(g_n)x, \lambda \rangle = \langle \varrho(a_n k'_n)x, \varrho^*(k_n^{-1})\lambda \rangle = \langle \varrho(a_n)y_n, \mu_n \rangle \rightarrow \langle z, \mu \rangle$$

implying $|\langle z, \mu \rangle| \geq \varepsilon$, which in particular means that $z \neq 0$. \square

Lemma 9.3 (Generalized Mautner Lemma). *Suppose that $\{a_n\}$ and h in G satisfy $a_n^{-1}ha_n \rightarrow 1_G$ in G . If $y, z \in B$ are such that $\varrho(a_n)y \xrightarrow{w} z$ then $\varrho(h)z = z$. In particular, if $\varrho(a_n)z = z$ then $\varrho(h)z = z$.*

Proof. (Strong) continuity of ϱ gives

$$\|\varrho(ha_n)y - \varrho(a_n)y\| = \|\varrho(a_n^{-1}ha_n)y - y\| \rightarrow 0$$

At the same time $\varrho(a_n)y \xrightarrow{w} z$ and $\varrho(ha_n)y \xrightarrow{w} \varrho(h)z$. Hence $\varrho(h)z = z$. \square

We can now prove Theorem 9.1 in the case of $G = \mathbf{SL}_2(k)$ where k is a local field. Assuming that $\varrho : \mathbf{SL}_2(k) \rightarrow \mathbf{O}(B)$ has some matrix coefficient not vanishing at infinity, we get by Lemma 9.2 a sequence $a_n \rightarrow \infty$ in A (here A is the diagonal semigroup $\{\text{diag}(a, 1/a) : a \geq 1\}$), and *non zero* vectors $y, z \in B$ with $\varrho(a_n)y \xrightarrow{w} z$.

Let H denote the upper triangular unipotent subgroup (the horocyclic subgroup). It is normalized by a_n and $a_n^{-1}ha_n \rightarrow e$ for all $h \in H$, so by Lemma 9.3 z is a (non-trivial) $\varrho(H)$ -invariant vector. We may assume that $\|z\| = 1$. The matrix coefficient $f(g) = \langle \varrho(g)z, z^* \rangle$ is a continuous function on G , which is bi- H -invariant:

$$\begin{aligned} \text{(a)} \quad f(gh) &= \langle \varrho(g)\varrho(h)z, z^* \rangle = \langle \varrho(g)z, z^* \rangle = f(g) \\ \text{(b)} \quad f(hg) &= \langle \varrho(g)z, \varrho^*(h^{-1})z^* \rangle = \langle \varrho(g)z, z^* \rangle = f(g) \end{aligned}$$

for all $g \in G$ and $h \in H$. The proof can be now completed as in the original unitary Howe-Moore Theorem. By (a) f can be viewed as a continuous function f_0 on the punctured plane $G/H = k^2 - \{(0, 0)\}$, and by (b) f_0 is constant on each horizontal line $\ell_s = \{(t, s) : t \in k\}$, $s \neq 0$. By continuity, f_0 is a constant on $\{(t, 0) : t \neq 0\}$. Since $f_0(0, 1) = f(e) = 1$ this constant is 1.

This implies that z is $\varrho(A)$ -invariant because $\langle \varrho(a)z, z^* \rangle = f(a) = f(e) = 1$ whilst z^* attains its norm only on z .

Thus z is fixed by the upper triangular group $AH \subseteq G$ and f descends to a continuous function f_1 on the projective line $\mathbf{P}(k^2) = G/AH$. The H -action on $\mathbf{P}(k^2)$ has a dense orbit. Thus f_1 is constant 1, and so is f :

$$\langle \varrho(g)z, z^* \rangle = f(g) = f(e) = 1 \quad (g \in G)$$

Thus the unit vector z is $\varrho(G)$ -invariant, completing the proof in the case of $G = \mathbf{SL}_2(k)$.

The proof of the unitary Howe-Moore theorem for semisimple Lie group $G = \prod G_i$ (c.f. Zimmer [Z], Margulis [M5]) relies only on the reduction to the Cartan

subgroup (Lemma 9.2), the structure of such groups, the case of $\mathbf{SL}_2(k)$ and on Mautner Lemma. Thus the “unitary” argument can be applied almost *verbatim* to the present setup of ucus Banach spaces. \square

REFERENCES

- [A] R. Alperin, Locally compact groups acting on trees and property T . *Monatsh. Math.* **93** (1982), no. 4, 261–265.
- [Ba] S. Banach, *Théorie des opérations linéaires*, Warsaw, 1932.
- [Be] M.E.B. Bekka, *On uniqueness of invariant means*, Proc. Amer. Math. Soc. **126** (1998), no. 2, 507–514.
- [BV] M.E.B. Bekka and A. Valette, Kazhdan’s property (T) and amenable representations, *Math. Z.* **212** (1993), no. 2, 293–299.
- [BHV] M.E.B. Bekka, P. de la Harpe and A. Valette, *Kazhdan’s Property (T)*, forthcoming.
- [BL] Y. Benyamini and J. Lindenstrauss, *Geometric nonlinear functional analysis. Vol. 1*. American Mathematical Society Colloquium Publications, **48**. American Mathematical Society, Providence, RI, 2000.
- [BK] I.N. Bernstein and D.A. Kazhdan, *The one-dimensional cohomology of discrete subgroups* Funkcional. Anal. i Priložen. 4 1970 no. 1, 1–5.
- [B1] N. Bourbaki, *Éléments de mathématique. Fasc. X. Première partie. Livre III: Topologie générale. Chapitre 10: Espaces fonctionnels (deuxième édition, entièrement refondue)*, Actualités Sci. Indust., No. 1084. Hermann, Paris, 1961.
- [B2] N. Bourbaki, *Éléments de mathématique, Fascicule XXIX. Livre IV, Intégration, Chapitres 7 et 8*, Hermann (Act. Sc. Ind. 1306), Paris, 1963.
- [BP] M. Bourdon and H. Pajot, Cohomologie l_p et espaces de Besov. *J. Reine Angew. Math.*, 558:85–108, 2003.
- [BDCK] J. Bretagnolle, D. Dacunha-Castelle, and J.-L. Krivine, *Lois stables et espaces L^p* , Ann. Inst. H. Poincaré Sect. B (N.S.) **2** (1965/1966), 231–259.
- [BMd] M. Burger and N. Monod, *Continuous bounded cohomology and applications to rigidity theory (with an appendix by M. Burger and A. Iozzi)*, Geom. Funct. Anal. **12** (2002), no. 2, 219–280.
- [BMz] M. Burger and S. Mozes, *Groups acting on trees: from local to global structure*, Inst. Hautes Études Sci. Publ. Math. (2000), no. 92, 113–150 (2001).
- [C] A. Connes, A factor of type II_1 with countable fundamental group, *J. Operator Theory* **4** (1980), no. 1, 151–153.
- [CW] A. Connes and B. Weiss, Property T and asymptotically invariant sequences, Israel J. Math. **37** (1980), no. 3, 209–210.
- [D] P. Delorme, 1-cohomologie des représentations unitaires des groupes de Lie semisimples et résolubles. Produits tensoriels continus et représentations, *Bull. Soc. Math. France* **105** (1977), 281–336.
- [DU] J. Diestel and J.J. Uhl, Jr. *Vector measures*. Mathematical Surveys, No. 15. American Mathematical Society, Providence, R.I., 1977.
- [FM1] D. Fisher and G.A. Margulis, Almost isometric actions, property (T) and local rigidity, *preprint* (arXiv:math.DS/0312386).
- [FM2] D. Fisher and G. A. Margulis, Local rigidity of affine actions of higher rank groups and lattices, *preprint* (arXiv:math.DS/0408213).
- [FJ] R.J. Fleming and J.E. Jamison, *Isometries on Banach spaces: function spaces* Monographs and Surveys in Pure and Applied Mathematics, 129. Chapman and HALL/CRC, 2003.
- [GW] E. Glasner and B. Weiss, Kazhdan’s property T and the geometry of the collection of invariant measures, *GFAA, Geom. Func. Anal.* **7** (1997), 917–935.
- [Gr1] M. Gromov, *Hyperbolic groups. Essays in group theory*, 75–263, Math. Sci. Res. Inst. Publ., 8, Springer, New York, 1987.

- [Gr2] M. Gromov, Random walk in random groups, *Geom. Funct. Anal.* **13** (2003), no. 1, 73–146.
- [Gu] A. Guichardet, Sur la cohomologie des groupes topologiques II. *Bull. Soc. Math.* **96** (1972) 305–332.
- [Ha] C.D. Hardin, Isometries on subspaces of L_p , *Indiana Univ. Math. J.* **30** (1981), 449–465.
- [HV] P. de la Harpe and A. Valette, *La propriété (T) de Kazhdan pour les groupes localement compacts* (avec un appendice de Marc Burger), Astérisque No. **175** (1989), 158 pp.
- [Hj] G. Hjorth, A converse to Dye's Theorem., to appear in *Trans. AMS* (2005).
- [KK] S. Kakutani and K. Kodaira, *Über das Haarsche Maß in der lokal bikompakten Gruppe*, Proc. Imp. Acad. Tokyo **20** (1944), 444–450.
- [Ka] D.A. Kazhdan, Connection of the dual space of a group with the structure of its closed subgroups, *Functional Analysis and Application* **1** (1967), 63–65.
- [Kl] V. Klee, *Circumspheres and inner products*, *Math. Scand.* **8** (1960), 363–370.
- [Lg] V. Lafforgue, unpublished.
- [Li] J. Lamperti, On the isometries of some function spaces, *Pacific J. Math.* **8** (1958), 459–466.
- [LT] J. Lindenstrauss and L. Tzafriri, Classical Banach spaces I and II, *Classics in Mathematics*, Springer-Verlag.
- [Lu] A. Lubotzky, *Discrete groups, expanding graphs and invariant measures*, (With an appendix by Jonathan D. Rogawski), Progress in Mathematics, **125**. Birkhäuser Verlag, Basel, 1994.
- [M1] G.A. Margulis, Explicit constructions of expanders, *Problemy Peredači Informacii* **9** (1973), no. 4, 71–80.
- [M2] G.A. Margulis, Finiteness of quotient groups of discrete subgroups, *Funktsional. Anal. i Prilozhen.* **13** (1979), no. 3, 28–39.
- [M3] G. Margulis, *Some remarks on invariant means*, *Monatshafte für Mathematik*, **90** (1980), 233–235.
- [M4] G.A. Margulis, On the decomposition of discrete subgroups into amalgams. Selected translations, *Selecta Math. Soviet.* **1** (1981), no. 2, 197–213.
- [M5] G.A. Margulis, *Discrete subgroups of semisimple Lie groups*, Springer-Verlag, 1991.
- [Mi] I. Mineyev, Straightening and bounded cohomology of hyperbolic groups, *GAFSA* **11** 807–839 (2001).
- [Mo] N. Monod, Superrigidity for irreducible lattices and geometric splitting, *preprint* 2004.
- [MS] N. Monod and Y. Shalom, *Cocycle superrigidity and bounded cohomology for negatively curved spaces*, *J. Differential Geometry* **67** (2004), 395–455.
- [N] A. Navas, Actions de groupes de Kazhdan sur le cercle, *Ann. Sci. École Norm. Sup. (4)* **35** (2002), no. 5, 749–758.
- [Pa] P. Pansu, Cohomologie L^p : invariance sous quasiisométrie, *preprint* (1995), 23 pages.
- [Po] S. Popa, Strong rigidity of II_1 factors arising from malleable actions of w-rigid groups, *preprint* (2004).
- [R] B. Rémy, *Integrability of induction cocycles for Kac-Moody groups*, Preprint, 2004.
- [RS] G. Robertson and T. Steger, Negative definite kernels and a dynamical characterization of property (T) for countable groups, *Ergodic Theory Dynam. Systems* **18** (1998), no. 1, 247–253.
- [Se] J.-P. Serre, *Trees*, Springer-Verlag, Berlin-New York, 1980.
- [Sh] Y. Shalom, *Rigidity of commensurators and irreducible lattices*, *Invent. Math.* **141** (2000), no. 1, 1–54.
- [Su] D. Sullivan, *For $n > 3$ there is only one finitely additive rotationally invariant measure on the n -sphere on all Lebesgue measurable sets*, *Bull. AMS* **4** (1981), 121–123.
- [Wa] Y. Watatani, Property T of Kazhdan implies property FA of Serre, *Math. Japon.* **27** (1982), no. 1, 97–103.

- [WW] J.H. Wells and L.R. Williams, *Embeddings and Extensions in Analysis*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 84. Springer-Verlag, New York-Heidelberg, 1975.
- [Y] G. Yu, Hyperbolic groups admit proper affine isometric actions on l^p -spaces, *preprint* (arXiv:math.GR/0411234).
- [Z] R.J. Zimmer, *Ergodic theory and semisimple groups*, Birkhäuser Verlag, Basel, 1984.
- [Ž] A. Žuk, Property (T) and Kazhdan constants for discrete groups, *Geom. Funct. Anal.* (GAFA) **13** (2003), no. 3, 643–670.

U.B. TECHNION – ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA, ISRAEL

A.F. UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, USA

T.G. YALE UNIVERSITY, NEW HAVEN, USA

N.M. UNIVERSITY OF CHICAGO, CHICAGO, USA