

# Extending the reach of the point-to-set principle


Jack Lutz, Neil Lutz, **Elvira Mayordomo**

Iowa State University, Universidad de Zaragoza

Midwest Computability Seminar, November 22nd 2021

# Fractal dimensions

Given a (separable) metric space, Hausdorff dimension and packing dimension generalize the usual integer dimension idea



# Hausdorff definition of dimension

Let  $\rho$  be a metric on a set  $X$ .

- For  $E \subseteq X$  and  $\delta > 0$ , a  $\delta$ -cover of  $E$  is a collection  $\mathcal{U}$  such that for all  $U \in \mathcal{U}$ ,  $\text{diam}(U) < \delta$  and

$$E \subseteq \bigcup_{U \in \mathcal{U}} U.$$

- For  $s \geq 0$ ,

$$H^s(E) = \lim_{\delta \rightarrow 0} \inf_{\mathcal{U} \text{ is a } \delta\text{-cover of } E} \sum_{U \in \mathcal{U}} \text{diam}(U)^s$$

The Hausdorff dimension of  $E \subseteq X$  is

$$\dim_{\text{H}}(E) = \inf \{s \mid H^s(E) = 0\}.$$

# Gauged dimension

We can avoid infinite dimension by changing the scale /gauge function families

- A **gauge function** is a continuous, nondecreasing function from  $[0, \infty)$  to  $[0, \infty)$  that vanishes only at 0.
- A **gauge family** is a one-parameter family  $\varphi = \{\varphi_s \mid s \in (0, \infty)\}$  of gauge functions  $\varphi_s$  satisfying for  $s > t$ ,  $\varphi_s(\delta) = o(\varphi_t(\delta))$  as  $\delta \rightarrow 0^+$

Definition

$$H^{s,\varphi}(E) = \lim_{\delta \rightarrow 0} \inf_{\mathcal{U} \text{ is a } \delta\text{-cover of } E} \sum_{U \in \mathcal{U}} \varphi_s(\text{diam}(U))$$

$$\dim^\varphi(E) = \inf \{s \mid H^{s,\varphi}(E) = 0\}.$$

They generalize  $\theta_s(\delta) = \delta^s$  in Hausdorff dimension.

# Packing dimension

Let  $\rho$  be a metric on a set  $X$ .

- For  $E \subseteq X$  and  $\delta > 0$ , a  $\delta$ -packing of  $E$  is a collection  $\mathcal{U}$  of disjoint open balls  $U$  with centers in  $E$  and  $\text{diam}(U) < \delta$ .

- For  $s \geq 0$ ,

$$P_0^s(E) = \lim_{\delta \rightarrow 0} \sup_{\mathcal{U} \text{ is a } \delta\text{-packing of } E} \sum_{U \in \mathcal{U}} \text{diam}(U)^s$$

- For  $s \geq 0$ ,

$$P^s(E) = \inf \left\{ \sum_i P_0^s(E_i) \mid E \subseteq \cup E_i \right\}$$

The Packing dimension of  $E \subseteq X$  is

$$\dim_{\text{P}}(E) = \inf \{s \mid P^s(E) = 0\}.$$

# Packing dimension

Let  $\rho$  be a metric on a set  $X$ .

- For  $E \subseteq X$  and  $\delta > 0$ , a  $\delta$ -packing of  $E$  is a collection  $\mathcal{U}$  of disjoint open balls  $U$  with centers in  $E$  and  $\text{diam}(U) < \delta$ .

- For  $s \geq 0$ ,

$$P_0^s(E) = \lim_{\delta \rightarrow 0} \sup_{\mathcal{U} \text{ is a } \delta\text{-packing of } E} \sum_{U \in \mathcal{U}} \text{diam}(U)^s$$

- For  $s \geq 0$ ,

$$P^s(E) = \inf \left\{ \sum_i P_0^s(E_i) \mid E \subseteq \cup E_i \right\}$$

The Packing dimension of  $E \subseteq X$  is

$$\dim_{\text{P}}(E) = \inf \{s \mid P^s(E) = 0\}.$$

Can also be generalized using gauge functions

# Algorithmic dimensions on $2^\omega$

- Effectivizing the (several equivalent) definitions of fractal dimension we obtain algorithmic dimensions on Cantor and Euclidean spaces
- We can effectivize in several ways:
  - From a martingale/gale definition of dimension: an  $s$ -gale is  $d : 2^{<\omega} \rightarrow [0, \infty)$  with

$$d(w) = \frac{d(w0) + d(w1)}{2^s},$$

$$S^\infty[d] = \left\{ x \in 2^\omega \mid \limsup_n (x \upharpoonright n) = \infty \right\}.$$

$$S_{\text{strong}}^\infty[d] = \left\{ x \in 2^\omega \mid \liminf_n (x \upharpoonright n) = \infty \right\}.$$

- From a compression/decompression definition: Fix  $U$  a UTM. Let  $w \in 2^{<\omega}$ ,  $x \in 2^\omega$ ,  $\delta > 0$

$$K(w) = \min \{ |y| \mid U(y) = w \}$$

$$K_\delta(x) = \inf \{ K(q) \mid q \in \mathbb{Q}, |x - q| < \delta \}$$

# Effective dimension

## Definition

Let  $x \in 2^\omega$  and  $A \subseteq 2^\omega$

$$\dim(x) = \liminf_{\delta \rightarrow 0^+} \frac{K_\delta(x)}{\log(1/\delta)}.$$

$$\text{Dim}(x) = \limsup_{\delta \rightarrow 0^+} \frac{K_\delta(x)}{\log(1/\delta)}.$$

They are equivalent to:

$$\dim(x) = \inf \{s \mid \text{there is a lower semicomputable } s\text{-gale } d \text{ with } x \in S^\infty[d]\}$$

(an similarly for  $\text{Dim}$  and  $S_{\text{strong}}$ ).

All definitions relativize to any oracle  $B \subseteq \mathbb{N}$ .



# Effective dimension

These effectivizations are **pointwise**

- $\dim(A) = \sup_{x \in A} \dim(x)$
- $\text{Dim}(A) = \sup_{x \in A} \text{Dim}(x)$

# Effective dimension

## Definition

The  $\varphi$ -gauged algorithmic dimension of a point  $x \in 2^\omega$  is

$$\dim^\varphi(x) = \inf \left\{ s \mid \liminf_{\delta \rightarrow 0^+} 2^{K_\delta(x)} \varphi_s(\delta) = 0 \right\}.$$

# What does it have to do with fractal geometry?

$\dim_{\text{H}}(A)$  is the **Hausdorff dimension** of set  $A$

$\dim_{\text{P}}(A)$  is the **packing dimension** of set  $A$

For  $A \subseteq 2^{\omega}$

$$\dim_{\text{H}}(A) \leq \dim(A)$$

$$\dim_{\text{P}}(A) \leq \text{Dim}(A)$$

# What does it have to do with fractal geometry?

$$\dim_{\text{H}}(A) \leq \dim(A)$$

$$\dim_{\text{P}}(A) \leq \text{Dim}(A)$$

- Useful for upper bounds
- For sets of low complexity,  $\dim_{\text{H}}(A) = \dim(A)$  and  $\dim_{\text{P}}(A) \leq \text{Dim}(A)$  (correspondence principles)
- Partial randomness of points, quantitative analysis of sets ...

# Point-to-set principles

Theorem (Lutz Lutz 2018)

Let  $A \subseteq 2^\omega$ . Then

$$\dim_H(A) = \min_{B \subseteq \mathbb{N}} \dim^B(A).$$

Theorem (Lutz Lutz 2018)

Let  $A \subseteq 2^\omega$ . Then

$$\dim_P(A) = \min_{B \subseteq \mathbb{N}} \text{Dim}^B(A).$$

# Application of point to set principles to fractal geometry: projection formula

Theorem (Marstrand 1954)

*Let  $E \subseteq \mathbb{R}^2$  be an analytic set with  $\dim_{\text{H}}(E) = s$ . Then for almost every  $\theta \in (0, 2\pi)$ ,  $\dim_{\text{H}}(p_{\theta}E) = \min\{s, 1\}$*

# Application of point to set principles to fractal geometry: projection formula

Theorem (Marstrand 1954)

*Let  $E \subseteq \mathbb{R}^2$  be an analytic set with  $\dim_{\text{H}}(E) = s$ . Then for almost every  $\theta \in (0, 2\pi)$ ,  $\dim_{\text{H}}(p_{\theta}E) = \min\{s, 1\}$*

It does not hold for arbitrary  $E$  (assuming CH). Recently an extension using PSP

Theorem (N.Lutz Stull 2018)

*Let  $E \subseteq \mathbb{R}^2$  be an arbitrary set with  $\dim_{\text{H}}(E) = \dim_{\text{P}}(E) = s$ . Then for almost every  $\theta \in (0, 2\pi)$ ,  $\dim_{\text{H}}(p_{\theta}E) = \min\{s, 1\}$*

# Application of point to set principles to fractal geometry: projection formula

Theorem (Marstrand 1954)

*Let  $E \subseteq \mathbb{R}^2$  be an analytic set with  $\dim_{\text{H}}(E) = s$ . Then for almost every  $\theta \in (0, 2\pi)$ ,  $\dim_{\text{H}}(p_{\theta}E) = \min\{s, 1\}$*

It does not hold for arbitrary  $E$  (assuming CH). Recently an extension using PSP

Theorem (N.Lutz Stull 2018)

*Let  $E \subseteq \mathbb{R}^2$  be an arbitrary set with  $\dim_{\text{H}}(E) = \dim_{\text{P}}(E) = s$ . Then for almost every  $\theta \in (0, 2\pi)$ ,  $\dim_{\text{H}}(p_{\theta}E) = \min\{s, 1\}$*

Further extension in (Stull 2021)



# Intersection formula

Theorem (Kahane 1986, Mattila 1984)

Let  $E, F \subseteq \mathbb{R}^n$  be **Borel sets**. Then for almost every  $z \in \mathbb{R}^n$ ,

$$\dim_{\mathbb{H}}(E \cap (F + z)) \leq \max\{0, \dim_{\mathbb{H}}(E \times F) - n\}$$

where  $F + z = \{x + z \mid x \in F\}$ .

# Intersection formula

Theorem (Kahane 1986, Mattila 1984)

Let  $E, F \subseteq \mathbb{R}^n$  be **Borel sets**. Then for almost every  $z \in \mathbb{R}^n$ ,

$$\dim_{\mathbb{H}}(E \cap (F + z)) \leq \max\{0, \dim_{\mathbb{H}}(E \times F) - n\}$$

where  $F + z = \{x + z \mid x \in F\}$ .

Theorem (N.Lutz 2021)

Let  $E, F \subseteq \mathbb{R}^n$ . Then for almost every  $z \in \mathbb{R}^n$ ,

$$\dim_{\mathbb{H}}(E \cap (F + z)) \leq \max\{0, \dim_{\mathbb{H}}(E \times F) - n\}$$

where  $F + z = \{x + z \mid x \in F\}$ .

- (Lutz Stull 2020) results on Furstenberg sets
- (Slaman 2021) The Hausdorff dimensions of co-analytic sets are not carried by their closed subsets
- (Lutz 2021) There are Hamel bases ( $\mathbb{R}$  over  $\mathbb{Q}$ ) with any positive Hausdorff dimension

# Looking at other separable spaces

- Where can we effectivize dimension?
  - We can define Kolmogorov complexity/ effectivize Hausdorff measure if we have a **separator** (countable dense set)

# Looking at other separable spaces

- Where can we effectivize dimension?
  - We can define Kolmogorov complexity/ effectivize Hausdorff measure if we have a **separator** (countable dense set)

Definition (Kolmogorov complexity of  $x$  at precision  $\delta$ )

Let  $(X, \rho)$  be a separable metric space and let  $D \subseteq X$  be a countable dense set (fix  $f : 2^{<\omega} \rightarrow D$ )

$$K_\delta(x) = \inf \{K(w) \mid w \in 2^{<\omega}, \rho(x, f(w)) < \delta\}$$

# Looking at other separable spaces

## Definition

The *algorithmic dimension* and strong algorithmic dimension of a point  $x \in X$  is

$$\dim(x) = \liminf_{\delta \rightarrow 0^+} \frac{K_\delta(x)}{\log(1/\delta)},$$

$$\text{Dim}(x) = \limsup_{\delta \rightarrow 0^+} \frac{K_\delta(x)}{\log(1/\delta)}.$$

# Looking at other spaces: gauge dimension

## Definition

The  $\varphi$ -gauged algorithmic dimension and strong algorithmic dimension of a point  $x \in X$  is

$$\dim^\varphi(x) = \inf \left\{ s \mid \liminf_{\delta \rightarrow 0^+} 2^{K_\delta(x)} \varphi_s(\delta) = 0 \right\},$$

and the  $\varphi$ -gauged of  $x$  is

$$\text{Dim}^\varphi(x) = \inf \left\{ s \mid \limsup_{\delta \rightarrow 0^+} 2^{K_\delta(x)} \varphi_s(\delta) = 0 \right\},$$

# Looking at other spaces: gauge dimension

## Definition

The  $\varphi$ -gauged algorithmic dimension and strong algorithmic dimension of a point  $x \in X$  is

$$\dim^\varphi(x) = \inf \left\{ s \mid \liminf_{\delta \rightarrow 0^+} 2^{K_\delta(x)} \varphi_s(\delta) = 0 \right\},$$

and the  $\varphi$ -gauged of  $x$  is

$$\text{Dim}^\varphi(x) = \inf \left\{ s \mid \limsup_{\delta \rightarrow 0^+} 2^{K_\delta(x)} \varphi_s(\delta) = 0 \right\},$$

Ordinary Hausdorff/packing dimensions use the *canonical gauge family* defined by  $\theta_s(\delta) = \delta^s$



# A couple of restrictions on gauge families

$\varphi = \{\varphi_s \mid s \in (0, \infty)\}$  is a **strong gauge family** if the following hold:

- There is a precision family: a one-parameter family  $\alpha = \{\alpha_s \mid s \in (0, \infty)\}$  of functions  $\alpha_s : \mathbb{N} \rightarrow \mathbb{Q}^+$  that vanish as  $r \rightarrow \infty$  and satisfy
  - $\varphi_s(\alpha_s(r)) = O(\varphi_s(\alpha_s(r+1)))$  as  $r \rightarrow \infty$
  - $\sum_{r \in \mathbb{N}} \frac{\varphi_t(\alpha_s(r))}{\varphi_s(\alpha_s(r))} < \infty$  whenever  $s < t$ .
- $\varphi_t(2\delta) = O(\varphi_s(\delta))$  for all  $s \leq t$
- $\varphi_s(\delta) = O(1/\log \log(1/\delta))$  as  $\delta \rightarrow 0^+$

# General Point-to-set principles

Let  $(X, \rho)$  be a separable metric space,  $\varphi$  a strong gauge family

Theorem (Lutz Lutz M 2021)

Let  $A \subseteq X$ . Then

$$\dim_{\text{H}}^{\varphi}(A) = \min_{B \subseteq \mathbb{N}} \sup_{x \in A} \dim^{\varphi, B}(x).$$

Theorem (Lutz Lutz M 2021)

Let  $A \subseteq X$ . Then

$$\dim_{\text{P}}^{\varphi}(A) = \min_{B \subseteq \mathbb{N}} \sup_{x \in A} \text{Dim}^{\varphi, B}(x).$$

# The hyperspace

- Let  $(X, \rho)$  be a separable metric space
- Let  $\mathcal{K}(X)$  be the set of nonempty compact subsets of  $X$  together with the Hausdorff metric  $\text{dist}_H$  defined as follows

$$\text{dist}_H(U, V) = \max \left\{ \sup_{x \in U} \rho(x, V), \sup_{y \in V} \rho(y, U) \right\}.$$

$$(\rho(a, B) = \inf \{ \rho(a, b) \mid b \in B \})$$

# Relationship of the dimensions of $E$ and $\mathcal{K}(E)$

McClure (1995 and 1996) has several results relating Hausdorff and packing dimensions of a set  $E$  and  $\mathcal{K}(E)$  for

- $E$  self-similar
- $E$   $\sigma$ -compact

# Relationship of the dimensions of $E$ and $\mathcal{K}(E)$

McClure (1995 and 1996) has several results relating Hausdorff and packing dimensions of a set  $E$  and  $\mathcal{K}(E)$  for

- $E$  self-similar
- $E$   $\sigma$ -compact

$\mathcal{K}(E)$  has infinite dimension, a different gauge family is needed

# The jump of a gauge family

## Definition

The *jump* of a gauge family  $\varphi$  is the family  $\tilde{\varphi}$  given  $\tilde{\varphi}_s(\delta) = 2^{-1/\varphi_s(\delta)}$ .

For the canonical gauge family  $\theta_s(\delta) = \delta^s$ ,  $\tilde{\theta}_s(t) = 2^{-1/t^s}$ .

Theorem (McClure 1995)

Let  $E \subseteq X$  be  $\sigma$ -**compact**. Let  $\psi_s(t) = 2^{-1/t^s}$ . Then

$$\dim_{\mathbb{P}}^{\psi}(\mathcal{K}(E)) \geq \dim_{\mathbb{P}}(E).$$

# The jump of a gauge family

## Definition

The *jump* of a gauge family  $\varphi$  is the family  $\tilde{\varphi}$  given  $\tilde{\varphi}_s(\delta) = 2^{-1/\varphi_s(\delta)}$ .

For the canonical gauge family  $\theta_s(\delta) = \delta^s$ ,  $\tilde{\theta}_s(t) = 2^{-1/t^s}$ .

Theorem (McClure 1995)

Let  $E \subseteq X$  be  $\sigma$ -**compact**. Let  $\psi_s(t) = 2^{-1/t^s}$ . Then

$$\dim_{\mathbb{P}}^{\psi}(\mathcal{K}(E)) \geq \dim_{\mathbb{P}}(E).$$

We aim to extend the theorem to other  $E$  and to other gauge families beside the canonical one.

# Hyperspace packing dimension theorem

Theorem (LLM 2021)

*Let  $E \subseteq X$  be an analytic set, and let  $\varphi$  be strong gauge family, then*

$$\dim_{\mathbb{P}}^{\tilde{\varphi}}(\mathcal{K}(E)) \geq \dim_{\mathbb{P}}^{\varphi}(E).$$



# Where we use PSP in the hyperspace packing dimension theorem

- By the general point-to-set principle, let  $A$  be an oracle such that

$$\dim_{\mathbb{P}}^{\tilde{\varphi}}(\mathcal{K}(E)) = \sup_{L \in \mathcal{K}(E)} \text{Dim}^{\tilde{\varphi}, A}(L),$$

# Where we use PSP in the hyperspace packing dimension theorem

- By the general point-to-set principle, let  $A$  be an oracle such that

$$\dim_{\mathbb{P}}^{\tilde{\varphi}}(\mathcal{K}(E)) = \sup_{L \in \mathcal{K}(E)} \text{Dim}^{\tilde{\varphi}, A}(L),$$

- We recursively construct a single compact set  $L \in \mathcal{K}(E)$  (i.e., a single point in the hyperspace  $\mathcal{K}(E)$ ) that has high Kolmogorov complexity at infinitely many precisions, relative to oracle  $A$ .

$$\text{Dim}^{\tilde{\varphi}, A}(L) > s$$

# Where we use PSP in the hyperspace packing dimension theorem

- By the general point-to-set principle, let  $A$  be an oracle such that

$$\dim_{\mathbb{P}}^{\tilde{\varphi}}(\mathcal{K}(E)) = \sup_{L \in \mathcal{K}(E)} \text{Dim}^{\tilde{\varphi}, A}(L),$$

- We recursively construct a single compact set  $L \in \mathcal{K}(E)$  (i.e., a single point in the hyperspace  $\mathcal{K}(E)$ ) that has high Kolmogorov complexity at infinitely many precisions, relative to oracle  $A$ .

$$\text{Dim}^{\tilde{\varphi}, A}(L) > s$$

- For  $E$  compact, we can reach  $\text{Dim}^{\tilde{\varphi}, A}(L) > s$  for  $s = \dim_{\mathbb{P}}^{\varphi}(E)$

# Open questions

- Is there a more general hyperspace Hausdorff dimension theorem?  $\dim_{\mathbb{H}}^{\tilde{\varphi}}(\mathcal{K}(E))$  vs  $\dim_{\mathbb{H}}^{\varphi}(E)$  for interesting  $E$
- Are (the complexity or the good properties of) the two oracles in the PTSP related to hyperspace dimension theorems?

# References

- Jack H. Lutz and Neil Lutz, Who asked us? How the theory of computing answers questions about analysis, Ding-Zhu Du and Jie Wang (eds.), Complexity and Approximation: In Memory of Ker-I Ko, pp. 48-56, Springer, 2020.
- Jack H. Lutz and Elvira Mayordomo, Algorithmic fractal dimensions in geometric measure theory. In Vasco Brattka and Peter Hertling (eds.), Handbook of Computability and Complexity in Analysis, Springer-Verlag (2021).
- Jack H. Lutz, Neil Lutz, and Elvira Mayordomo, Extending the Reach of the Point-to-Set Principle.  
<https://arxiv.org/abs/2004.07798>