

BLOCH–KATO CONJECTURES FOR AUTOMORPHIC MOTIVES

FRANK CALEGARI, DAVID GERAGHTY, AND MICHAEL HARRIS

CONTENTS

1. Introduction	1
1.1. Acknowledgements	3
2. Relation with special values of periods	3
3. Vanishing Theorems	4
3.1. Betti Cohomology	4
3.2. Coherent Cohomology	6
4. Proofs	7
References	9

1. INTRODUCTION

Assume (for this paragraph only) the standard conjectures, and suppose that M is a pure irreducible Grothendieck motive over \mathbf{Q} with coefficients in (say) a totally real field E . We make no assumption on the regularity or self-duality of M . According to conjectures of Hasse–Weil, Langlands, Clozel, and others, one expects that the motive M is automorphic, and corresponds to an algebraic cuspidal automorphic representation π for $\mathrm{GL}(n)/\mathbf{Q}$ such that $L(\pi, s) = L(M, s)$. By a theorem of Jacquet and Shalika [JS81], the L -function

$$L(M \times M^\vee, s) = L(\pi \times \pi^\vee, s)$$

is meromorphic for $\mathrm{Re}(s) > 0$ and has a simple pole at $s = 1$. Let $\mathrm{ad}^0(M)$ be the pure motive of weight zero with coefficients in E such that $\mathrm{ad}^0(M) \oplus E = M \times M^\vee$. Then

$$L(\mathrm{ad}^0(M), s) = \frac{L(\pi \times \pi^\vee, s)}{\zeta(s)},$$

and $L(\mathrm{ad}^0(M), 1) \neq 0$ is finite. According to conjectures of Deligne and Bloch–Kato [BK90], for any pure de Rham representation V , there is an equality:

$$\dim H_f^1(G_{\mathbf{Q}}, V) - \dim H^0(G_{\mathbf{Q}}, V) = \mathrm{ord}_{s=1} L(V^*, s).$$

In particular, if we take $V = V^* = \mathrm{ad}^0(M)$, then we expect that $H_f^1(G_{\mathbf{Q}}, \mathrm{ad}^0(M))$ should vanish. This is a special case of the more general fact that $H_f^1(\mathbf{Q}, V)$ should be trivial for any p -adic representation V arising from a pure motive M of weight $w \geq 0$. One also conjectures

2010 *Mathematics Subject Classification.* 11F33, 11F80.

The first author was supported in part by NSF Grant DMS-1404620. The second author was supported in part by NSF Grants DMS-1200304 and DMS-1128155. The third author was supported in part by NSF DMS-1404769. M.H.’s research received funding from the European Research Council under the European Community’s Seventh Framework Programme (FP7/2007-2013) / ERC Grant agreement no. 290766 (AAMOT).

that the value of $L(\mathrm{ad}^0(M), 1)$, after normalization by some suitable period should lie in \mathbf{Q}^\times . Moreover, after equating M with its étale realization for some prime p , the normalized L function should have the same valuation as the order of a corresponding Selmer group $H_f^1(\mathbf{Q}, \mathrm{ad}^0(M) \otimes \mathbf{Q}_p/\mathbf{Z}_p)$.

No longer assuming any conjectures, suppose that $M = \{r_\lambda\}$ is now a weakly compatible system of n -dimensional irreducible Galois representations of $G_{\mathbf{Q}}$, and suppose moreover that M is automorphic, that is, it corresponds to a cuspidal form π for $\mathrm{GL}(n)/\mathbf{Q}$ in a manner compatible with the local Langlands correspondence. Then, even without the standard conjectures, it makes sense to ask, for a p -adic representation $r : G_{\mathbf{Q}} \rightarrow \mathrm{GL}_n(\mathcal{O})$ coming from M (for some finite extension K/\mathbf{Q}_p with ring of integers \mathcal{O}), if the Selmer group

$$H_f^1(\mathbf{Q}, \mathrm{ad}^0(r) \otimes K/\mathcal{O})$$

is finite. Theorems of this kind were first proved for $n = 2$ by Flach [Fla92], and they are also closely related to modularity lifting theorems as proved by Wiles [Wil95, TW95], see (in particular) [DFG04]. More precisely, the order of this group is related to the order of a congruence ideal between modular forms. In this paper, we prove versions of these results for modular abelian surfaces and (conditionally) compatible families of n -dimensional representations whose existence was only recently proved to exist [HLTT]. The main theorem is the following.

Theorem 1.1. *Let A/\mathbf{Q} be a semistable modular abelian surface with $\mathrm{End}(A) = \mathbf{Z}$. Let p be a prime such that:*

- (1) p is sufficiently large with respect to some constant depending only on A .
- (2) A is ordinary at p , and if α, β are the unit root eigenvalues of $D_{\mathrm{cris}}(V)$, then

$$(\alpha^2 - 1)(\beta^2 - 1)(\alpha - \beta)(\alpha^2\beta^2 - 1) \not\equiv 0 \pmod{p}.$$

Then

$$H_f^1(\mathbf{Q}, \mathrm{asp}^0(r) \otimes \mathbf{Q}_p/\mathbf{Z}_p) = 0$$

where $\mathrm{asp}^0(r)$ is the 10-dimensional adjoint representation of $\mathrm{PGSp}(4)$. Moreover, the set of primes p satisfying these conditions has density one.

For the families of Galois representations constructed in [HLTT], we prove the following result.

Theorem 1.2. *Let π be a weight zero regular algebraic cuspidal representation for $\mathrm{GL}(n)/F$ for a CM field F and coefficients in E . Let λ be a prime of \mathcal{O}_E dividing p , and let*

$$r = r_\lambda(\pi) : G_F \rightarrow \mathrm{GL}_n(\mathcal{O})$$

be a p -adic representation associated to π with determinant $\epsilon^{n(1-n)/2}$. Assume that

- (1) $\bar{r}|_{F(\zeta_p)}$ has enormous image in the sense of [CG18] §9.2,
- (2) Let $v \neq p$ be a prime at which π is ramified.
 - (a) π_v is an unramified twist of the Steinberg representation.
 - (b) The representations $r|_{G_{I_v}}$ and $\bar{r}|_{G_{I_v}}$ are unipotent. For a topological generator $\sigma_v \in I_v$ of tame inertia, $\bar{r}(\sigma)$ consists of a single block, namely:

$$\dim \ker(\bar{r}(\sigma) - \mathrm{id}_n) = 1.$$

- (3) p is sufficiently large with respect to some constant depending only on π .

Assume all of Conjecture B of [CG18] except assumption (4). Then the Selmer group $H_f^1(F, \text{ad}^0(r) \otimes K/\mathcal{O})$ is trivial.

Note that Conjecture B of [CG18] consists of five parts: The first part concern local–global compatibility at $v|p$, which is still open. The second and third parts concern local–global compatibility at finite v . Here there is work in characteristic zero by Varma [Var14], although arguments of this nature should also apply to the Galois representations constructed by Scholze [Sch15], at least for modularity lifting purposes (since for modularity lifting it is usually sufficient to have local–global compatibility up to N -semi-simplification). The fifth part is essentially addressed in [CG18], and also (in a different and arguably superior manner) in [KT17]. Hence the main remaining issue is local–global compatibility at $\ell = p$.

Unlike the case of Theorem 1.1, we do not know whether Theorem 1.2 applies for infinitely many p . One reason is that we do not even know that the representations r_λ are irreducible for sufficiently large p . Another is that we do not know whether r is a minimal deformation of \bar{r} at ramified primes v for sufficiently large p , although this is predicted to hold by some generalization of Serre’s conjecture. One example to which this does apply is to the Galois representations associated to symmetric powers of non-CM elliptic curves E over F . (The conclusion of the theorem holds for F if it holds for any extension F'/F , and any symmetric power of E/F is potentially modular over some CM extension F'/F in this case by [ACC⁺18]. We deduce our theorems from the modularity lifting results of [CG18] and [CG], of which we assume familiarity. One obstruction to directly applying the theorems of [CG18] is that the modularity results of *ibid.* require further unproven assumptions, namely, the vanishing of certain cohomology groups outside a prescribed range. The main observation here is that vanishing in these cases may be established for all sufficiently large p .)

For automorphic representations for $\text{GL}(n)$, we require the extra assumption of local–global compatibility at $v|p$, which is not yet known in full generality. Some results along this lines have very recently been announced in [ACC⁺18], although they are not strong enough to give a completely unconditional proof of Theorem 1.2. One problem is that [ACC⁺18, Theorem 4.5.1] requires the hypothesis that F contains an imaginary quadratic field in which p splits, which has to fail for a set of primes p of positive density. It may be possible to give an unconditional version of Theorem 1.2 under some such assumption on p , although we do not pursue this here, in part because we would still be unable to establish for a general π that condition (2b) holds for infinitely many p . (The nilpotent ideal of [Sch15] and [ACC⁺18] would also be an annoying complication.) Similarly the assumption (2a) that π_v is a twist of Steinberg representation can (in principle) be weakened to the weaker assumption that π_v is of the form $\text{Sp}_{n_1}(\chi_1) \boxplus \text{Sp}_{n_2}(\chi_2) \dots \boxplus \text{Sp}_{n_k}(\chi_k)$ for some partition $n = \sum_{i=1}^k n_i$. The main reason we do *not* do this is that it would require a more precise discussion of local–global compatibility at $v \nmid p$, and in particular a refinement of Conjecture B of [CG18].

1.1. Acknowledgements. The authors thank George Boxer for pointing out an oversight in the first version of this note.

2. RELATION WITH SPECIAL VALUES OF PERIODS

The Bloch–Kato conjecture actually gives a more precise prediction of the exact order of the Selmer group in terms of the value of the L -function divided by a certain motivic period. One can think of this as two separate conjectures. The first is to show that the normalization

of $L(1)$ by a suitable period is indeed rational. The second is to relate the corresponding p -adic valuation of this ratio to the order of a Selmer group. Our method naturally relates the order of a Selmer group to a certain tangent space. On the other hand, for most of the Galois representations we consider, it is not known whether there exists a corresponding motive, and so it is not clear exactly what it means to prove rationality. There are some formulations where one can establish certain forms of rationality (or even integrality) with respect to periods defined in terms of automorphic integrals (see, for example, [BR17], [GHL16], and also [Urb98]). However, it is not clear to the authors how these results exactly relate to the (sometimes conjectural) motivic periods. An interesting test case is the following. Suppose that

$$\rho : G_{\mathbf{Q}} \rightarrow \mathrm{GL}_2(\mathbf{C})$$

is an irreducible odd representation. According to the Artin conjecture (known in this case, see [BT99, Buz03, KW09a, KW09b, Kis09]), one knows that ρ is modular of weight one. If one chooses a prime p , and supposes that ρ has a model over \mathcal{O} , the finiteness of the Selmer group $H_f^1(\mathbf{Q}, \mathrm{ad}^0(\rho) \otimes K/\mathcal{O})$ is a consequence of the finiteness of the p -class group of $\mathbf{Q}(\ker(\rho))$. (The former is a quotient of the latter.) The methods of this paper (following [CG18]) show that, at all primes $p > 2$ such that ρ is unramified, the Selmer group $H_f^1(\mathbf{Q}, \mathrm{ad}^0(\rho) \otimes K/\mathcal{O})$ is detected by congruences between the modular form f and other Katz modular forms of weight one which may not lift to characteristic zero. In particular, there exist such congruences if and only if $H_f^1(\mathbf{Q}, \mathrm{ad}^0(\rho) \otimes K/\mathcal{O})$ is non-zero. However, unlike in the case of higher weight modular forms, there does not seem to be an *a priori* way to relate this to a normalization of the adjoint L function $L(\mathrm{ad}^0(\rho), 1)$ (which in this case is an Artin L -function). The issue is that all such constructions (following Hida [Hid81]) proceed by understanding various pairings on the Betti cohomology of arithmetic groups in characteristic zero, whereas weight one Katz modular forms only have an interpretation in terms of coherent cohomology. Even in cases where one does have access to Betti cohomology, say for regular algebraic cuspidal automorphic representations for $\mathrm{GL}(n)/F$ (even for $\mathrm{GL}(2)$ over imaginary quadratic fields F), it is not so clear whether the cohomological pairings one can define give the “correct” regulators or merely the regulators up to some finite multiple related to the torsion classes in cohomology. Since we have nothing to say about how to resolve these issues, we follow Wittgenstein’s dictum ([Wit21] §7) and say no more about them.

3. VANISHING THEOREMS

The main idea of this paper is to note that the various vanishing theorems which are required inputs for the method of [CG18, CG] may be established at least for p sufficiently large. This is not so useful for applications to modularity — if p is sufficiently large, then any completion of the appropriate Hecke ring \mathbf{T} at a maximal ideal \mathfrak{m} of residue characteristic p will be formally smooth of dimension one, and so the only characteristic zero representation one can prove is modular is the representation one must assume is modular in the first place. However, with respect to Selmer groups, this statement does have content — it says that these representations will have no infinitesimal deformations.

3.1. Betti Cohomology. Let F be an imaginary CM field of degree $2d$. Let

$$\begin{aligned} l_0 &:= d(\mathrm{rank}(\mathrm{SL}_n(\mathbf{C})) - \mathrm{rank}(\mathrm{SU}_n(\mathbf{C}))) = d(n - 1), \\ 2q_0 + l_0 &= d(\mathrm{dim}(\mathrm{SL}_n(\mathbf{C})) - \mathrm{dim}(\mathrm{SU}_n(\mathbf{C}))) = d(n^2 - 1), \end{aligned}$$

$$q_0 = \frac{d(n^2 - n)}{2}.$$

Fix a tempered cuspidal automorphic representation π for $\mathrm{PGL}(n)/F$ of weight zero with coefficients in E . Let $Y = Y(K)$ be the corresponding arithmetic orbifold considered in §9 of [CG18], where K is chosen to be maximal at all unramified primes for π and Iwahori level Iw_v for all ramified primes. Let \mathbf{T} denote the (anemic) Hecke algebra defined as the \mathbf{Z} -subalgebra of

$$\mathrm{End} \bigoplus_{k,m} H^k(Y(K), \mathbf{Z}/m\mathbf{Z})$$

generated by Hecke endomorphisms $T_{\alpha,i}$ for $i \leq n$ and α which are units at primes dividing the level. (cf [CG18, Definition 9.1].) For a prime v of \mathcal{O}_E , let

$$\bar{r}_v : G_F \rightarrow \mathrm{GL}_n(k)$$

be the corresponding semi-simple Galois representation, and let \mathfrak{m} denote the corresponding maximal ideal of \mathbf{T} .

Lemma 3.1. *For all sufficiently large v , and $\mathcal{O} = \mathcal{O}_{E,v}$, we have $H^i(Y, \mathcal{O}/\varpi^k)_{\mathfrak{m}} = 0$ unless $i \in [q_0, \dots, q_0 + l_0]$.*

Proof. Assume otherwise. Pick a neat finite index subgroup $K' \subset K$, and a corresponding Galois cover $Y' = Y(K') \rightarrow Y = Y(K)$ where Y' is now a manifold. It follows that $H^*(Y', \mathbf{Z})$ is finitely generated, and thus $H^*(Y, \mathbf{Z}[1/M])$ is also finitely generated where M denotes the product of primes dividing $[K : K']$. We now assume that v has residue characteristic prime to $[K : K']$. Since $H^*(Y, \mathbf{Z}[1/M])$ is finitely generated, the groups $H^*(Y, \mathcal{O}) = H^*(Y, \mathbf{Z}[1/M]) \otimes \mathcal{O}$ are torsion free and of finite rank over \mathcal{O} for all i when \mathcal{O} has sufficiently large residue characteristic. Moreover, there exist only finitely many systems of eigenvalues which occur in $H^*(Y, \mathbf{R})$. Assuming that the result is false (and there are infinitely many v), we deduce that there exists an eigenclass $[c]$ in $H^i(Y, \mathcal{O}_E)$ with $i \notin [q_0, \dots, q_0 + l_0]$ such that the action of \mathbf{T} on $[c]$ has support at \mathfrak{m} for infinitely many primes v of \mathcal{O}_E . By the Chinese remainder theorem, the Hecke eigenvalues of $[c]$ coincide with those of π . We now show that $[c]$ corresponds to an automorphic form Π which must simultaneously be non-tempered and yet isomorphic to π , giving a contradiction. Eigenclasses in cohomology may be realized by isobaric automorphic representations (see [FS98, Thm 2.3]). Suppose that $[c]$ corresponds to such an automorphic representation Π . Because of the degree where $[c]$ occurs, we deduce (from [BW80, Ch.II, Prop 3.1] and [Clo90, Lemma 3.14]) that Π is not tempered. Yet by strong multiplicity one [JS81], there is an isomorphism $\Pi \simeq \pi$. \square

For a more detailed discussion (in a more general setting) relating the cohomology of local systems to tempered automorphic representations, see the proof of [ACC⁺18, Thm 2.4.9].

Theorem 3.2. *Suppose that $H^i(Y, \mathcal{O}/\varpi^n)_{\mathfrak{m}} = 0$ unless $i \in [q_0, \dots, q_0 + l_0]$. Let Q be a finite collection of primes x such that $\bar{r}(\mathrm{Frob}_x)$ has distinct eigenvalues and $N(x) \equiv 1 \pmod{p}$. Then*

$$H^i(Y_1(Q), \mathcal{O}/\varpi^n)_{\mathfrak{m}_\alpha} = 0$$

for all $i \notin [q_0, \dots, q_0 + l_0]$, where:

- (1) $\alpha = \{\alpha_x\}$ is a choice of eigenvalues of $\bar{r}(\mathrm{Frob}_x)$ for each x dividing Q .
- (2) The localization takes place with respect to the Hecke algebra \mathbf{T}_Q consisting of the Hecke operators prime to p and prime to the level together with $U_x - \alpha_x$ for all $x \in Q$.

In particular, the conclusions of this theorem apply for all sufficiently large p .

Proof. We first note that the assumption (of absolute irreducibility) on \bar{r} ensures that the cohomology of the boundary vanishes after localization at \mathfrak{m} . This is because the cohomology of the boundary may be computed inductively from the cohomology of Levi subgroups and then of GL_{n_i} for $n_i < n$ (see, for example, §3 and in particular Prop. 3.3 of [CLH16]), and so the corresponding Galois representations associated to these classes are reducible.

By Poincaré duality (and the discussion above concerning the vanishing of the boundary cohomology localized at \mathfrak{m}), it suffices to prove the result for $i < q_0$. Let i be the smallest integer for which the inequality is violated. Then, by the Hochschild–Serre spectral sequence, we deduce that

$$H^i(Y_0(Q), k)_{\mathfrak{m}_\alpha} \neq 0.$$

As in §9.4 of [CG18] (see also Lemma 6.25(4) of [KT17]), we deduce that $H^i(Y_0(Q), k)_{\mathfrak{m}_\alpha} \simeq H^i(Y, k)_{\mathfrak{m}}$. The result then follows by Lemma 3.1. \square

The modularity lifting theorems of [CG18] are proved by constructing sets of so-called “Taylor–Wiles primes” which have the property that imposing local conditions at these primes annihilates (as much as possible) the dual Selmer group. The assumption that $\bar{r}_v|_{F(\zeta_p)}$ has enormous image implies that there exists arbitrarily many sets Q of auxiliary Taylor–Wiles primes satisfying the hypothesis that $\bar{r}(\mathrm{Frob}_x)$ has distinct eigenvalues. In particular, Theorem 3.2 serves as a replacement for Conjecture B(4) of [CG18]. (For a different (and somewhat cleaner) treatment of Taylor–Wiles primes using the enormous image hypothesis, see [KT17], which is also used in [ACC⁺18].)

3.2. Coherent Cohomology. Let \mathcal{O} denote the ring of integers in some finite extension of \mathbf{Q}_p . Let X denote a toroidal compactification of a Siegel 3-fold Y of level prime to p over $\mathrm{Spec} \mathcal{O}$, and let Z denote the minimal compactification. Let $\pi : \mathcal{A} \rightarrow Y$ denote the universal abelian variety, let $\mathcal{E} = \pi_* \Omega_{\mathcal{A}/X}^1$, let $\omega = \det \mathcal{E}$, and, by abuse of notation, also let ω denote the canonical extension of ω to X or the corresponding ample line bundle on Z . Fix a cuspidal automorphic representation π for $\mathrm{GSp}(4)/\mathbf{Q}$ corresponding to a modular abelian surface A which we assume has endomorphism ring \mathbf{Z} over \mathbf{Q} , and hence to a cuspidal Siegel modular form of scalar weight 2. Let

$$\bar{r}_p : G_{\mathbf{Q}} \rightarrow \mathrm{GSp}_4(\mathbf{F}_p)$$

be the corresponding semi-simple representation for each prime p . Let \mathfrak{m} denote the corresponding maximal ideal of \mathbf{T} .

Lemma 3.3. *For all sufficiently large p , and any set Q of auxiliary primes, we have*

$$H^i(X, \omega_{\mathcal{O}/\varpi^n}^2)_{\mathfrak{m}_Q} = 0$$

for $i = 2$ and 3 , where \mathfrak{m}_Q denotes the maximal ideal in the Hecke algebra where operators at Q and p have been omitted.

Proof. This is a consequence of the proof of Theorem 7.11 of [CG]. We give a brief sketch here of the idea: as in the proof of Lemma 3.1, we otherwise deduce that there exists a characteristic zero form in $H^i(X, \omega_{\mathbf{C}}^2)$ giving rise to infinitely many of these classes. The representation \bar{r}_p will be irreducible for all sufficiently large p (because $\mathrm{End}_{\mathbf{Q}}(A) = \mathbf{Z}$ — see also the proof of Lemma 4.1). It follows that the transfer of this form to $\mathrm{GL}(4)$ must be cuspidal, and moreover (by multiplicity one) coincide with the transfer of the representation coming from the holomorphic Siegel modular form. But such a representation only contributes to cohomology

in degrees 0 and 1. (For details relating the coherent cohomology of Siegel threefolds and their relation to automorphic representations we refer the reader back to [CG].) \square

Lemma 3.4. *Suppose that $H^2(X, \omega_{\mathcal{O}/\varpi^n}^2)_{\mathfrak{m}_Q} = 0$ for any set of auxiliary primes Q , as in the statement of Lemma 3.3. Suppose, moreover, that \bar{r}_p is absolutely irreducible. Then, for $i \geq 2$,*

$$H^i(X_1(Q), \omega_{\mathcal{O}/\varpi^n}^2)_{\mathfrak{m}_\alpha} = H^i(X_0(Q), \omega_{\mathcal{O}/\varpi^n}^2)_{\mathfrak{m}_\alpha} = 0,$$

where Q is any collection of primes where $\bar{r}(\text{Frob}_x)$ has distinct eigenvalues, $N(x) \equiv 1 \pmod{p}$, and $\alpha = \{\alpha_x\}$ is any collection of eigenvalues of $\rho(\text{Frob}_x)$ for x dividing Q . In particular, the conclusions of this theorem apply for all sufficiently large p .

Proof. In [CG], a somewhat elaborate version of this result is proved in Lemmas 7.4 and 7.5. We instead, however, use the modified treatment of Taylor–Wiles primes by Khare and Thorne (cf. [KT17, Lem. 6.25]), as adapted for $\text{GSp}(4)$ in §2.4 and §7.9 of [BCGP18], which leads to a great simplification. By dévissage, we can reduce to the case where the coefficients are a finite field k . To deduce vanishing for $X_1(Q)$, we first use Serre duality to reduce the problem to vanishing of $H^i(X_1(Q), \omega(-\infty))_{\mathfrak{m}_\alpha^*}$ for $i \leq 1$. (Serre duality is only Hecke equivariant up to a twist by a power of the cyclotomic character, which is recorded by the star.) Equivalently, it suffices to show that, if i denotes the smallest integer such that $H^i(X_1(Q), \omega(-\infty))_{\mathfrak{m}_\alpha^*}$ is non-zero, then $H^i(X, \omega(-\infty))_{\mathfrak{m}_Q}$ is also non-zero. By Hochschild–Serre applied to the map $X_1(Q) \rightarrow X_0(Q)$, it suffices to show that, if i denotes the smallest integer such that $H^i(X_0(Q), \omega(-\infty))_{\mathfrak{m}_\alpha^*}$ is non-zero, then $H^i(X, \omega(-\infty))_{\mathfrak{m}_Q}$ is also non-zero. The result now follows as in the proof of Proposition 7.9.8 of [BCGP18], which identifies these groups for all i under the Taylor–Wiles hypothesis using Lemmas 2.4.36 and 2.4.37 of [BCGP18]. \square

4. PROOFS

Let R denote the minimal deformation ring of \bar{r} defined as follows:

- (1) **Coherent Case:** R is the minimal ordinary deformation ring denoted by $R^{\min} = R_\emptyset$ in §4 of [CG18].
- (2) **Betti Case:** R is the minimal ordinary deformation ring corresponding to the following conditions:
 - (a) If v is a prime of bad reduction (so we are assuming, for a topological generator σ of tame inertia, that $\bar{r}(\sigma)$ is unipotent with a single block), then we take the local deformation ring to be the ring R_v^1 in §8.5.1 of [CG18]. Note that, if $r(\sigma)$ on A^n has characteristic polynomial $(X - 1)^n$, then (given our assumption on \bar{r}) this deformation problem coincides with the minimal condition in Definition 2.4.14 of [CHT08], namely, that the map

$$\ker(r(\sigma) - \text{id}_n)^r \otimes_R k \rightarrow \ker(\bar{r}(\sigma) - \text{id}_n)^r$$

is an isomorphism (equivalently, surjection) for all $r \leq n$. (One can see this equivalence by induction — $\bar{r}(\sigma)$ has a unique eigenvector over k , which lifts to a unique eigenvector over A whose mod- p reduction is non-trivial; now take the representation of A^{n-1} and k^{n-1} given by quotienting out by this eigenvector.)

- (b) If $v|p$ and p is sufficiently large with respect to n and the primes which ramify in F , we take deformations which are Fontaine–Laffaille of weight $[0, 1, \dots, n-1]$.

If one has an isomorphism $R \simeq \mathbf{T}$ for all sufficiently large p satisfying the required hypothesis, then since one also will have an isomorphism $\mathbf{T} \simeq \mathcal{O}$, this would immediately imply that the tangent space to R along the projection to \mathcal{O} is trivial, and hence the corresponding adjoint Selmer groups are trivial. Theorem 1.2 is now a consequence of Theorem 5.16 of [CG18] and Theorem 6.4 of [CG18], where we use the fact that the corresponding local deformation rings are formally smooth (as follows from Corollary 2.4.3 of [CHT08] and Lemma 2.4.19 of [CHT08]), and where we use Theorem 3.2 as a substitute for the vanishing assumption required in Conjecture B of *ibid.* Equally, Theorem 1.1 follows as in Theorem 1.2 of [CG], where the vanishing result of Lemma 3.4 replaces the vanishing results of Lan–Suh [LS13] for other low weight local systems used in [CG]. The modularity argument above requires a large image hypothesis which we assume in Theorem 1.2 and which we are required to prove (for sufficiently large p) for Theorem 3.2. Furthermore, we must justify the claim in Theorem 3.2 that the assumptions hold for a set of primes p of density one. Hence it remains to prove the following:

Lemma 4.1. *Let A be a semistable abelian surface of conductor N with $\text{End}(A) = \mathbf{Z}$. Then:*

- (1) *For sufficiently large p , the residual representation $\bar{r}_p : G_{\mathbf{Q}} \rightarrow \text{GSp}_4(\mathbf{F}_p)$ is surjective with minimal conductor N .*
- (2) *For a set of primes p of density one, we have*

$$\alpha\beta(\alpha^2 - 1)(\beta^2 - 1)(\alpha - \beta)(\alpha^2\beta^2 - 1) \not\equiv 0 \pmod{p}.$$

Proof. Since $\text{End}(A) = \mathbf{Z}$, the residual image is surjective for all sufficiently large p by [Ser00], Corollaire au Théorème 3. In order to ensure that the conductor of \bar{r}_p at a prime ℓ dividing N matches that of A , it suffices to take p co-prime to the (finite) order of the component group Φ_A of the Néron model of A^\vee at ℓ . This proves the first claim.

For the second claim, let the characteristic polynomial of Frobenius (for p not dividing the discriminant on the étale cohomology $V_\ell = H^1(A, \bar{\mathbf{Q}}_\ell)$ at any prime $\ell \neq p$) be

$$X^4 + a_p X^3 + (2p + b_p)X^2 + pa_p X + p^2.$$

Let the roots of this polynomial be α , β , $\alpha^{-1}p$ and $\beta^{-1}p$ respectively; by the Riemann hypothesis for curves (Weil bound) they are Weil numbers of absolute value \sqrt{p} . Note that $2p + b_p$ is the trace of Frob_p on $\wedge^2 V_\ell$ for all but finitely many ℓ , and a_p^2 is the trace of Frob_p on $V_\ell \otimes V_\ell$. We use the following lemma, which is essentially an observation of Ogus (2.7.1 of [DMOS82]).

Lemma 4.2. *There is no fixed linear relation between 1 , p , b_p , a_p , and a_p^2 which can hold for a set $p \in S$ of positive density.*

Proof. From such an equality, we can build two finite dimensional representations A_ℓ and B_ℓ built out of copies of $\wedge^2 V_\ell$, \mathbf{Q}_p , $\mathbf{Q}_p(1)$, V_ℓ , and $V_\ell \otimes V_\ell$ respectively which have equal trace on Frob_p for infinitely many p . There must be at least one quadratic field with a positive density of inert primes in S , twisting by this representation we arrive at a representation W_ℓ with a set S of positive density such that Frob_p has trace zero for $p \in S$. For sufficiently large ℓ , our assumptions on A implies ([Ser00]) that the image of $G_{\mathbf{Q}}$ on V_ℓ is $\text{GSp}_4(\mathbf{Z}_\ell)$ if $\text{End}(A) = \mathbf{Z}$. By Chebotarev, it follows that the appropriate identity must also hold on an subset of these groups of positive measure. Yet the distribution of the appropriate eigenvalues for $\text{GSp}_4(\mathbf{Z}_\ell)$ does not have any atomic measure. In particular, writing the eigenvalues in either case as x , y , δ/x , and δ/y , we would obtain an relation between the polynomials

$$1, \quad \delta, \quad xy + \delta x/y + \delta y/x + \delta^2/xy, \quad x + y + \delta/x + \delta/y, \quad (x + y + \delta/x + \delta/y)^2$$

that holds on an open set (and consequently holds everywhere). There are no such relations by inspection. \square

Returning to the proof of Lemma 4.1, we deduce from the Weil bounds that $|a_p|^2 \leq 16p$ and $|b_p| \leq 4p$. Hence, if we have any linear expression in a_p , a_p^2 , b_p and 1 which is congruent to zero modulo p for a set of positive density, then it must also equal a constant multiple of p for a set of positive density, and we would obtain a contradiction by Lemma 4.2. We show that this holds in each of the possible cases when our congruence above holds. (We take advantage of the symmetry in α and β and consider a reduced number of cases.)

- (1) Suppose that neither β and $\beta^{-1}p$ are units. Then $b_p \equiv 0 \pmod{p}$.
- (2) Suppose that $\alpha\beta \equiv \varepsilon \pmod{p}$ for some fixed $\varepsilon \in \{\pm 1\}$. Then $b_p \equiv \varepsilon \pmod{p}$.
- (3) Suppose that $\alpha = \varepsilon \pmod{p}$ for some fixed $\varepsilon \in \{\pm 1\}$. Then $b_p - \varepsilon a_p + 1 \equiv 0 \pmod{p}$.
- (4) Suppose that $\alpha - \beta \equiv 0 \pmod{p}$. Then $4b_p - a_p^2 \equiv 0 \pmod{p}$.

\square

REFERENCES

- [ACC⁺18] Patrick Allen, Frank Calegari, Ana Caraiani, Toby Gee, David Helm, Bao Le Hung, James Newton, Peter Scholze, Richard Taylor, and Jack Thorne, *Potential automorphy over CM fields*, preprint, 2018.
- [BCGP18] George Boxer, Frank Calegari, Toby Gee, and Vincent Pilloni, *Abelian surfaces over totally real fields are potentially modular*, preprint, 2018.
- [BK90] Spencer Bloch and Kazuya Kato, *L-functions and Tamagawa numbers of motives*, The Grothendieck Festschrift, Vol. I, Progr. Math., vol. 86, Birkhäuser Boston, Boston, MA, 1990, pp. 333–400. MR 1086888
- [BR17] Baskar Balasubramanyam and A. Raghuram, *Special values of adjoint L-functions and congruences for automorphic forms on $GL(n)$ over a number field*, Amer. J. Math. **139** (2017), no. 3, 641–679. MR 3650229
- [BT99] Kevin Buzzard and Richard Taylor, *Companion forms and weight one forms*, Ann. of Math. (2) **149** (1999), no. 3, 905–919. MR 1709306 (2000j:11062)
- [Buz03] Kevin Buzzard, *Analytic continuation of overconvergent eigenforms*, J. Amer. Math. Soc. **16** (2003), no. 1, 29–55 (electronic). MR 1937198 (2004c:11063)
- [BW80] Armand Borel and Nolan R. Wallach, *Continuous cohomology, discrete subgroups, and representations of reductive groups*, Annals of Mathematics Studies, vol. 94, Princeton University Press, Princeton, N.J., 1980. MR 554917 (83c:22018)
- [CG] Frank Calegari and David Geraghty, *Modularity lifting for non-regular symplectic representations*, preprint.
- [CG18] Frank Calegari and David Geraghty, *Modularity lifting beyond the Taylor–Wiles method*, Inventiones mathematicae **211** (2018), no. 1, 297–433.
- [CHT08] Laurent Clozel, Michael Harris, and Richard Taylor, *Automorphy for some l -adic lifts of automorphic mod l Galois representations*, Publ. Math. Inst. Hautes Études Sci. (2008), no. 108, 1–181, With Appendix A, summarizing unpublished work of Russ Mann, and Appendix B by Marie-France Vignéras. MR 2470687 (2010j:11082)
- [CLH16] Ana Caraiani and Bao V. Le Hung, *On the image of complex conjugation in certain Galois representations*, Compos. Math. **152** (2016), no. 7, 1476–1488. MR 3530448
- [Clo90] Laurent Clozel, *Motifs et Formes Automorphes : Applications du Principe de Fonctorialité*, Automorphic forms, Shimura varieties, and L-functions, Vol. I (Ann Arbor, MI, 1988), Perspect. Math., vol. 10, Academic Press, Boston, MA, 1990, pp. 77–159.
- [DFG04] Fred Diamond, Matthias Flach, and Li Guo, *The Tamagawa number conjecture of adjoint motives of modular forms*, Ann. Sci. École Norm. Sup. (4) **37** (2004), no. 5, 663–727. MR 2103471 (2006e:11089)

- [DMOS82] Pierre Deligne, James S. Milne, Arthur Ogus, and Kuang-yen Shih, *Hodge cycles, motives, and Shimura varieties*, Lecture Notes in Mathematics, vol. 900, Springer-Verlag, Berlin-New York, 1982. MR 654325 (84m:14046)
- [Fla92] Matthias Flach, *A finiteness theorem for the symmetric square of an elliptic curve*, Invent. Math. **109** (1992), no. 2, 307–327. MR 1172693 (93g:11066)
- [FS98] Jens Franke and Joachim Schwermer, *A decomposition of spaces of automorphic forms, and the Eisenstein cohomology of arithmetic groups*, Math. Ann. **311** (1998), no. 4, 765–790. MR 1637980
- [GHL16] Harald Grobner, Michael Harris, and Erez Lapid, *Whittaker rational structures and special values of the asai L -function*, Contemporary Math. **664** (2016), 119–134.
- [Hid81] Haruzo Hida, *Congruence of cusp forms and special values of their zeta functions*, Invent. Math. **63** (1981), no. 2, 225–261. MR 610538 (82g:10044)
- [HLTT] Michael Harris, Kai-Wen Lan, Richard Taylor, and Jack Thorne, *Galois representations for regular algebraic cusp forms over CM-fields*, in preparation.
- [JS81] H. Jacquet and J. A. Shalika, *On Euler products and the classification of automorphic forms. II*, Amer. J. Math. **103** (1981), no. 4, 777–815. MR 623137 (82m:10050b)
- [Kis09] Mark Kisin, *Modularity of 2-adic Barsotti-Tate representations*, Invent. Math. **178** (2009), no. 3, 587–634. MR 2551765 (2010k:11089)
- [KT17] Chandrashekhara B. Khare and Jack A. Thorne, *Potential automorphy and the Leopoldt conjecture*, Amer. J. Math. **139** (2017), no. 5, 1205–1273. MR 3702498
- [KW09a] Chandrashekhara Khare and Jean-Pierre Wintenberger, *Serre’s modularity conjecture. I*, Invent. Math. **178** (2009), no. 3, 485–504. MR 2551763 (2010k:11087)
- [KW09b] ———, *Serre’s modularity conjecture. II*, Invent. Math. **178** (2009), no. 3, 505–586. MR 2551764 (2010k:11088)
- [LS13] Kai-Wen Lan and Junecue Suh, *Vanishing theorems for torsion automorphic sheaves on general PEL-type Shimura varieties*, Adv. Math. **242** (2013), 228–286. MR 3055995
- [Sch15] Peter Scholze, *On torsion in the cohomology of locally symmetric varieties*, Ann. of Math. (2) **182** (2015), no. 3, 945–1066. MR 3418533
- [Ser00] Jean-Pierre Serre, *Lettre à Marie-France Vignéras du 10/2/1986*, Œuvres. Collected papers. IV, Springer-Verlag, Berlin, 2000, 1985–1998, pp. viii+657. MR 1730973
- [TW95] Richard Taylor and Andrew Wiles, *Ring-theoretic properties of certain Hecke algebras*, Ann. of Math. (2) **141** (1995), no. 3, 553–572. MR 1333036 (96d:11072)
- [Urb98] Eric Urban, *Module de congruences pour $GL(2)$ d’un corps imaginaire quadratique et théorie d’Iwasawa d’un corps CM biquadratique*, Duke Math. J. **92** (1998), no. 1, 179–220. MR 1611003 (98m:11035)
- [Var14] Ila Varma, *Local-global compatibility for regular algebraic cuspidal automorphic representations when $\ell \neq p$* , Preprint, 2014.
- [Wil95] Andrew Wiles, *Modular elliptic curves and Fermat’s last theorem*, Ann. of Math. (2) **141** (1995), no. 3, 443–551. MR 1333035 (96d:11071)
- [Wit21] Ludwig Wittgenstein, *Tractatus logico-philosophicus*, Harcourt, Brace & company, 1921.