

Topology and arithmetic of resultants, II: the resultant = 1 hypersurface

Benson Farb and Jesse Wolfson

With an appendix by Christophe Cazanave¹

ABSTRACT

We consider the moduli space \mathcal{R}_n of pairs of monic, degree n polynomials whose resultant equals 1. We relate the topology of these algebraic varieties to their geometry and arithmetic. In particular, we compute their étale cohomology, the associated eigenvalues of Frobenius, and the cardinality of their set of \mathbb{F}_q -points. When q and n are coprime, we show that the étale cohomology of $\mathcal{R}_n/\overline{\mathbb{F}_q}$ is pure, and of Tate type if and only if $q \equiv 1 \pmod n$. We also deduce the values of these invariants for the finite field counterparts of the moduli spaces \mathcal{M}_n of $\mathrm{SU}(2)$ monopoles of charge n in \mathbb{R}^3 , and the associated moduli space X_n of strongly centered monopoles.

An appendix by Cazanave gives an alternative and elementary computation of the point counts.

1. Introduction

Consider two monic, degree $n \geq 1$ complex polynomials

$$\phi(z) = z^n + a_{n-1}z^{n-1} + \cdots + a_1z + a_0$$

and

$$\psi(z) = z^n + b_{n-1}z^{n-1} + \cdots + b_1z + b_0.$$

A beautiful classical fact is that the condition for ϕ and ψ to have a common root is polynomial in the coefficients a_i and b_j . More precisely, ϕ and ψ have a common root if and only if

$$\mathcal{R}(\phi, \psi) := \mathcal{R}(a_0, \dots, a_{n-1}, b_0, \dots, b_{n-1}) = 0 \tag{1.1}$$

Lab. de Mathématiques J.A. Dieudonné, Université de Nice Sophia-Antipolis, France
E-mail: cazanave@unice.fr

2010 Mathematics Subject Classification 55R80 (Primary), 11G20 (Secondary)

Keywords: Resultant, monopoles, étale cohomology.

B.F. is supported in part by NSF Grant Nos. DMS-1105643 and DMS-1406209. J.W. is supported in part by NSF Grant No. DMS-1400349.

where \mathcal{R} is the *resultant*, given by

$$\mathcal{R}(\phi, \psi) = \det \begin{bmatrix} a_0 & a_1 & \cdots & a_{n-1} & 1 & 0 & \cdots & 0 \\ 0 & a_0 & \cdots & \cdots & a_{n-1} & 1 & \cdots & 0 \\ \vdots & \vdots \\ b_0 & b_1 & \cdots & b_{n-1} & 1 & 0 & \cdots & 0 \\ 0 & b_0 & \cdots & \cdots & b_{n-1} & 1 & \cdots & 0 \\ \vdots & \vdots \\ 0 & 0 & \cdots & b_0 & \cdots & \cdots & b_{n-1} & 1 \end{bmatrix}$$

This is a homogeneous polynomial of degree n in the a_i and similarly in the b_i . It has integer coefficients. Fix a field k , and denote by \mathbb{A}^n the affine space over k . The resultant can be thought of as a map

$$\mathcal{R} : \mathbb{A}^{2n} \longrightarrow \mathbb{A}^1$$

from the space \mathbb{A}^{2n} of pairs of monic, degree n polynomials to k . The *resultant locus* $\mathcal{M}_n := \mathbb{A}^{2n} \setminus \mathcal{R}^{-1}(0)$ is a classically studied object. It is isomorphic to the moduli space of degree n rational maps $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ taking ∞ to 1. Harder to understand is the “*resultant = 1*” hypersurface $\mathcal{R}_n := \mathcal{R}^{-1}(1)$ in \mathbb{A}^{2n} .

Since the polynomial \mathcal{R} has integer coefficients, we can extend scalars to \mathbb{C} and consider the complex points $\mathcal{R}_n(\mathbb{C})$, and we can also reduce modulo p for any prime p . This gives a variety defined over \mathbb{F}_p , and for any positive power $q = p^d$ we can consider both the \mathbb{F}_q -points as well as the $\overline{\mathbb{F}}_q$ -points of \mathcal{R}_n , where $\overline{\mathbb{F}}_q$ is the algebraic closure of \mathbb{F}_q . Three of the most fundamental arithmetic invariants attached to a such a variety \mathcal{R}_n are:

- (i) The cardinality $|\mathcal{R}_n(\mathbb{F}_q)|$.
- (ii) The étale cohomology $H_{et}^*(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$, where ℓ is a prime not dividing q .
- (iii) The eigenvalues of the (geometric) Frobenius morphism

$$\text{Frob}_q : H_{et}^*(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell) \longrightarrow H_{et}^*(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell).$$

Our main theorems compute the étale cohomology of \mathcal{R}_n as well as the associated eigenvalues of Frobenius, building on the topological work of Segal and Selby [SS96]. We then apply this to compute the cardinality of finite field versions of these moduli spaces; that is, of $\mathcal{R}_n(\mathbb{F}_q)$ and $X_n(\mathbb{F}_q)$, where \mathbb{F}_q is a finite field.

There is a canonical μ_n -action on \mathcal{R}_n ; see Section 2. This induces a μ_n -action on $H_{et}^*(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$. Fix an algebraic closure $\overline{\mathbb{Q}}_\ell$. Since \mathbb{Q}_ℓ has characteristic 0, it follows that $H_{et}^*(\mathcal{R}_n/\overline{\mathbb{F}}_q; \overline{\mathbb{Q}}_\ell) := H_{et}^*(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell) \otimes_{\mathbb{Q}_\ell} \overline{\mathbb{Q}}_\ell$ decomposes into a direct sum of irreducible representations of μ_n . The irreducible representations of μ_n are parametrized by integers m with $0 \leq m < n$, corresponding to $\xi \mapsto \zeta^m \xi$, where ζ is a primitive n^{th} root of unity; denote this irreducible representation by V_m . Let $H_{et}^*(\mathcal{R}_n)_m$ denote the isotypic component of $H_{et}^*(\mathcal{R}_n; \overline{\mathbb{Q}}_\ell)$ corresponding to V_m . Denote by $H_{et}^i(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)^{\mu_n}$ the subspace of μ_n -fixed vectors.

Denote by $\mathbb{Q}_\ell(-i)$ the rank 1 $\text{Gal}(\overline{\mathbb{F}}_q/\mathbb{F}_q)$ -representation on which Frobenius acts by q^i . Also, recall that the étale cohomology of a variety X is *pure* if the absolute values of the eigenvalues of Frob_q on H_{et}^i are all $q^{\frac{i}{2}}$; as Deligne showed, this is always the case when X is smooth and projective. The group $H_{et}^i(X_{\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ is of *Tate type* if the eigenvalues of Frob_q are all equal to integer powers of q .

THEOREM 1.1 Étale cohomology of \mathcal{R}_n . Let $n \geq 1$. For all primes p not dividing n , and for all positive powers $q = p^d$:

- (i) The ℓ -adic cohomology of $\mathcal{R}_n/\overline{\mathbb{F}}_q$ is pure.
- (ii) The μ_n -invariants are concentrated in degree 0:

$$H_{et}^i(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)^{\mu_n} \cong \begin{cases} \mathbb{Q}_\ell(0) & i = 0 \\ 0 & i \neq 0 \end{cases}$$

- (iii) $H_{et}^{2i+1}(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell) = 0$ for all i .
- (iv) For $i > 0$, $H_{et}^{2i}(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$ is nonzero if and only if $i < n$ and $(n-i)|n$. In this case, it is of Tate type if and only if $q \equiv 1 \pmod{\frac{n}{n-i}}$. More precisely, let $\mathcal{O}_a := \{1 \leq m \leq n \mid (m, n) = a\}$. Then we have a Frob_q -invariant decomposition

$$H_{et}^{2i}(\mathcal{R}_n/\overline{\mathbb{F}}_q; \overline{\mathbb{Q}}_\ell) \cong \left(\bigoplus_{m \in \mathcal{O}_{n-i}} \overline{\mathbb{Q}}_\ell \right) \otimes_{\mathbb{Q}_\ell} \mathbb{Q}_\ell(-i)$$

where Frob_q acts on the direct sum by $(\overline{\mathbb{Q}}_\ell)_m \xrightarrow{1} (\overline{\mathbb{Q}}_\ell)_{qm \bmod n}$, where $(\overline{\mathbb{Q}}_\ell)_m$ denotes the summand corresponding to a given $m \in \mathcal{O}_{n-i}$. In particular:

- (a) If $(n-i) \nmid n$, then the rank of $H_{et}^{2i}(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$ is 0. If $i < n$ and $(n-i)|n$, then $H_{et}^{2i}(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$ has rank equal to Euler's totient function $\phi(\frac{n}{n-i})$.
- (b) The Frob_q action is given by the permutation representation coming from the action of Frob_q on \mathcal{O}_{n-i} .
- (c) When $q \not\equiv 1 \pmod{\frac{n}{n-i}}$, the trace of this representation is 0.
- (d) When $q \equiv 1 \pmod{\frac{n}{n-i}}$, Frob_q acts on $H_{et}^{2i}(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$ by multiplication by q^i , and the trace is $q^i \cdot \phi(\frac{n}{n-i})$.

Remark 1.2.

- (i) The variety $\mathcal{R}_n/\overline{\mathbb{F}}_q$ is smooth (so long as $(q, n) = 1$), but not projective, and thus purity does not follow from Deligne.
- (ii) The failure of $H_{et}^{2i}(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$ to be of Tate type is precisely the failure of the group-scheme $\mu_{\frac{n}{n-i}}$ to be of Tate type over \mathbb{F}_q . See below for details.

COROLLARY 1.3 Isotypic decomposition when $q \equiv 1 \pmod n$. Let $n \geq 1$ as above. Then for $q \equiv 1 \pmod n$, the isotypic decomposition of $H_{et}^*(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$ is Frob_q -invariant, and for any $1 \leq m \leq n-1$:

$$H_{et}^{2i}(\mathcal{R}_n/\overline{\mathbb{F}}_q)_m \cong \begin{cases} \mathbb{Q}_\ell(-i) & \gcd(m, n) = n-i \\ 0 & \text{else} \end{cases}$$

Thus, for each $i \geq 0$:

$$H_{et}^{2i}(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell) \cong \bigoplus_{\{m \in \mathcal{O}_{n-i}\}} \mathbb{Q}_\ell(-i)$$

Note that for n prime and $q \equiv 1 \pmod n$,

$$H_{et}^i(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell) \begin{cases} \mathbb{Q}_\ell(1-n)^{\oplus(n-1)} & i = 2n-2 \\ \mathbb{Q}_\ell(0) & i = 0 \\ 0 & \text{else} \end{cases}$$

Theorem 1.1 gives not only the étale cohomology of the varieties \mathcal{R}_n , but it also computes the eigenvalues of Frobenius acting on these varieties over $\overline{\mathbb{F}}_q$. Applying the Grothendieck-Lefschetz trace formula, we conclude the following.

COROLLARY 1.4 Cardinality of $\mathcal{R}_n(\mathbb{F}_q)$. *Let $n \geq 1$. Then for all primes $p \nmid n$, for each positive power q of p , let*

$$\mathcal{F}(q, n) : \{a : a|n \text{ and } q \equiv 1 \pmod{\frac{n}{a}}\}.$$

Then

$$|\mathcal{R}_n(\mathbb{F}_q)| = q^{2n-1} \left(\sum_{a \in \mathcal{F}(q, n)} \phi\left(\frac{n}{a}\right) \cdot q^{a-n} \right).$$

In particular, for $(q-1, n) = 1$,

$$|\mathcal{R}_n(\mathbb{F}_q)| = q^{2n-1},$$

and for $q \equiv 1 \pmod n$,

$$|\mathcal{R}_n(\mathbb{F}_q)| = q^{2n-1} \left(\sum_{i=0}^{n-1} \phi\left(\frac{n}{n-i}\right) \cdot q^{-i} \right).$$


where we define Euler's totient function to be identically 0 on $\mathbb{Q} \setminus \mathbb{N}$. When n is prime and $q \equiv 1 \pmod n$, this gives

$$|\mathcal{R}_n(\mathbb{F}_q)| = q^{2n-1} + (n-1)q^n.$$

We also include an appendix, by Cazanave, in which the above point count (as well as the case when $(q, n) \neq 1$) is deduced by elementary means, i.e. without using étale cohomology.

The results of this paper provide an example of a broader program applying this viewpoint to make concrete calculations for various moduli spaces. For more, see [FW16]. The varieties \mathcal{R}_n are closely related to some moduli spaces studied in physics, namely moduli spaces of magnetic monopoles. In §3 we apply the results above to deduce similar theorems for these moduli spaces.

Remark on the proofs. One novelty of the proofs in this paper is that we obtain information about various algebraic varieties Z defined over \mathbb{Z} by traversing, in different directions, a “triangle” of viewpoints: arithmetic ($|Z(\mathbb{F}_q)|$); topological ($H^*(Z(\mathbb{C}); \mathbb{Q})$); and geometric (Frob $_q$ acting on $H_{et}^*(Z/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$). As an example, the logic of the starting point of the proofs of Theorem 1.1 and Corollary 1.4 is as follows:



The upper horizontal deduction is a special case of the results of the first paper in this series [FW16]. We take this as a starting point in order to make the remaining deductions.

Acknowledgements. We are grateful to Sasha Beilinson, Weiyang Chen, Jordan Ellenberg, Matt Emerton, Nir Gadish, Jackson Hance, Sean Howe, Peter May, Joel Specter, Shmuel Weinberger, and Melanie Wood for helpful conversations. We thank Joe Silverman for helpful comments on an earlier draft. We would especially like to thank Christophe Cazanave, not only for contributing the appendix, but also for his many comments, corrections and suggestions; they have greatly improved this paper. Finally, we thank the anonymous referee for many useful comments which greatly improved the paper.

2. Proof of Theorem 1.1

In this section we prove Theorem 1.1. Throughout this section we fix $n \geq 1$ and a prime power $q = p^d, d \geq 1$.

The μ_n -action on \mathcal{R}_n . The multiplicative group \mathbb{G}_m acts on \mathcal{M}_n by²

$$\lambda \cdot (\phi, \psi) := (\psi + \lambda(\phi - \psi), \psi).$$

Because the resultant is a homogeneous of degree n with respect to this action, this induces an action of the group μ_n of n^{th} -roots of unity on \mathcal{R}_n . We study this action on the étale cohomology of \mathcal{R}_n .

Recall that the resultant gives a morphism $\mathcal{R} : \mathbb{A}^{2n} \rightarrow \mathbb{A}^1$. Note that

$$\mathcal{R}_1 = \left\{ \frac{z + a_0}{z + b_0} : a_0 - b_0 = 1 \right\} \cong \left\{ \frac{z + a_0}{z + (a_0 - 1)} \right\} \cong \mathbb{A}^1$$

as \mathbb{F}_q -varieties. Thus the theorem is true when $n = 1$. We now show the theorem for $n > 1$. Our analysis proceeds in a series of steps.

Step 1 (The μ_n -isotypic decomposition of $H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$): For each $i \geq 0$ there is a decomposition of $H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ into μ_n -isotypic components :

$$H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell) \cong H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)^{\mu_n} \bigoplus H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q})^{\mu_n^\perp} \quad (2.1)$$

where

$$H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q}; \overline{\mathbb{Q}}_\ell)^{\mu_n^\perp} := \bigoplus_{m=1}^{n-1} H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q})_m.$$

The decomposition (2.1) is invariant under the action of Frob_q . However, for $m > 0$ the subspace $H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q})_m$ is in general *not* Frob_q -invariant. In fact, this failure of invariance is at the crux of the proof of the theorem. We begin by finding the Frob_q -invariant subspaces.

To this end, for any factor a of n , define

$$\mathcal{O}_a := \{m \mid 1 \leq m \leq n - 1, (m, n) = a\}$$

²Note that this action is isomorphic to the one considered by [SS96] under conjugation by the Möbius transformation $z \mapsto z + 1$.

and define

$$H_{et}^i(\mathcal{R}_{n/\mathbb{F}_q})_a := \bigoplus_{m \in \mathcal{O}_a} H_{et}^i(\mathcal{R}_{n/\mathbb{F}_q})_m.$$

Fix $n \geq 1$ and assume that $(q, n) = 1$. We claim that for each $i \geq 0$ the splitting

$$H_{et}^i(\mathcal{R}_{n/\mathbb{F}_q}; \overline{\mathbb{Q}}_\ell)^{\mu_n^\perp} \cong \bigoplus_{a|n} H_{et}^i(\mathcal{R}_{n/\mathbb{F}_q})_a \quad (2.2)$$

is Frob_q -equivariant. To see this, note that since $(q, n) = 1$, multiplication of q acts by an automorphism of $\mathbb{Z}/n\mathbb{Z}$, and so preserves the order of elements in $\mathbb{Z}/n\mathbb{Z}$. For $m \in \mathcal{O}_a$, the order of m in $\mathbb{Z}/n\mathbb{Z}$ equals n/a . Thus the order of qm in $\mathbb{Z}/n\mathbb{Z}$ is also n/a , and thus the greatest common divisor of $qm \bmod n$ and n equals a , proving the claim.

Step 2 (The μ_n -invariant part of $H_{et}^i(\mathcal{R}_{n/\mathbb{F}_q}; \mathbb{Q}_\ell)$): First note that $\mathcal{R} - 1$ is an irreducible polynomial. Indeed, \mathcal{R} is an irreducible polynomial [vdW50, Section 77] and

LEMMA 2.1. *Let $\Phi(x_1, \dots, x_n, y_1, \dots, y_m)$ be an irreducible, bi-homogeneous polynomial of bi-degree (p, q) . Then $\Phi - 1$ is irreducible.*

Proof. Suppose $\Phi - 1 = PQ$. Without loss of generality, we can assume that P and Q are of total degrees c and d . Write $P = P_0 + P_1$ where P_0 is homogenous of total degree c and $\deg(P_1) < c$, and, similarly, write $Q = Q_0 + Q_1$. Then $c + d = p + q$, and

$$\begin{aligned} \Phi &= P_0Q_0, \text{ and} \\ -1 &= P_0Q_1 + P_1Q_0 + P_1Q_1 \end{aligned}$$

The irreducibility of Φ implies that, without loss of generality, $P_0 = 1$ and thus $P_1 = 0$. Therefore, we have that $Q = \Phi - 1$ with $Q_0 = \Phi$ and $Q_1 = -1$. \square

It follows that $H_{et}^0(\mathcal{R}_{n/\mathbb{F}_q}; \mathbb{Q}_\ell) \cong \mathbb{Q}_\ell(0)$. Since this group is generated by the constant function 1, it follows that $H_{et}^0(\mathcal{R}_{n/\mathbb{F}_q}; \mathbb{Q}_\ell) \subseteq H_{et}^i(\mathcal{R}_{n/\mathbb{F}_q}; \mathbb{Q}_\ell)^{\mu_n}$. We now prove the reverse inclusion.

Recall that we have defined $\mathcal{M}_n := \mathbb{A}^{2n} \setminus \mathcal{R}^{-1}(0)$. The variety \mathcal{M}_n admits a free action of the multiplicative group $\mathbb{G}_m = \text{GL}_1$ via $\lambda \cdot \frac{\phi}{\psi} := \frac{\lambda\phi}{\psi}$. Because the group \mathbb{G}_m is special (cf. [Gro58, p. 12]), This gives a Zariski-locally trivial fibering

$$\mathbb{G}_m \longrightarrow \mathcal{M}_n \longrightarrow \mathcal{M}_n/\mathbb{G}_m \cong \mathcal{R}_n/\mu_n. \quad (2.3)$$

where $\mathcal{M}_n/\mathbb{G}_m$ is a fiber bundle over \mathbb{P}^{n-1} .³

Transfer now gives:

$$H_{et}^i(\mathcal{R}_{n/\mathbb{F}_q}; \mathbb{Q}_\ell)^{\mu_n} \cong H_{et}^i(\mathcal{R}_n/\mu_n/\mathbb{F}_q; \mathbb{Q}_\ell) \cong H_{et}^i(\mathcal{M}_n/\mathbb{G}_m/\mathbb{F}_q; \mathbb{Q}_\ell).$$

It is therefore enough to prove that $H_{et}^i(\mathcal{M}_n/\mathbb{G}_m/\mathbb{F}_q; \mathbb{Q}_\ell)$ vanishes except when $i = 0$, in which case (as observed above) it is isomorphic to $\mathbb{Q}_\ell(0)$. To this end, we apply the Leray spectral sequence in étale cohomology⁴ to the fibering (2.3). This spectral sequence has

$$E_2^{i,j} = \begin{cases} H_{et}^i(\mathcal{M}_n/\mathbb{G}_m/\mathbb{F}_q; H_{et}^j(\mathbb{G}_m/\mathbb{F}_q; \mathbb{Q}_\ell)) & \text{if } i, j \geq 0 \\ 0 & \text{else} \end{cases}$$

³Note that the projection $\frac{\phi}{\psi} \mapsto \phi$ induces a \mathbb{G}_m -equivariant fibering $\mathcal{M}_n \longrightarrow (\mathbb{A}^n - \{0\})$, and $\mathcal{M}_n/\mathbb{G}_m \longrightarrow \mathbb{P}^{n-1}$ is the quotient of this \mathbb{G}_m -action.

⁴One reference for this spectral sequence is Theorem 12.7 of [Mil13].

and the spectral sequence converges to $H_{et}^i(\mathcal{M}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$. This cohomology is computed as a special case of Theorem 1.2 of [FW16], where in the notation of [FW16] the variety \mathcal{M}_n was called $\text{Poly}_1^{n,2}$. It is as follows:

$$H_{et}^i(\mathcal{M}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell) \cong \begin{cases} \mathbb{Q}_\ell(0) & i = 0 \\ \mathbb{Q}_\ell(-1) & i = 1 \\ 0 & \text{else} \end{cases} \quad (2.4)$$

We use this to compute $E_2^{i,j}$. To start, we claim that the monodromy action on $H_{et}^j(\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ is trivial. To see this, first recall

$$H_{et}^j(\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell) \cong \mathbb{Q}_\ell(-j)$$

for $j = 0, 1$ and equals 0 for $j > 1$. Over \mathbb{C} , $\pi_1(\mathcal{M}_n/\mathbb{G}_m) \cong \pi_1(\mathcal{R}_n/\mu_n) \cong \mu_n$ since $\pi_1(\mathcal{R}_n) = 0$. Now use the fact that the μ_n action on $H_{et}^j(\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ is the restriction of the action of \mathbb{G}_m induced by left multiplication. Because $H^0(\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell) = \mathbb{Q}_\ell$, this action is trivial. Therefore, the μ_n action on $H_{et}^j(\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ is trivial as well. We have thus shown:

$$E_2^{i,j} = \begin{cases} H_{et}^i(\mathcal{M}_n/\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell(-j)) & \text{if } j = 0, 1 \\ 0 & \text{else} \end{cases}$$

The differential $d_2^{i,j} : E_2^{i,j} \rightarrow E_2^{i+2,j-1}$ thus gives, for each $i \geq 0$, a homomorphism

$$d_2^{i,1} : H_{et}^i(\mathcal{M}_n/\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell(-j)) \rightarrow H_{et}^{i+2}(\mathcal{M}_n/\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell(1-j)).$$

Since $E_2^{i,j} = 0$ for $i > 1$ and $j < 0$, the only nontrivial differentials occur on the E_2 page, and, for each $i > 0$:

$$H_{et}^i(\mathcal{M}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell) \cong \ker(d_2^{i-1,1}) \oplus H_{et}^i(\mathcal{M}_n/\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell(0))/\text{image}(d_2^{i-1,1}) \quad (2.5)$$

while $H_{et}^0(\mathcal{M}_n/\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell(0)) \cong H_{et}^0(\mathcal{M}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell) \cong \mathbb{Q}_\ell(0)$.

Equation (2.4) now gives that $H_{et}^i(\mathcal{M}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell) \cong \mathbb{Q}_\ell(-i)$ for $i = 0, 1$ and equals 0 for $i > 1$. Now, the target of d on $E_2^{i,j}$ is 0 for $i = 4n, 4n - 1$, so these entries vanish. Working backwards, starting at $i = 4n$ and working down to $i = 1$, we can apply Equation (2.5) using that the left-hand side equals 0, to conclude that $H_{et}^i(\mathcal{M}_n/\mathbb{G}_{m/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell) = 0$ for $i \geq 1$. This concludes the computation of $H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)^{\mu_n}$.

Step 3 (The $H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q})_a$): In this step we analyze the individual summands $H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q})_a$ of the decomposition in Equation (2.2). We will prove:

$$H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q})_a \cong \begin{cases} (\bigoplus_{m \in \mathcal{O}_a} \overline{\mathbb{Q}}_\ell(0)) \otimes_{\mathbb{Q}_\ell} \mathbb{Q}_\ell(a-n) & j - 2(n-a) = 0 \\ 0 & j \neq 0 \end{cases}$$

Given this claim, the bijection

$$\begin{aligned} \mathcal{O}_a &\xrightarrow{\cong} \{m' \leq \frac{n}{a} \mid \gcd(m', \frac{n}{a}) = 1\} \\ m &\mapsto \frac{m}{a} \end{aligned}$$

implies that $H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q}; \mathbb{Q}_\ell)$ has rank $\phi(\frac{n}{n-i})$. To prove the claim, recall that for $a|n$ we defined

$$\mathcal{O}_a := \{m \mid 1 \leq m \leq n-1, (m, n) = a\}.$$

For any $m \in \mathcal{O}_a$ note that the order of $e^{2\pi im/n}$ is n/a . For each $a|n$, define

$$Y_{n,a} := \left\{ \frac{\phi}{\psi} \in \mathcal{R}_n : \psi(z) = \chi(z)^{n/a} \text{ for some } \chi(z) \in k[z], \deg(\chi) = a \right\}.$$

Over any field K containing a *primitive* n^{th} root of unity, Segal and Selby [SS96, Proposition 2.1] construct an isomorphism⁵

$$Y_{n,a/K} \cong \mu_n \times_{\mu_a} (\mathcal{R}_{a/K} \times \mathbb{A}_K^{n-a}). \quad (2.6)$$

In fact, as we now show, these varieties are isomorphic over $K = \mathbb{F}_q$ for any q .

PROPOSITION 2.2. *The isomorphism (2.6) is defined over \mathbb{F}_q , i.e.*

$$Y_{n,a} \cong \mu_n \times_{\mu_a} (\mathcal{R}_a \times \mathbb{A}^{n-a})$$

as \mathbb{F}_q -varieties.

Proof. The homogeneity of the resultant implies that $\mathcal{R}(\phi, \chi^{n/a}) = \mathcal{R}(\phi, \chi)^{n/a}$. Thus, for any $\frac{\phi}{\psi} \in Y_{n,a}$, $\mathcal{R}(\phi, \chi)$ is an n/a^{th} root of unity. Over $\overline{\mathbb{F}}_q$, this gives a decomposition

$$Y_{n,a} \cong \coprod_{\lambda \in \mu_{n/a}} Y_{n,a,\lambda}$$

where $Y_{n,a,\lambda} := \left\{ \frac{\phi}{\psi} \in Y_{n,a} : \psi(z) = \chi(z)^{n/a} \text{ with } \mathcal{R}(\phi, \chi) = \lambda \right\}$. Following Segal and Selby, given $\frac{\phi}{\chi^{n/a}} \in Y_{n,a}$ we can write

$$\phi = \phi_0 \chi + \phi_1$$

where $\deg(\phi_0) < n - a$ and $\deg(\phi_1) < a$. Using this, the assignment

$$\frac{\phi}{\chi^{n/a}} \mapsto \left(\frac{\phi_1}{\chi}, \phi_0 \right)$$

defines an isomorphism over $\overline{\mathbb{F}}_q$

$$Y_{n,a,\lambda} \cong \mathcal{R}_a^{-1}(\lambda) \times \mathbb{A}^{n-a}.$$

Now a primitive n^{th} root of unity ζ gives an isomorphism of $\overline{\mathbb{F}}_q$ -varieties

$$\begin{aligned} \mathcal{R}_a^{-1}(\zeta^{ja}) &\xrightarrow[\cong]{\zeta^{-j}} \mathcal{R}_a \\ \frac{\phi}{\psi} &\mapsto \frac{\zeta^{-j}\phi}{\psi}. \end{aligned}$$

Taken together, these isomorphisms define a μ_n -equivariant map

$$Y_{n,a} \xrightarrow{\cong} \coprod_{\lambda \in \mu_{n/a}} Y_{n,a,\lambda} \xrightarrow{\cong} \coprod_{\lambda \in \mu_{n/a}} \mathcal{R}_a^{-1}(\lambda) \times \mathbb{A}^{n-a} \xrightarrow{\sqcup_j \zeta^{-j}} \mathcal{R}_a \times \mathbb{A}^{n-a} \quad (2.7)$$

where the μ_n action on $\mathcal{R}_a^{-1}(\lambda)$ factors through the action of μ_a . By Galois descent, it suffices to show that the map (2.7) is Frob_q -equivariant. But, by the definition of Frobenius, we have

$$\text{Frob}_q\left(\frac{\zeta^{-j}\phi}{\psi}\right) = \zeta^{-jq} \text{Frob}_q\left(\frac{\phi}{\psi}\right)$$

⁵While Proposition 2.1 of [SS96] is stated only over the field \mathbb{C} , the proof works *verbatim* over any field K containing a primitive n^{th} root of unity.

which shows that, for any $\lambda \in \mu_{n/a}$ the square

$$\begin{array}{ccc} Y_{n,a,\lambda} & \xrightarrow{\text{Frob}_q} & Y_{n,a,\lambda^q} \\ \lambda^{-1/a} \downarrow & & \downarrow \lambda^{-q/a} \\ Y_{n,a,1} & \xrightarrow{\text{Frob}_q} & Y_{n,a,1} \end{array},$$

and thus the map (2.7) is Frob_q -equivariant, and thus defined over \mathbb{F}_q . Combining this with the resultant

$$\mathcal{R}: Y_{n,a} \longrightarrow \mu_{n/a}$$

we obtain a map of \mathbb{F}_q -varieties

$$Y_{n,a} \xrightarrow{\pi} \mu_{n/a} \times (\mathcal{R}_a \times \mathbb{A}^{n-a})$$

which splits the torsor (2.8) below, over $\overline{\mathbb{F}_q}$. By inspection, this is an isomorphism over $\overline{\mathbb{F}_q}$, and because π is Frob_q -equivariant, π^{-1} is as well. Thus we conclude that π is an isomorphism over \mathbb{F}_q .

By inspection, we see that π induces an isomorphism of μ_n -varieties over \mathbb{F}_q

$$Y_{n,a} \cong \mu_n \times_{\mu_a} (\mathcal{R}_a \times \mathbb{A}^{n-a}). \quad (2.8)$$

□

Remark 2.3. We remark that we are using here, in a crucial way, the condition that $\mathcal{R}(\phi, \psi) = 1$ (as opposed, say, to $\mathcal{R}(\phi, \psi) = 2$), since $\mathcal{R}(\phi, \psi) = \mathcal{R}(\phi, \chi)^{n/a}$, and so $\mathcal{R}(\phi, \chi)$ is an n/a^{th} root of unity.

Proposition 2.2 implies that, as μ_n -representations,

$$H_{\text{et}}^i(Y_{n,a}/\overline{\mathbb{F}_q}; \mathbb{Q}_\ell) \cong \text{Ind}_{\mu_a}^{\mu_n} H_{\text{et}}^i(\mathcal{R}_a/\overline{\mathbb{F}_q}; \mathbb{Q}_\ell) \quad (2.9)$$

for each $i \geq 0$.

LEMMA 2.4. *For any $a \mid n$, we have for every positive power q of p :*

$$H_{\text{et}}^*((\mathcal{R}_n - Y_{n,a})/\overline{\mathbb{F}_q})_a := \bigoplus_{m \in \mathcal{O}_a} H_{\text{et}}^*((\mathcal{R}_n - Y_{n,a})/\overline{\mathbb{F}_q})_m = 0.$$

Proof. The analogous lemma over \mathbb{C} is Proposition 2.2 in [SS96]. We now adapt their proof to the étale setting.

For any $n_1, n_2 > 0$ with $n = n_1 + n_2$, let $\mathcal{U}_{n_1, n_2} \longrightarrow \mathcal{R}_n$ be the étale neighborhood consisting of all triples

$$\mathcal{U} := \{(\phi, \psi_1, \psi_2) \mid \text{deg}(\psi_i) = n_i \text{ and } \mathcal{R}(\psi_1, \psi_2) = \mathcal{R}(\phi, \psi_1\psi_2) = 1\}.$$

Given $(\phi, \psi_1, \psi_2) \in \mathcal{U}_{n_1, n_2}$, there is a unique decomposition

$$\frac{\phi}{\psi_1\psi_2} = \frac{\phi_1}{\psi_1} + \frac{\phi_2}{\psi_2}.$$

Let G be the group

$$G := \{(\zeta_1, \zeta_2) \in \mathbb{G}_m \times \mathbb{G}_m \mid \zeta_1^{n_1} \zeta_2^{n_2} = 1\}.$$

The group G acts on \mathcal{U}_{n_1, n_2} via

$$(\zeta_1, \zeta_2) \cdot (\phi, \psi_1, \psi_2) := \frac{\zeta_1 \phi_1}{\psi_1} + \frac{\zeta_2 \phi_2}{\psi_2}.$$

The action of (ζ_1, ζ_2) multiplies $\mathcal{R}(\phi, \psi_1 \psi_2)$ by $\zeta_1^{n_1} \zeta_2^{n_2} = 1$ since:

$$\begin{aligned} \mathcal{R}(\phi, \psi_1 \psi_2) &= \mathcal{R}(\phi_1 \psi_2 + \phi_2 \psi_1, \psi_1 \psi_2) \\ &= (-1)^{n_1 n_2} \mathcal{R}(\phi_1, \psi_1) \mathcal{R}(\phi_2, \psi_2) \mathcal{R}(\psi_1, \psi_2)^2. \end{aligned}$$

Let $d = \gcd(n_1, n_2)$, and let b, c be the integers such that $bn_1 + cn_2 = d$. Observe that we have an isomorphism of algebraic groups

$$\begin{aligned} G &\xrightarrow{\cong} \mathbb{G}_m \times \mu_d \\ (\zeta_1, \zeta_2) &\mapsto (\zeta_1^c \zeta_2^{-b}, \zeta_1^{n_1/d} \zeta_2^{n_2/d}). \end{aligned}$$

The action of G extends the action of μ_n , which can be identified with the diagonal subgroup of G . Because $H_{et}^0(\mathbb{G}_m/\mathbb{F}_q; \mathbb{Q}_\ell) = \mathbb{Q}_\ell$, for any generator $g \in \mu_n$, g^d acts trivially on $H_{et}^*(\mathcal{U}_{n_1, n_2}/\mathbb{F}_q; \mathbb{Q}_\ell)$; therefore $H_{et}^*(\mathcal{U}_{n_1, n_2}/\mathbb{F}_q; \mathbb{Q}_\ell)_m = 0$ for $m \in \mathcal{O}_a$ unless $a \mid d$.

To conclude the proof of the lemma, we now observe that $\mathcal{R}_n - Y_{n,a}$ admits an étale cover by the collection $\{\mathcal{U}_{n_1, n_2} \mid n = n_1 + n_2 \text{ with } a \nmid n_1, n_2\}$. Using what we just proved as the base case, we now conclude the lemma by induction on the number of étale neighborhoods \mathcal{U}_{n_1, n_2} , using the Mayer-Vietoris sequence in étale cohomology (see e.g. [AM69, Corollary 8.15]). \square

We apply the long exact sequence of a pair in étale cohomology; see Corollary 16.2 of [Mil13], and then we take the direct sum of the m -isotypic components for $m \in \mathcal{O}_a$. In Milne's notation, setting $c = n - a$; $Z = Y_{n,a}$; $X = \mathcal{R}_n$; $U = \mathcal{R}_n - Y_{n,a}$, for any $0 \leq j \leq 2(n - a) - 2$, we obtain that

$$H_{et}^j(\mathcal{R}_n/\mathbb{F}_q)_a \cong H_{et}^j((\mathcal{R}_n - Y_{n,a})/\mathbb{F}_q)_a$$

and, above degree $2(n - a) - 2$, we have a long exact sequence

$$\begin{aligned} 0 &\longrightarrow H_{et}^{2(n-a)-1}(\mathcal{R}_n/\mathbb{F}_q)_a \longrightarrow H_{et}^{2(n-a)-1}((\mathcal{R}_n - Y_{n,a})/\mathbb{F}_q)_a \longrightarrow \\ &H_{et}^0(Y_{n,a}/\mathbb{F}_q)_a \otimes \mathbb{Q}_\ell(a - n) \longrightarrow H_{et}^{2(n-a)}(\mathcal{R}_n/\mathbb{F}_q)_a \longrightarrow H_{et}^{2(n-a)}((\mathcal{R}_n - Y_{n,a})/\mathbb{F}_q)_a \cdots \\ \cdots &H_{et}^{j-2(n-a)}(Y_{n,a}/\mathbb{F}_q)_a \otimes \mathbb{Q}_\ell(a - n) \longrightarrow H_{et}^j(\mathcal{R}_n/\mathbb{F}_q)_a \longrightarrow H_{et}^j((\mathcal{R}_n - Y_{n,a})/\mathbb{F}_q)_a \cdots \end{aligned}$$

Lemma 2.4 gives that the terms $H_{et}^i((\mathcal{R}_n - Y_{n,a})/\mathbb{F}_q)_a$ vanish. It follows that $H_{et}^j(\mathcal{R}_n/\mathbb{F}_q)_a = 0$ for $j \leq 2(n - a) - 1$, and that, for $j \geq 2(n - a)$,

$$H_{et}^j(\mathcal{R}_n/\mathbb{F}_q)_a \cong H_{et}^{j-2(n-a)}(Y_{n,a}/\mathbb{F}_q)_a \otimes \mathbb{Q}_\ell(a - n) \quad (2.10)$$

We therefore have

$$H_{et}^j(\mathcal{R}_{n/\overline{\mathbb{F}}_q})_a \cong H_{et}^{j-2(n-a)}(Y_{n,a/\overline{\mathbb{F}}_q})_a \otimes \mathbb{Q}_\ell(a-n) \quad \text{by (2.10)}$$

$$\cong (\text{Ind}_{\mu_a}^{\mu_n} H_{et}^{j-2(n-a)}(\mathcal{R}_{a/\overline{\mathbb{F}}_q})_a \otimes \mathbb{Q}_\ell(a-n)) \quad \text{by (2.9)}$$

$$\cong \text{Hom}_{\mu_n}(\bigoplus_{m \in \mathcal{O}_a} V_m, \text{Ind}_{\mu_a}^{\mu_n} H_{et}^{j-2(n-a)}(\mathcal{R}_{a/\overline{\mathbb{F}}_q}) \otimes \mathbb{Q}_\ell(a-n))$$

where V_m is the m -isotypic irrep

$$\cong \bigoplus_{m \in \mathcal{O}_a} \text{Hom}_{\mu_a}(\text{Res}_{\mu_a}^{\mu_n} V_m, H_{et}^{j-2(n-a)}(\mathcal{R}_{a/\overline{\mathbb{F}}_q}) \otimes \mathbb{Q}_\ell(a-n))$$

by Frobenius Reciprocity

$$\cong \bigoplus_{m \in \mathcal{O}_a} \text{Hom}_{\mu_a}(V_0, H_{et}^{j-2(n-a)}(\mathcal{R}_{a/\overline{\mathbb{F}}_q}) \otimes \mathbb{Q}_\ell(a-n))$$

since a divides m

$$\cong \bigoplus_{m \in \mathcal{O}_a} H_{et}^{j-2(n-a)}(\mathcal{R}_{a/\overline{\mathbb{F}}_q})^{\mu_a} \otimes \mathbb{Q}_\ell(a-n)$$

which is, as we have shown above,

$$\cong \begin{cases} (\bigoplus_{m \in \mathcal{O}_a} \overline{\mathbb{Q}}_\ell) \otimes \mathbb{Q}_\ell(a-n) & j - 2(n-a) = 0 \\ 0 & j \neq 0 \end{cases}$$

as claimed above.

Step 4 (The permutation action of \mathbf{Frob}_q): We complete our analysis of $H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q})_a$ as a Frob_q -module. The analysis of Step 3 shows that Frob_q acts by q^{n-a} times the action of Frob_q on the factor $\bigoplus_{m \in \mathcal{O}_a} \overline{\mathbb{Q}}_\ell$ of $H_{et}^{2(n-a)}(\mathcal{R}_n)_a$. We claim that this is given by a permutation action

$$\text{Frob}_q: (\overline{\mathbb{Q}}_\ell)_m \xrightarrow{1} (\overline{\mathbb{Q}}_\ell)_{qm \bmod n}, \quad (2.11)$$

where $(\overline{\mathbb{Q}}_\ell)_m$ denotes the summand corresponding to $m \in \mathcal{O}_a$.

Granting this, we conclude that the variety \mathcal{R}_n is pure. Further, we see that \mathcal{R}_n is of Tate type if and only if $q \equiv 1 \pmod n$.

To prove the claim, recall that the actions of μ_n and of Frob_q on $H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q})$ do not commute. This is because Frob_q acts on $\mu_n(\overline{\mathbb{F}}_q)$ as an automorphism. For any vector $v \in H_{et}^i(\mathcal{R}_{n/\overline{\mathbb{F}}_q})$ and any $\sigma \in \mu_n$:

$$\text{Frob}_q(\sigma \cdot v) = \text{Frob}_q(\sigma) \cdot \text{Frob}_q(v). \quad (2.12)$$

To be more explicit, let λ be any primitive n^{th} root of 1 in $\overline{\mathbb{F}}_q$. Then we can write each element of μ_n as λ^j for some $0 \leq j \leq n-1$. The action of Frob_q on μ_n is given by $\text{Frob}_q(\lambda^j) = \lambda^{jq}$ where jq is taken mod n , and so

$$\text{Frob}_q(\lambda^j \cdot v) = \lambda^{aj} \cdot \text{Frob}_q(v) \quad (2.13)$$

for any $0 \leq j \leq n-1$. This, combined with Step 3 and Proposition 2.2 above prove the claim.

Step 5 (Computing the trace of \mathbf{Frob}_q): To conclude the proof we must compute the trace of Frob_q , which by the analysis above equals q^{n-a} times the number of Frob_q -fixed vectors. The permutation action given in Equation (2.11) has a fixed vector precisely when $(q-1)m \equiv 0 \pmod n$,

i.e. when $(q-1)m/a \equiv 0 \pmod{n/a}$. If $q \equiv 1 \pmod{n/a}$ then this equation has no solutions since by assumption $(m, n) = a$. It follows in this case that Frob_q acts with trace 0. If $q \not\equiv 1 \pmod{n/a}$ then this equation has $\phi(n/a)$ solutions, where ϕ is Euler's totient function. In particular, in this case Frob_q acts on $H_{et}^{2(n-a)}(\mathcal{R}_n; \mathbb{Q}_\ell)$ by multiplication by q^{n-a} .

3. Moduli space of magnetic monopoles

The varieties \mathcal{R}_n are closely related to some moduli spaces studied in physics, and the results above can be used to deduce arithmetic properties of these spaces, as we now explain.

The moduli space \mathcal{M}_n of $\text{SU}(2)$ monopoles of charge n in \mathbb{R}^3 , and the associated moduli space X_n of strongly centered monopoles, have a rich geometric and topological structure. These complex algebraic varieties have been studied both by physicists and mathematicians; see [SS96] and the references therein.

Let \mathcal{M}_n be the *moduli space of based $\text{SU}(2)$ monopoles in \mathbb{R}^3 of charge n* . Elements of \mathcal{M}_n are pairs (A, Φ) , where A is a smooth connection on the trivial $\text{SU}(2)$ bundle $E \rightarrow \mathbb{R}^3$, and Φ is a smooth section of the vector bundle associated to E via the adjoint representation. The pair (A, Φ) is a monopole if it satisfies two conditions. First, it must give a solution to the *Bogomolnyi equation*

$$*F_A = D_A\Phi$$

where $*$ is the Hodge star operator, D_A is the covariant derivative operator defined by A , and F_A is the curvature of A . Second, (A, Φ) must satisfy a regularity and boundary condition. See, e.g. Chapter 1 of [AH88] for details. These spaces connect to the present paper because of a different description of \mathcal{M}_n , due to Donaldson.

As explained for example by Manton and Murray, there are many ways of describing \mathcal{M}_n , each of which leads to the moduli space of degree n rational maps $\mathbb{P}^1 \rightarrow \mathbb{P}^1$. This is summarized in [MM94] by the following diagram:

$$\begin{array}{ccc}
 \text{Monopoles} & \longleftrightarrow & \text{Holomorphic bundles} \\
 \searrow & & \swarrow \\
 \updownarrow & \text{Rational Maps} & \updownarrow \\
 \nearrow & & \nwarrow \\
 \text{Nahm data} & \longleftrightarrow & \text{Spectral curves}
 \end{array}$$

The moduli space \mathcal{M}_n is a $2n$ -dimensional complex manifold. Donaldson [Don84] proved that there is a diffeomorphism of \mathcal{M}_n with the moduli space of degree n rational maps $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ that send ∞ to 0:

$$\mathcal{M}_n \cong \left\{ \frac{\phi}{\psi} = \frac{a_{n-1}z^{n-1} + \cdots + a_1z + a_0}{z^n + b_{n-1}z^{n-1} + \cdots + b_1z + b_0} : \phi, \psi \in \mathbb{C}[z] \text{ have no common root} \right\} \quad (3.1)$$

Thus the diffeomorphism (3.1) endows \mathcal{M}_n with the structure of a smooth, complex-algebraic variety of (complex) dimension $2n$. There is a subvariety X_n of \mathcal{M}_n , called the *reduced moduli space* of $\text{SU}(2)$ monopoles of charge n in \mathbb{R}^3 , or the *moduli space of strongly centered monopoles*,

given by

$$X_n \cong \left\{ \frac{\phi}{\psi} \in \mathcal{M}_n : b_{n-1} = 0 \text{ and } \mathcal{R}(\phi, \psi) = 1 \right\}$$

The algebraic variety X_n is a smooth hypersurface in \mathbb{C}^{2n-1} . It admits an action by algebraic automorphisms, of the cyclic group μ_n of n^{th} roots of unity. Segal–Selby [SS96] computed the isotypic components under this action of the rational singular cohomology groups $H^*(X_n; \mathbb{Q})$.

When n is invertible in \mathbb{F}_q there is a Zariski-locally trivial fibration $\mathbb{G}_a \rightarrow \mathcal{R}_1 \rightarrow X_n$. This together with Theorem 1.4 implies the following.

COROLLARY 3.1. *With notation as above, when $\gcd(q, n) = 1$,*

$$H_{\text{et}}^*(X_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell) \cong H_{\text{et}}^*(\mathcal{R}_n/\overline{\mathbb{F}}_q; \mathbb{Q}_\ell)$$

as given in Theorem 1.1 and

$$|X_n(\mathbb{F}_q)| = |\mathcal{R}_n(\mathbb{F}_q)|/q$$

as given in Corollary 1.4.

Appendix A. An elementary count of the cardinality of $\mathcal{R}_n(\mathbb{F}_q)$ (by C. Cazanave)

We present a more direct approach to count the cardinality of the set $\mathcal{R}_n(\mathbb{F}_q)$. It is elementary (no use of étale cohomology is needed) and we obtain a formula valid for all values of q and n .

Our main tool is the “addition law” \oplus of pointed rational functions that was introduced in [Caz12, Proposition 3.1]. For sake of completeness, we first recall it briefly.

Let \mathcal{F}_n denote the scheme of degree n rational functions which send ∞ to ∞ .⁶ Given two degrees $n_1, n_2 \geq 0$, we define a map

$$\oplus : \mathcal{F}_{n_1} \times \mathcal{F}_{n_2} \rightarrow \mathcal{F}_{n_1+n_2}$$

as follows. Two rational functions $\frac{A_i}{B_i} \in \mathcal{F}_{n_i}$, for $i = 1, 2$, *uniquely* define two pairs (U_i, V_i) of polynomials with $\deg U_i \leq n_i - 2$ and $\deg V_i \leq n_i - 1$ and satisfying Bzout identities $A_i U_i + B_i V_i = 1$ (this is true over any ring because A_i is monic). Define polynomials A_3, B_3, U_3 and V_3 as:

$$\begin{bmatrix} A_3 & -V_3 \\ B_3 & U_3 \end{bmatrix} := \begin{bmatrix} A_1 & -V_1 \\ B_1 & U_1 \end{bmatrix} \cdot \begin{bmatrix} A_2 & -V_2 \\ B_2 & U_2 \end{bmatrix}.$$

One easily checks that $\frac{A_3}{B_3}$ is in $\mathcal{F}_{n_1+n_2}$. Over a field, this addition law is closely related to continued fraction expansion of rational functions. This gives:

LEMMA A.1. *Let k be any field and let $\frac{A}{B}$ be a element of $\mathcal{F}_n(k)$. Then there exists a unique family of monic polynomials P_1, \dots, P_r and a unique family of scalars $a_1, \dots, a_r \in k^\times$ such that*

$$\frac{A}{B} = \frac{P_1}{a_1} \oplus \dots \oplus \frac{P_r}{a_r}.$$

Moreover:

- (i) Let n_i denote the degrees of the P_i . Then $n = n_1 + \dots + n_r$.
- (ii) There exists a sign $\varepsilon(n) = \pm 1$ (depending only on the n_i 's) such that

$$\mathcal{R}(A, B) = \varepsilon(n) a_1^{n_1} \dots a_r^{n_r}.$$

⁶Note that \mathcal{F}_n is isomorphic to \mathcal{M}_n under the Möbius transformation $\frac{A(z)}{B(z)} \mapsto \frac{B(z)-A(z)}{B(z)}$.

Proof. The decomposition of $\frac{A}{B}$ as a \oplus -sum of polynomials is explained in [Caz12, Example 3.3]. The formula expressing the resultant $\mathcal{R}(A, B)$ in terms of the resultants $\mathcal{R}(P_i, a_i)$ can be seen by induction by noting that if P is a monic polynomial of degree d , $a \in k^\times$ and $\frac{A}{B} \in \mathcal{F}_n(k)$ one has

$$\mathcal{R}\left(\frac{P}{a} \oplus \frac{A}{B}\right) = (-1)^{nd} a^d \mathcal{R}(A, B).$$

□

Remark A.2. The precise expression of the sign ε will not be needed in the sequel. It will be enough note that when all the n_i are even the sign $\varepsilon(\underline{n})$ is equal to 1.

Over a field k , specifying a pointed degree n rational function $f = \frac{A}{B}$ such that $\mathcal{R}(A, B) = 1$ is thus equivalent to specifying:

- an integer $1 \leq r \leq n$
- an ordered decomposition $n = n_1 + \cdots + n_r$ (with integers $n_i \geq 1$)
- monic polynomials P_1, \dots, P_r of degrees n_1, \dots, n_r
- units $a_1, \dots, a_r \in k^\times$ such that $a_1^{n_1} \cdots a_r^{n_r} \varepsilon(\underline{n}) = 1$.

We now specialize to $k = \mathbb{F}_q$. For $x \in \mathbb{F}_q^\times$, let us denote by $\mathcal{C}(\underline{n}, x)$ the cardinality of the finite set $\{(a_1, \dots, a_r) \in (\mathbb{F}_q^\times)^r, a_1^{n_1} \cdots a_r^{n_r} = x\}$. We therefore have:

$$|\mathcal{R}_n(\mathbb{F}_q)| = \sum_{r=1}^n \sum_{n=n_1+\dots+n_r} q^n \mathcal{C}(\underline{n}, \varepsilon(\underline{n})).$$

(The factor $q^n = q^{n_1+\dots+n_r}$ counts the choices for the polynomials P_i .)

The following lemma is useful:

LEMMA A.3. *Let $\underline{n} = \{n_1, \dots, n_r\}$ as above. Then*

- (i) *One has $\mathcal{C}(\underline{n}, 1) = (q-1)^{r-1} \gcd(q-1, n_1, \dots, n_r)$.*
- (ii) *If one of the n_i is odd, the one has the equality $\mathcal{C}(\underline{n}, 1) = \mathcal{C}(\underline{n}, -1)$.*

Proof. (i) Recall that the group of units \mathbf{F}_q^\times is cyclic of order $q-1$. Therefore, $\mathcal{C}(\underline{n}, 1)$ is also the number of solutions $(x_1, \dots, x_r) \in (\mathbb{Z}/(q-1)\mathbb{Z})^r$ of the linear equation

$$n_1 x_1 + \cdots + n_r x_r = 0.$$

To count the number solutions of this equation, one can use the invertible changes of variables dictated by the Euclidean algorithm to reduce it to the equivalent equation $\gcd(n_1, \dots, n_r) x_1 = 0$.

- (ii) Suppose n_i is an odd integer. Then one has an explicit bijection between the two sets in question given by $a_i \mapsto -a_i$.

□

Combining Remark A.2 and Lemma A.3, we get rid of the signs $\varepsilon(\underline{n})$ to obtain:

$$|\mathcal{R}_n(\mathbb{F}_q)| = q^n \sum_{r=1}^n \sum_{n=n_1+\dots+n_r} (q-1)^{r-1} \gcd(q-1, \underline{n}). \quad (\text{A.1})$$

For a fixed r , all decompositions \underline{n} with same gcd contribute equally, so we regroup these as follows.

NOTATION A.4. Let $n, r \geq 1$ be two integers.

- Let $\pi_r(n)$ be the number of decompositions of n of length r . One has $\pi_r(n) = \binom{n-1}{r-1}$.
- For d a divisor of n , let $\pi_r(n, d)$ be the number of length r decompositions of n with gcd equal to d . One has $\pi_r(n, d) = \pi_r(\frac{n}{d}, 1)$.

From the identity $\pi_r(n) = \sum_{d|n} \pi_r(n, d) = \sum_{d|n} \pi_r(\frac{n}{d}, 1)$, the Mbius inversion formula gives $\pi_r(\cdot, 1) = \mu \star \binom{\cdot-1}{r-1}$, where μ is the Mbius function and \star denotes the Dirichlet convolution product. Inserting this into equation (A.1) leads to :

$$|\mathcal{R}_n(\mathbb{F}_q)| = q^n \left[\mu \star \left(\sum_{r=1}^n q^{r-1} \binom{\cdot-1}{r-1} \right) \star \gcd(\cdot, q-1) \right] (n) = q^n [\mu \star q^{-1} \star \gcd(\cdot, q-1)] (n).$$

We have thus proved the following theorem.

THEOREM A.5. *Let n be any positive integer and let q be any prime power. Then*

$$|\mathcal{R}_n(\mathbb{F}_q)| = q^n \left(\sum_{abc=n} \mu(a) q^{b-1} \gcd(c, q-1) \right).$$

We end up briefly showing that the point count of Theorem A.5 coincides with that of Corollary 1.4. The two formulas are similar. The only point is to check the following lemma.

LEMMA A.6. *Let $m \geq 1$ be an integer and let $\delta_m : \mathbb{N} \rightarrow \mathbb{N}$ be the function*

$$\delta_m : n \mapsto \begin{cases} 1 & \text{if } n | m \\ 0 & \text{otherwise.} \end{cases}$$

Then we have the identity $\delta_m \phi = \mu \star \gcd(\cdot, m)$.

Proof. Let $n \geq 1$ be an integer. We prove that these two functions agree on n . We distinguish cases according to whether n divides m or not.

In the first case, the function $\gcd(\cdot, m)$ coincides with the identity and the equality follows from $\phi = \mu \star \text{id}$.

In the second case, we have to show that:

$$\sum_{d|n} \mu(d) \gcd(d, m) = 0.$$

Replacing m by $\gcd(n, m)$ does not change the value, so one can assume that m divides (strictly) n . Let p be a prime dividing $\frac{n}{m}$. We conclude noting that:

- in the above sum, only the divisors without square factor have a non-zero contribution;
- the remaining divisors can be split into those divisible by p and those not. These two sets have opposite contributions in the sum.

□

Remark A.7. Given $x \in \mathbb{F}_q^\times$, we could in the same way write a formula for the number of pointed degree n rational functions $\frac{A}{B} \in \mathcal{F}_n(\mathbb{F}_q)$ with resultant $\mathcal{R}(A, B) = x$. This number equals

$$q^n \left(\sum_{abc=n} \mu(a) q^{b-1} g(c) \right)$$

where o is the order of x in \mathbb{F}_q^\times and $g : \mathbb{N} \rightarrow \mathbb{N}$ is the function

$$g : n \mapsto \begin{cases} \gcd(n, q-1) & \text{if } \gcd(n, q-1) \text{ divides } \frac{q-1}{o} \\ 0 & \text{otherwise.} \end{cases}$$

However, it is interesting to note that, as a function of x , this cardinal always reaches its maximum for $x = 1$. This is because the functions $\mathcal{C}(\underline{n}, r)$ also reach their maximum at $x = 1$. The reason comes from linear algebra: a homogeneous linear equation $a_1x_1 + \dots + a_rx_r = 0$ has always at least as many solutions as the corresponding inhomogeneous one $a_1x_1 + \dots + a_rx_r = k$.

REFERENCES

- AH88 M. Atiyah and N. Hitchin, *The geometry and dynamics of magnetic monopoles*, Princeton University Press, 1988.
- AM69 M. Artin and B. Mazur, *Étale homotopy*, Springer, Lecture Notes in Mathematics **100** (1969).
- Caz12 Christophe Cazanave, *Algebraic homotopy classes of rational functions*, Ann. Sci. Éc. Norm. Supér. (4) **45** (2012), no. 4, 511–534 (2013). MR 3059240
- Don84 S. Donaldson, *Nahm’s equations and the classification of monopoles*, Comm. Math. Phys. **96** (1984), no. 3, 387–407.
- FW16 B. Farb and J. Wolfson, *Topology and arithmetic of resultants, I*, New York Journal of Mathematics **22** (2016), pp. 801–821.
- Gro58 A. Grothendieck, *Torsion homologique et sections rationnelles*, Séminaire Claude Chevalley, tome **3** (1958), no. 5, 1–29.
- Mil13 J. Milne, *Lectures on étale cohomology*, Version 2.21, March 22 2013.
- MM94 N. Manton and M. Murray, *Symmetric monopoles*, <http://arxiv.org/abs/hep-th/9407102v1>, 1994.
- SS96 G. Segal and A. Selby, *The cohomology of the space of magnetic monopoles*, Comm. Math. Phys. **177** (1996), no. 3, 775–787.
- vdW50 B. L. van der Waerden, *Modern Algebra*, Ungar (1950).

Benson Farb farb@math.uchicago.edu
 Dept. of Mathematics, University of Chicago, USA

Jesse Wolfson wolfson@math.uchicago.edu
 Dept. of Mathematics, University of Chicago, USA